



Modeling the output of a central pattern generator

Ronald Calabrese, *Emory University*

Paul Garcia, *Emory University*

Brian Norris, *Emory University*

Terrence Wright, *Emory University*

Journal Title: BMC Neuroscience

Volume: Volume 8, Number Suppl 2

Publisher: BioMed Central | 2007, Pages P166-P166

Type of Work: Article | Final Publisher PDF

Publisher DOI: 10.1186/1471-2202-8-S2-P166

Permanent URL: <https://pid.emory.edu/ark:/25593/pqpm>

Final published version: <http://dx.doi.org/10.1186/1471-2202-8-S2-P166>

Copyright information:

© 2007 Calabrese et al

This is an Open Access article distributed under the terms of the Creative Commons Attribution 4.0 International License (

<http://creativecommons.org/licenses/by/4.0/>), which permits distribution of derivative works, making multiple copies, distribution, public display, and publicly performance, provided the original work is properly cited. This license requires copyright and license notices be kept intact, credit be given to copyright holder and/or author.



Accessed December 7, 2021 10:16 PM EST

Poster presentation

Open Access

Modeling the output of a central pattern generator

Ronald L Calabrese*, Paul S García, Brian J Norris and Terrence M Wright

Address: Department of Biology, Emory University, Atlanta, Georgia, USA

Email: Ronald L Calabrese* - ronald.calabrese@emory.edu

* Corresponding author

from Sixteenth Annual Computational Neuroscience Meeting: CNS*2007
Toronto, Canada. 7–12 July 2007

Published: 6 July 2007

BMC Neuroscience 2007, 8(Suppl 2):P166 doi:10.1186/1471-2202-8-S2-P166

© 2007 Calabrese et al; licensee BioMed Central Ltd.

Experimental analysis in our lab has provided a quantitative description of the spatiotemporal pattern of inhibitory synaptic input from the heartbeat central pattern generator (CPG) to segmental motor neurons that drive heartbeat in the medicinal leech. To begin the process of elucidating the relative roles of this pattern of input and motor neuron intrinsic properties and electrical coupling in the elaboration of the heartbeat fictive motor pattern, we constructed a conductance-based ensemble model of all the segmental heart motor neurons and their known synaptic inputs. Our focus was intersegmental and side-to-side coordination of the asymmetric motor pattern: motor neurons on one side fire nearly in synchrony (synchronous coordination), while on the other they fire in a rear-to-front progression (peristaltic coordination). The model reproduces the general trends of the two intersegmental phase relations among motor neurons, but the match with the living system is quantitatively poor, particularly for the peristaltic coordination mode where the phase progression among the segmental motor neurons in the model is only half that observed in the living system. Thus the realistic inputs (experimentally determined) do not produce similarly realistic output in our model.

Modeling experiments, indicate that the most important determinant of the intersegmental and side-to-side phase relations among the heart motor neurons in the model was the spatiotemporal pattern of synaptic inputs, yet phasing was influenced by electrical coupling between the motor neurons in each segment, intersegmental conduction delays in the premotor interneurons, intra-burst syn-

aptic plasticity, and intrinsic membrane currents of the motor neurons.

Understanding the shortcomings of the model required that we establish experimentally the precise timing of motor neuron activity in each segment with respect to CPG activity. This analysis show quantitatively how motor neurons in the model fail to fire at the appropriate time with respect to their synaptic inputs and suggest that the intrinsic properties of the model motor neuron are simplistic.