



EMORY
LIBRARIES &
INFORMATION
TECHNOLOGY

OpenEmory

Noscapine Induced Apoptosis via Downregulation of Survivin in Human Neuroblastoma Cells Having Wild Type or Null p53

Shiwang Li, *Huazhong University of Science & Technology*
Jing He, *Huazhong University of Science & Technology*
Shuai Li, *Huazhong University of Science & Technology*
Guoqing Cao, *Huazhong University of Science & Technology*
Shaotao Tang, *Huazhong University of Science & Technology*
Qiangsong Tong, *Huazhong University of Science & Technology*
[Harish Joshi](#), *Emory University*

Journal Title: PLoS ONE

Volume: Volume 7, Number 7

Publisher: Public Library of Science | 2012-07-26, Pages e40076-e40076

Type of Work: Article | Final Publisher PDF

Publisher DOI: 10.1371/journal.pone.0040076

Permanent URL: <https://pid.emory.edu/ark:/25593/s5gfc>

Final published version: <http://dx.doi.org/10.1371/journal.pone.0040076>

Copyright information:

© 2012 Li et al.

This is an Open Access work distributed under the terms of the Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/>).

Accessed October 18, 2019 4:27 PM EDT

Noscapine Induced Apoptosis via Downregulation of Survivin in Human Neuroblastoma Cells Having Wild Type or Null p53

Shiwang Li¹, Jing He^{2*}, Shuai Li¹, Guoqing Cao¹, Shaotao Tang¹, Qiangsong Tong¹, Harish C. Joshi³

1 Department of Pediatric Surgery, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China, **2** Institute of Hematology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China, **3** Department of Cell Biology, Emory University, Atlanta, Georgia, United States of America

Abstract

Neuroblastoma is the most common extracranial solid tumor of childhood. It accounts for 15% of pediatric cancer deaths. Chemotherapy is the mainstay of treatment in children with advanced neuroblastoma. Noscapine, a nontoxic natural compound, can trigger apoptosis in many cancer types. We now show that p53 is dispensable for Noscapine-induced cell death in neuroblastoma cell lines, proapoptotic response to this promising chemopreventive agent is mediated by suppression of survivin protein expression. The Noscapine treatment increased levels of total and Ser¹⁵-phosphorylated p53 protein in SK-SY5Y cells, but the proapoptotic response to this agent was maintained even after knockdown of the p53 protein level. Exposure of SK-SY5Y and LA1-5S cells to Noscapine resulted in a marked decrease in protein and mRNA level of survivin as early as 12 hours after treatment. Ectopic expression of survivin conferred statistically significant protection against Noscapine-mediated cytoplasmic histone-associated apoptotic DNA fragmentation. Also, the Noscapine-induced apoptosis was modestly but statistically significantly augmented by RNA interference of survivin in both cell lines. Furthermore, Noscapine-induced apoptotic cell death was associated with activation of caspase-3 and cleavage of PARP. In conclusion, the present study provides novel insight into the molecular circuitry of Noscapine-induced apoptosis to indicate suppression of survivin expression as a critical mediator of this process.

Citation: Li S, He J, Li S, Cao G, Tang S, et al. (2012) Noscapine Induced Apoptosis via Downregulation of Survivin in Human Neuroblastoma Cells Having Wild Type or Null p53. PLoS ONE 7(7): e40076. doi:10.1371/journal.pone.0040076

Editor: Wael El-Rifai, Vanderbilt University Medical Center, United States of America

Received: November 17, 2011; **Accepted:** June 5, 2012; **Published:** July 26, 2012

Copyright: © 2012 Li et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: Grant support: the National Natural Science Foundation of China (NSFC) grants 81070429 (Jing He), 30600278 and 30772359 (Qiangsong Tong). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: jing937@hotmail.com

Introduction

Neuroblastoma is a pediatric cancer of the developing sympathetic nervous system that most often affects young children. It remains an important pediatric problem because it accounts for approximately 15% of childhood cancer mortality [1]. Although improvement of multimodal therapy has been achieved by the progress of treatment, prognosis of this cancer, especially high-risk neuroblastoma, remains poor. On account of poor treatment outcome, new treatment strategies are constantly sought for neuroblastoma.

Among anticancer agents, antimicrotubules constitute one of the promising chemotherapeutic agents for treatment of different cancers. During cell division, α/β -tubulin polymerizes into dynamic structures called microtubules. Inhibitors of tubulin either target polymerization (vinca alkaloids and colchicine) or depolymerization (taxanes and epothilones). These two functional classes bind to different regions of the α/β -tubulin heterodimer and allosterically regulate tubulin oligomerization. Colchicine is deeply buried at the heterodimer interface and prevents a conformational change needed for polymerization [2], whereas taxanes bind in a shallow groove found on the β -subunit of tubulin in microtubules [3]. Small-molecule inhibitors of tubulin have been in clinical use since 1965. Many classes of tubulin inhibitors are

known, and it appears that the presence of multiple allosteric binding sites makes tubulin particularly amenable to inhibition by small molecules [4]. Clinical use of currently available antimicrotubular agents has been limited due to drug-resistance, prolonged i.v. infusion and associated adverse side effects [5,6]. Among antimicrotubule agents, noscapinoids constitute an emerging class of compounds receiving considerable attention for treating cancers due to oral available, improved patient compliance and minimal side effects compared to taxanes [7,8,9,10]. Noscapine (Nos) attenuates microtubule dynamics just enough to activate the mitotic checkpoints to stop cell cycle and do not alter the steady state monomer/polymer ratio of tubulin [11] and demonstrated promising in vitro and in vivo antitumor activity against variety of cancers including resistant type [11,12,13,14,15,16]. Based upon anticancer activity and non-toxic attributes, Noscapine is already in Phase I/II clinical trials.

We now show that p53 is dispensable for Noscapine-induced cell death, proapoptotic response to this promising chemopreventive agent is mediated by suppression of survivin protein expression. The Noscapine treatment increased levels of total and Ser¹⁵-phosphorylated p53 protein in NB cells, but the proapoptotic response to this agent was maintained even after knockdown of the p53 protein level. Exposure of SK-SY5Y and

LA1-5S cells to Noscapine resulted in a marked decrease in protein level of survivin as early as 12 hours after treatment. Ectopic expression of survivin conferred statistically significant protection against Noscapine-mediated cytoplasmic histone-associated apoptotic DNA fragmentation in both cell lines, while knock-down of endogenous survivin by survivin siRNA sensitizes both cells to Noscapine-induced apoptosis. Furthermore, Noscapine-induced apoptotic cell death in SK-SY5Y and LA1-5S cells was associated with activation of caspase-3 and cleavage of PARP. In conclusion, the present study provides novel insight into the molecular circuitry of Noscapine-induced apoptosis to indicate suppression of survivin expression as a critical mediator of this process.

Results and Discussion

Noscapinoids represent a new generation of anticancer agents that modulate microtubule dynamics but do not significantly alter the total polymer mass of tubulin. To evaluate the efficacy of Noscapine in neuroblastoma cells, we first examined the ability of Noscapine to inhibit cellular proliferation in a comprehensive panel of neuroblastoma cells with variable but well-characterized genotypes. The panel included SK-SY5Y, SH-EP1, SK-N-MC, SK-N-AS, LA1-55N, LA1-5S, NB1643, NB1691, SK-N-SH and IMR32 neuroblastoma cells. These 10 different neuroblastoma cell lines were treated with gradient concentrations of Noscapine and the extent of cell proliferation was measured by the SRB assay, which is based on the stoichiometric binding of SRB dye to all cellular protein components [17]. As shown in Figure 1A, Noscapine effectively suppressed cellular proliferation of neuroblastoma cells. The neuroblastoma cell lines included in the study had well-characterized p53 status, namely, wild type (represented by SK-SY5Y, SH-EP1, NB1643, NB1691), null (LA1-55N, LA1-5S) and mutant (such as SK-N-AS). Although Noscapine has been shown to induce a p53-dependent apoptosis in colon cancer cells [18], these data show that the half-maximal growth inhibitory concentrations of Noscapine did not correlate with the p53 status in neuroblastoma cells. For example, LA1-5S that lacks endogenous p53 showed similar sensitivity as SK-SY5Y (wild type p53) or SK-N-AS (mutant p53) cells. The IC₅₀ values for most neuroblastoma cell lines studied were in the range of 21–101 μ M. The relatively less-sensitive neuroblastoma cells included SK-N-SH (101.2 μ M), LA1-55N (IC₅₀ = ~81.4 μ M) and SK-N-MC (IC₅₀ = ~72.3 μ M). We chose SK-SY5Y (wild type p53) and LA1-5S (p53null) for further studies to gain insights into cellular and molecular mechanisms of Noscapine action. Phase-contrast microscopic analysis of cell morphology showed that while vehicle-treated SK-SY-5Y (Figure 1B first panel) and LA1-5S (Figure 1B second panel) cells proliferated normally, Noscapine treatment impaired their proliferation capacity. Cells first appeared rounded-up (12 hrs and 24 hrs treatment (Figure 2C), followed by a fragmented morphology (48 hrs treatment, Figure 1B, 2C).

To investigate the precise mechanism responsible for Noscapine-mediated antiproliferative effects, we examined the cell-cycle distribution profile of Nos-treated SK-SY5Y neuroblastoma cells over time. A flow-cytometric assay using the DNA intercalator dye, propidium iodide (PI), was utilized to monitor cell-cycle progression on the basis of status of DNA amounts. Figure 2Ai shows the time dependent effects of Noscapine on cell-cycle profile of SK-SY5Y cells in a three-dimensional representation. The x-axis shows amount of DNA depicting different phases of the cell-cycle. While 2N and 4N DNA complements represent G₀/G₁ and G₂/M cell populations respectively, S phase is characterized by variable DNA (between 2N and 4N) and sub-G₁ population is

usually indicative of degraded DNA, a hallmark of apoptosis. The y-axis represents the number of cells containing that amount of DNA and the z-axis shows the time of drug-exposure. 0 hr depicts cell-cycle profile of control cells. As we go along in time, Noscapine treatment caused a significant inhibition of cell-cycle progression in SK-SY5Y cells resulting in an accumulation of cells in the G₂/M phase compared to control cells (Figure 2Ai). The G₂/M population achieved a maximum at 12 hrs (~67%) and was still about ~48% at 48 hrs of Noscapine exposure (Figure 2Aii). This increased population of cells with 4N DNA perhaps correlated with concomitant losses from G₀/G₁ phases (Figure 2Ai). Following this, a disappearance of the G₂/M population and an emergence of a characteristic hypodiploid DNA content peak (sub-G₁) was observed beginning 12 hrs, indicative of apoptotic cells (Figure 2Ai). The sub-G₁ population at 12 hrs (~15%) increased to ~61% at 48 hrs post-treatment (Figure 2Ai).

Fluorescent-activated cell sorting (FACS) analyses of DNA content showed a decline in G₁ and a concomitant rise in G₂/M. It also showed a distinct shift of cells to a sub-G₁ DNA content of <2 N at 48 h Noscapine-treatment suggesting DNA-degradation associated with apoptosis (Figure 2Aii). Careful confocal microscopic examination of the MT arrays and the chromosomes in drug-treated cells revealed a typical mitotic arrest with bipolar prometaphase spindles with tightly condensed chromosomes. In contrast, highly disorganized, often multipolar mitoses were visible with less condensed chromosomes and appeared to die 48-hours post treatment (Figure 2C).

To determine the extent of the apoptotic onset caused by Noscapine in SK-SY5Y cells, we analyzed the early apoptotic cells that externalize the normally internal membrane lipid, phosphatidylserine (PS), by binding to a fluorescently conjugated PS-binding protein, Annexin V-Alexa Fluor488. These cells can be distinguished from the late apoptotic cells that also allow the DNA dye, propidium iodide (PI) to penetrate the cell membranes allowing intracellular DNA to bind it. As shown in Figure 2B, 70.1% cells were in early and late stages of apoptosis after a 48 hour Noscapine treatment (50 μ M) as compare to a modest 20.2% at vehicle control treatment (Figure 2Bi, Bii). The same results were also confirmed in LA1-5S cell lines (data not shown). We conclude that Noscapine caused programmed cell death both in SK-SY5Y and LA1-5S neuroblastoma cell lines.

We used the same cell line to test whether Noscapine-induced apoptosis was influenced by p53, which is a well-accepted facilitator of apoptotic cell death by different stimuli [19]. As can be seen in Figure 3A, Noscapine exposure caused a time-dependent increase in the levels of total as well as Ser¹⁵-phosphorylated p53. The Ser¹⁵ phosphorylation of p53 has been implicated in apoptosis [19]. Next, we used siRNA technology to directly test possible involvement of p53 in regulation of Noscapine-induced apoptosis. Transient transfection of SK-SY5Y cells with a p53-targeted siRNA resulted in complete silencing of the p53 protein expression (Figure 3B). Moreover, the Noscapine-mediated induction of p53 protein was abolished in SK-SY5Y cells transfected with the p53-specific siRNA (Figure 3B). However, enrichment of cytoplasmic histone-associated apoptotic DNA fragmentation resulting from Noscapine exposure (50 μ mol/L for 24 h) over DMSO-treated control was comparable in SK-SY5Y cells transfected with the control nonspecific siRNA and p53-targeted siRNA (Figure 3C). Knock-down of p53 using p53 siRNA didn't alter the apoptotic response of SK-SY5Y cells to Noscapine treatment. Collectively, these results indicated that the Noscapine-induced apoptosis was not influenced by the p53 status at least in the SK-SY5Y cell line.




Figure 1. Noscapine suppresses proliferation of human neuroblastoma cells. **A.** A panel of Neuroblastoma cells (SK-SY5Y, SH-EP1, SK-N-MC, SK-N-AS, LA1-55N, LA1-5S, NB1643, NB1691, SK-N-SH and IMR32) were treated with gradient concentrations of Noscapine for 72 hrs and the IC₅₀ values were evaluated by the sulforhodamine B (SRB) *in vitro* cell proliferation assay. The values and error bars shown in the graph represent average and standard deviations, respectively, of three independent experiments ($p < 0.05$). **B.** Phase-contrast images (20 \times) of SK-SY5Y cells treated with 50 μ M Noscapine for 0, and 48 hrs. doi:10.1371/journal.pone.0040076.g001

Since knockdown of p53 had no influence on proapoptotic response to Noscapine in SK-SY5Y cells, IAP proteins, including cIAP1, XIAP, and survivin have emerged as critical regulators of apoptotic cell death [20–21], we raised the question of whether the Noscapine induced apoptosis in neuroblastoma cells was accompanied by alterations in expression of IAP family proteins. The IAPs play important roles in adaptive response to cellular stress, differentiation, motility, and immune response [22]. This family of proteins is characterized by the presence of baculovirus IAP repeat (BIR) domains [23]. The survivin protein expression was markedly decreased on treatment of SK-SY5Y and LA1-5S cells with Noscapine at 50 μ mol/L concentrations. The Noscapine-mediated decline in survivin protein level was evident as early as 12 hours after treatment, and this effect was maintained for the duration of the experiment (48 h after treatment) in both cell lines (Figure 4). The level of cIAP1 protein was increased on Noscapine treatment in both SK-SY5Y and LA1-5S cell lines, but this effect was not sustainable (Figure 4). For example, Noscapine-mediated induction of cIAP1 protein was nearly completely abolished at the 48-hour time point in the SK-SY5Y cell line. On the other hand, the Noscapine treatment did not have an appreciable effect on XIAP protein level both in SK-SY5Y and LA1-5S cell lines. We also test the level of survivin in p53 knockout experiment. As shown in figure 3B, survivin levels didn't have apparent changes whether p53 was knockout or not, but do have changes when treated with Noscapine. Collectively, these results indicated that

Noscapine treatment caused a sustained decrease in the level of survivin protein in both cell lines.

Survivin, a 17 kD protein that contains only a single BIR and no RING domain, is the smallest member of IAPs and stands out for its clear association with cancer [24,25]. Reactivation of the survivin gene occurs frequently during tumorigenesis and in most cases correlates with drugs resistance and shorter disease free and poor overall survival [26,27]. Synthesis and degradation of survivin are modulated in a cell cycle-dependent manner. Transcription increases during G₁ and reaches a peak at G₂/M, which is controlled by specific sequences in the promoter region involving Sp1 and cell cycle-dependent element binding sites [28]. One of the first characteristics of survivin to be described was its ability to bind to microtubules through the carboxy-terminal coiled-coil domain, and to assist in chromosomal segregation and cytokinesis during mitosis [29,30].

Although pioneering studies of Hoffman and colleagues [31], and Mirza and colleagues [32] identified survivin as one of the relatively few known genes that is actively repressed by wild type p53, we failed to identify this effectiveness in SK-SY5Y cell line which with wild type p53. After deletion of p53 with siRNA in SK-SY5Y cells, the survivin protein was not apparently changed (data not shown). To reveal the precise mechanism by which Noscapine treatment suppresses expression of survivin protein, we have had carefully examination of the survivin mRNA using rt-PCR and a significant down-regulation (of 35~70% in SK-SY5Y and 30~75% in LA1-5S) of survivin mRNA expression were observed at 12, 24 and 48 h post-




Figure 2. Noscapine perturbs cell-cycle progression of Sk-SY5Y cells and cause apoptosis. **A.** Effect of 25 μ M EM011 on cell-cycle progression of A549 cells over time shown in a three-dimensional disposition. X-axis, intensity of propidium iodide fluorescence, is indicative of the total DNA content of cells in various phases of the cell-cycle. Y-axis, number of cells detected for a given DNA content. Z-axis, time points (i.e., 0, 12, 24, and 48 hrs). Representative results of three independent experiments. **B.** Quantitative graphical representation of the percentage of G2/M and sub-G1 cell population. Points, average of three independent experiments; bars, SD ($p < 0.05$). **C.** Confocal immunofluorescence analyses of cells. Microtubules and DNA were visualized using fluorescent anti- α -tubulin antibody (green) and fluorescent DNA dye propidium iodide (red). SK-SY5Y cells begin to show mitotic arrest as early as 12 hours after drug treatment and begin to show disorganized arrest and chromatin decondensation or even death at 48 hours of treatment. doi:10.1371/journal.pone.0040076.g002

treatment ($P < 0.05$) (Figure 4C, 4D). Because Noscapine treatment exhibited the most striking effect on survivin protein as well as mRNA expression in both cell lines (Figure 4), we designed experiments to determine functional significance of these observations. After screening all the neuroblastoma cell lines in our lab, we found notable difference between the cell lines (figure 3D). We use another two cell lines (NB1691 and LA1-55N) that have least basal level of survivin for purpose of further over expression experiment. Transient transfection of NB1691 cells with a vector encoding for survivin resulted in ~ 1.8 -fold increase in its protein level compared with empty vector-transfected cells (Figure 5A). The Noscapine treatment (50 μ mol/L for 24 h) caused a decrease in the level of survivin protein both in empty vector-transfected NB1691 cells and in survivin-overexpressing cells (Figure 5A). In addition, overexpression of survivin conferred significant protection against

Noscapine-induced cytoplasmic histone-associated DNA fragmentation in NB1691 cells (Figure 5B). Statistically significant inhibition of Noscapine-induced cytoplasmic histone-associated DNA fragmentation by forced overexpression of survivin was also observed in the LA1-55N cell line (Figure 5C, 5D). These results indicated that survivin was a target of Noscapine-induced apoptosis in neuroblastoma cells, at least in NB1691 and LA1-55N cell lines.

If survivin is important to Noscapine-induced apoptosis, knock-down of endogenous survivin expression should render SK-SY5Y and LA1-55N cells to become more susceptible to Noscapine treatment. Thus, to determine whether downregulation of survivin contributes to Noscapine-induced apoptosis, we used a plasmid construct encoding survivin specific siRNA to selectively knock-down endogenous survivin in SK-SY5Y and LA1-55N cells. Exposure of nonspecific siRNA-transfected SK-SY5Y cells to




Figure 3. Noscapine cause p53 independent survivin downregulation. A, immunoblotting for total and phospho-(Ser¹⁵)-p53 using lysates from SK-SY5Y cells treated with DMSO (control) or Noscapine (50 μmol/L) for the indicated time periods. B, immunoblotting for p53 using lysates from SK-SY5Y cells transfected with a control nonspecific siRNA or a p53-targeted siRNA and treated for 24 h with DMSO (control) or 50 μmol/L Noscapine. C, cytoplasmic histone-associated apoptotic DNA fragmentation in SK-SY5Y cells transiently transfected with a control nonspecific siRNA or a p53-targeted siRNA and treated for 24 h with DMSO (control) or 50 μmol/L Noscapine. Columns, mean (n = 3); bars, SD. *, significantly different (P < 0.05) compared with nonspecific siRNA-transfected cells treated with DMSO by one-way ANOVA followed by Bonferroni's test. The results were consistent in three independent experiments, and representative data from one such experiment are shown. D, Immunoblotting for survivin using lysates from a series of NB cell lines.

doi:10.1371/journal.pone.0040076.g003

50 μmol/L Noscapine for 24 hours resulted in ~3-fold increase in survivin protein level compared with DMSO-treated control (Figure 6A). The level of survivin protein was decreased by >95% in SK-SY5Y cells transfected with a survivin-targeted siRNA (Figure 6A). Moreover, the Noscapine-mediated induction of survivin protein expression was nearly completely abolished in survivin siRNA transfected cells. As shown in Figure 6B, the cytoplasmic histone-associated DNA fragmentation enrichment resulting from Noscapine exposure over DMSO-treated control was modestly but statistically significantly greater in survivin silenced cells in comparison with nonspecific siRNA transfected SK-SY5Y cells (Figure 6B). On the other hand, knockdown of survivin protein level (Figure 6C) also have an appreciable effect on Noscapine-induced apoptosis in the LA1-5S cell line

(Figure 6D). We conclude that induction of survivin is marginally cytoprotective against Noscapine-mediated cell death in the SK-SY5Y and LA1-5S cells.

Upon cleavage by upstream proteases in an intracellular cascade, the activation of caspase-3 is considered as a hallmark of the apoptotic process. The levels of cleaved active subunits of executioner caspase-3 were evaluated by immunoblotting cell lysates following Noscapine treatment for 0, 12, 24, and 48 hrs. Noscapine caused a significant increase in activated caspase-3 following 48 hrs of Noscapine exposure (Figure 7). To confirm the involvement of caspase-3, the active form of the cysteine protease was monitored using a small conserved modified peptide substrate that becomes luminogenic upon cleavage. Noscapine treatment caused a time-dependent activation of caspase-3 in SK-SY5Y cells




Figure 4. Noscapine selectively cause survivin downregulation in SK-SY5Y and LA1-5S cells. Immunoblotting for survivin, cIAP1 and XIAP using lysates from SK-SY5Y (A) and LA1-5S (B) cells treated with DMSO (control) or Noscapine (50 $\mu\text{mol/L}$) for the indicated time periods. Immunoblotting for each protein was done at least twice using independently prepared lysates. Representative data from one such experiment are shown. Histograms represent the increase or decrease of survivin mRNA expression in SK-SY5Y (C) and LA1-5S (D) cells incubated with noscapine (50 μM) for 12, 24, and 48h. Measurements were performed in triplicate. Bars represent standard errors and asterisks designate statistically significant results ($P < 0.05$).

doi:10.1371/journal.pone.0040076.g004

(Figure 7). Next, we examined the activation mediated cleavage of caspase-3 substrate, poly(ADP-ribose) polymerase (PARP), which is a reliable marker of apoptosis. Utilizing their cysteine protease activity, caspases separate N-terminal DNA-binding domain of PARP from its C-terminal catalytic domain (89 kDa) [33]. A time-dependent increase in cleaved PARP was observed upon probing with a cleaved PARP specific antibody (Figure 7). Overall, these results show activation of caspase-3 and PARP cleavage suggesting that Noscapine induced apoptotic cell death in SK-SY5Y cells. Same results were also observed in LA1-5S cell line (data not shown).

In conclusion, we provide experimental evidence to show that the Noscapine-induced apoptosis in human neuroblastoma cells is independent of p53 but mediated by suppression of survivin protein level. The cell death program is mediated through downregulation of survivin, in that knock-down of survivin sensitized cells to undergo apoptosis whereas overexpression of survivin reduced Noscapine-induced apoptosis. The inhibition of cellular proliferation is perhaps due to induction of a robust transient mitotic arrest. This is followed by activating caspase machinery by the cleavage of downstream targets such as PARP. This study provides evidence for the potential usefulness of

Noscapine, a non-toxic microtubule-modulating agent, in chemotherapy of neuroblastoma.

Materials and Methods

Chemicals and Reagents

Noscapine, Sulforhodamine B (SRB), 4,6-diamidino-2-phenylindole (DAPI), propidium iodide (PI), RNase, bovine serum albumin (BSA), and the mouse monoclonal antibody against α -tubulin were from Sigma-Aldrich (St. Louis, MO, US). anti-p53 antibody was from Calbiochem; antibodies against cIAP1, phospho-p53 (Ser¹⁵), caspase-3, cleaved PARP were obtained from Cell Signaling (Beverly, MA); anti-survivin antibody was from Novus Biologicals; antibody against X-linked inhibitor of apoptosis (XIAP) was from BD Biosciences. Horseradish peroxidase-conjugated anti-rabbit and anti-mouse secondary antibodies were purchased from Sigma. Fluorescein-conjugated anti-mouse antibody was from Jackson ImmunoResearch, Inc. (West Grove, PA). RPMI 1640 and MEM were purchased from Cellgro. The p53- and survivin-targeted small interfering RNA (siRNA) were purchased from Santa Cruz Biotechnology. A nonspecific control siRNA was from Qiagen. FuGENE6 transfection reagent and a kit




Figure 5. Overexpression of survivin conferred protection Noscipine-induced apoptosis. Noscipine-induced apoptosis. Immunoblotting for survivin protein using lysates from NB1691(A) and LA1-55N (C) cells transiently transfected with empty pcDNA3.1 vector or pcDNA3.1 vector encoding for survivin and treated for 24 h with DMSO (control) or 50 μmol/L Noscipine. The numbers above the immunoreactive bands represent change in protein level relative to empty vector-transfected cells treated with DMSO (lane 1). Cytoplasmic histone-associated apoptotic DNA fragmentation in NB1691 (B) and LA1-55N (D) cells transiently transfected with empty pcDNA3.1 vector or vector encoding for survivin and treated for 24 h with DMSO (control) or 50 μmol/L Noscipine. Columns, mean (n=3); bars, SD. Significantly different (P<0.05) compared with empty vector-transfected cells treated with DMSO (a) and empty vector-transfected cells treated with Noscipine (b) by one-way ANOVA followed by Bonferroni's test. Comparable results were observed in three independent experiments. Representative data from one such experiment are shown. doi:10.1371/journal.pone.0040076.g005

for quantification of cytoplasmic histone-associated apoptotic DNA fragmentation were procured from Roche Applied Sciences.

Cell Lines and Plasmids

Human neuroblastoma cells (SK-SY5Y, SH-EP1, SK-N-MC, SK-N-AS, LA1-55N, LA1-5S, NB1643, NB1691, SK-N-SH and IMR32) were well established cell lines [34,35,36,37,38] and obtained from Dr. Muxiang Zhou's laboratory, Children oncology/hematology, Emory University. All cells were maintained in RPMI-1640 media supplemented with 10% fetal bovine serum (Invitrogen Carlsbad, CA) and 1% antibiotics. The plasmid vector with survivin siRNA sequence 5'-GGCTGGCTTCATC-CACTGCCC-3' was generated by cloning the synthesized oligonucleotide into pSilencer 2.1-U6Neo plasmid (Ambion Inc., Austin, TX). Control pSilencer 2.1-U6 Neo plasmid vector containing a scrambled siRNA sequence, 5'-ACTACCGTTGT-TATAGGTGT-3', was obtained from Ambion Inc. The survivin plasmid construct was cloned into the pcDNA3 vector and pcDNA3 vector alone served as control. All transfections were done using Lipofectamine 2000 following the manufacturers' instructions.

In vitro Cell Proliferation Assay

Neuroblastoma cells were seeded in 96-well plates at a density of 5×10^3 cells/well followed by next day treatment with increasing gradient concentrations of Noscipine ranging from 10 nm to 100 mM. After 48 hrs of drug treatment, cells were fixed with 50% trichloroacetic acid and stained with 0.4% sulforhodamine B (SRB) dissolved in 1% acetic acid. Cells were then washed with 1% acetic acid to remove the unbound dye. Essentially, the SRB assay measures cell density by quantitating colored SRB bound to cellular proteins fixed to the plates by trichloroacetate. The protein-bound dye was extracted with 10 mM Tris-base to determine the absorbance at 564 nm wavelength [17,39]. The percentage of cell survival as a function of drug concentration was then plotted to determine IC₅₀ values (drug concentration needed to prevent cell proliferation by 50%).

Cell-cycle Analysis

SK-SY5Y and LA1-5S cells were seeded in culture dishes and grown until ~70% confluence. The medium was then replaced with new medium containing either vehicle (0.01% DMSO) or 25 μM Noscipine for 6, 12, 18, 24, 36 and 48 hrs. After the incubation period, cells were centrifuged, washed twice with ice-




Figure 6. Knock-down endogenous survivin render SK-SY5Y and LA1-5S cells more susceptible to Noscapine. Immunoblotting for survivin using lysates from SK-SY5Y (A) and LA1-5S (C) cells transiently transfected with a control nonspecific siRNA or a survivin-specific siRNA and treated with DMSO (control) or 50 µmol/L Noscapine for 24 h. Cytoplasmic histone-associated DNA fragmentation in SK-SY5Y (B) and LA1-5S (D) cells transiently transfected with a control nonspecific siRNA or a survivin-specific siRNA and treated with DMSO (control) or 50 µmol/L Noscapine for 24 h. Columns, mean (n = 3); bars, SD. Significantly different ($P < 0.05$) compared with control nonspecific siRNA-transfected cells treated with DMSO (a) and control nonspecific siRNA-transfected cells treated with Noscapine (b) by one-way ANOVA followed by Bonferroni's test. Results were consistent in two experiments, and representative data from one such experiment are shown. doi:10.1371/journal.pone.0040076.g006




Figure 7. Noscapine induced apoptosis via activation of caspase-3 and PARP cleavage. Immunoblot analysis of SK-SY5Y and LA1-5S cells treated with noscapine for 0, 12, 24 and 48 hrs. After the indicated times, cells were lysed and total protein was extracted, separated by SDS-PAGE, electrotransferred onto polyvinylidene difluoride membrane, and subjected to immunoblotting with the indicated primary antibodies followed by incubation with horseradish peroxidase-conjugated secondary antibodies. β -actin was used as a loading control. doi:10.1371/journal.pone.0040076.g007

cold PBS, and fixed in 70% ethanol. Tubes containing the cell pellets were stored at 4°C for at least 24 hrs. Cells were then centrifuged at 100×g for 10 minutes, and the supernatant was discarded. Pellets were washed twice with PBS and stained with PI in the presence of RNase A for 45 minutes in dark. Samples were analyzed on a FACSCalibur flow-cytometer (Beckman Coulter, Inc., Fullerton, CA).

Immunofluorescence Confocal Microscopy

SK-SY5Y and LA1-5S cells were grown on poly(L-lysine)-coated glass coverslips for immunofluorescence microscopy as described previously [10,40]. After treatment, cells were fixed with cold (−20°C) methanol for 5 minutes and blocked by incubating with 2% BSA/PBS at 37°C for 1 hr. A mouse monoclonal antibody against α -tubulin (DM1A, Sigma) was diluted 1:500 in 2% BSA/PBS and incubated with the coverslips for 2 hrs at 37°C. Cells were then washed with 2% BSA/PBS for 10 minutes at room temperature before incubating with a 1:200 dilution of a FITC-labeled goat anti-mouse IgG antibody (Jackson ImmunoResearch, Inc., West Grove, PA) at 37°C for 1 hr. Coverslips were then rinsed with 2% BSA/PBS for 10 minutes and incubated with PI (0.5 µg/mL) for 15 minutes at room temperature before they were mounted with Aquamount (Lerner Laboratories, Pittsburgh, PA)

containing 0.01% 1,4-diazobicyclo(2,2,2)octane (Sigma). Cells were examined using confocal microscopy for microtubule and nuclear morphology using a 63× objective (numerical aperture, 1.4).

Western Blot Analysis

Proteins were resolved by polyacrylamide gel-electrophoresis and transferred onto polyvinylidene difluoride membranes (Millipore). Membranes were blocked in Trisbuffered saline containing 0.2% Tween-20 and 5% fatfree dry milk and incubated first with primary antibodies and then with horseradish peroxidase-conjugated secondary antibodies. Specific proteins were visualized with enhanced chemiluminescence detection reagent according to the manufacturer's instructions (Pierce Biotechnology).

Quantitative Real-time Reverse Transcriptase-polymerase Chain Reaction

Total cellular RNA was extracted from SK-SY5Y and LA1-5S cells using Trizol reagent (Gibco) according to the manufacturer's instructions. Preservation of 28S and 18S rRNA species was used to assess RNA integrity. Only samples with prominent 28S and 18S rRNA components were included in the study. Of the total RNA, 1 µg from each sample was reverse transcribed to complementary DNA (cDNA) using Moloney Murine Leukemia Virus reverse transcriptase (Invitrogen, Life Technologies, Paisley, UK) and random primers (Invitrogen). Survivin mRNA expression was evaluated by real-time polymerase chain reaction using FastStart Universal SYBR Green Master (ROX) (Roche, Mannheim, Germany) in a iCycler Optical Module (Bio-Rad, Hercules, CA, USA). Reactions were performed in triplicate using 2 µL of cDNA per reaction and primers specific for survivin (forward: 5' CGA GGC TGG CTT CAT CCA 3'; reverse: 5' GCA ACC GGA CGA ATG CTT T 3'), while human porphobilinogen deaminase was used as a housekeeping gene (forward: 5' AGA GTG ATT CGC GTG GGT ACC 3'; reverse: 5' GGC TCC GAT GGT GAA GCC 3').

References

1. Maris JM, Matthay KK (1999) Molecular biology of neuroblastoma. *J Clin Oncol* 17: 2264–2279.
2. Duflos A, Kruczynski A, Barret JM (2002) Novel aspects of natural and modified vinca alkaloids. *Current medicinal chemistry Anti-cancer agents* 2: 55–70.
3. Rowinsky EK (1997) The development and clinical utility of the taxane class of antimicrotubule chemotherapy agents. *Annual review of medicine* 48: 353–374.
4. Arkin M (2005) Protein-protein interactions and cancer: small molecules going in for the kill. *Curr Opin Chem Biol* 9: 317–324.
5. Crown J, O'Leary M (2000) The taxanes: an update. *Lancet* 355: 1176–1178.
6. Zhou J, Giannakakou P (2005) Targeting microtubules for cancer chemotherapy. *Current medicinal chemistry Anti-cancer agents* 5: 65–71.
7. Anderson JT, Ting AE, Boozer S, Brunden KR, Crumrine C, et al. (2005) Identification of novel and improved antimetabolic agents derived from noscapine. *Journal of medicinal chemistry* 48: 7096–7098.
8. Anderson JT, Ting AE, Boozer S, Brunden KR, Danzig J, et al. (2005) Discovery of S-phase arresting agents derived from noscapine. *Journal of medicinal chemistry* 48: 2756–2758.
9. Aneja R, Zhou J, Vangapandu SN, Zhou B, Chandra R, et al. (2006) Drug-resistant T-lymphoid tumors undergo apoptosis selectively in response to an antimicrotubule agent, EM011. *Blood* 107: 2486–2492.
10. Zhou J, Gupta K, Aggarwal S, Aneja R, Chandra R, et al. (2003) Brominated derivatives of noscapine are potent microtubule-interfering agents that perturb mitosis and inhibit cell proliferation. *Molecular pharmacology* 63: 799–807.
11. Landen JW, Lang R, McMahon SJ, Rusan NM, Yvon AM, et al. (2002) Noscapine alters microtubule dynamics in living cells and inhibits the progression of melanoma. *Cancer research* 62: 4109–4114.
12. Zhou J, Gupta K, Yao J, Ye K, Panda D, et al. (2002) Paclitaxel-resistant human ovarian cancer cells undergo c-Jun NH2-terminal kinase-mediated apoptosis in response to noscapine. *The Journal of biological chemistry* 277: 39777–39785.
13. Ke Y, Ye K, Grossniklaus HE, Archer DR, Joshi HC, et al. (2000) Noscapine inhibits tumor growth with little toxicity to normal tissues or inhibition of immune responses. *Cancer immunology, immunotherapy* : CII 49: 217–225.

Gene Silencing with Small Interfering RNAs

siRNA oligonucleotides were purchased from Ambion/Applied Biosystems (Austin, TX). All siRNAs were reconstituted according to the manufacturer's instructions. Briefly, The SK-SY5Y and LA1-5S cells were seeded in six-well plates and transfected at ~50% confluency with 100 nmol/L of p53- or survivin-targeted siRNA using Oligofectamine. Twenty-four hours after transfection, the cells were treated for 24 hours with either DMSO (control) or 50 µmol/L Noscapine. Subsequently, the cells were collected and processed for immunoblotting and flow cytometric analysis.

Ectopic Expression of Survivin by Transient Transfection

The SK-SY5Y and LA1-5S cells were transiently transfected at ~50% confluency with the empty pcDNA3.1 vector or pcDNA3.1 encoding for survivin (Addgene) using FuGENE6 transfection reagent. Twentyfour hours after transfection, the cells were treated with DMSO (control) or Noscapine for specified time period. Cells were collected and processed for immunoblotting and measurement for apoptosis.

Statistical Analysis

Each experiment was repeated at least twice with triplicate measurements for quantitative comparisons. Statistical significance of difference in measured variables between control and treated groups was determined by t test or one-way ANOVA. Difference was considered significant at P<0.05.

Acknowledgments

We thank Qinglan Ruan, Hong Mei, Ying Yang for experimental discussion and assistance.

Author Contributions

Conceived and designed the experiments: SL JH. Performed the experiments: SL GC JH. Analyzed the data: HCJ ST. Contributed reagents/materials/analysis tools: HCJ QT. Wrote the paper: SL.

14. Heidari N, Goliaei B, Moghaddam PR, Rahbar-Roshandel N, Mahmoudian M (2007) Apoptotic pathway induced by noscapine in human myelogenous leukemic cells. *Anti-cancer drugs* 18: 1139–1147.
15. Landen JW, Hau V, Wang M, Davis T, Ciliax B, et al. (2004) Noscapine crosses the blood-brain barrier and inhibits glioblastoma growth. *Clinical cancer research : an official journal of the American Association for Cancer Research* 10: 5187–5201.
16. Ye K, Ke Y, Keshava N, Shanks J, Kapp JA, et al. (1998) Opium alkaloid noscapine is an antitumor agent that arrests metaphase and induces apoptosis in dividing cells. *Proceedings of the National Academy of Sciences of the United States of America* 95: 1601–1606.
17. Vichai V, Kirtikara K (2006) Sulforhodamine B colorimetric assay for cytotoxicity screening. *Nature protocols* 1: 1112–1116.
18. Aneja R, Ghaleb AM, Zhou J, Yang VW, Joshi HC (2007) p53 and p21 determine the sensitivity of noscapine-induced apoptosis in colon cancer cells. *Cancer research* 67: 3862–3870.
19. Vousden KH, Lu X (2002) Live or let die: the cell's response to p53. *Nature reviews Cancer* 2: 594–604.
20. Srinivasula SM, Ashwell JD (2008) IAPs: what's in a name? *Molecular cell* 30: 123–135.
21. Ghavami S, Hashemi M, Ande SR, Yeganeh B, Xiao W, et al. (2009) Apoptosis and cancer: mutations within caspase genes. *Journal of medical genetics* 46: 497–510.
22. Dubrez-Daloz L, Dupoux A, Cartier J (2008) IAPs: more than just inhibitors of apoptosis proteins. *Cell cycle* 7: 1036–1046.
23. Rothe M, Pan MG, Henzel WJ, Ayres TM, Goeddel DV (1995) The TNFR2-TRAF signaling complex contains two novel proteins related to baculoviral inhibitor of apoptosis proteins. *Cell* 83: 1243–1252.
24. Ambrosini G, Adida C, Altieri DC (1997) A novel anti-apoptosis gene, survivin, expressed in cancer and lymphoma. *Nature medicine* 3: 917–921.

25. Tamm I, Wang Y, Sausville E, Scudiero DA, Vigna N, et al. (1998) IAP-family protein survivin inhibits caspase activity and apoptosis induced by Fas (CD95), Bax, caspases, and anticancer drugs. *Cancer research* 58: 5315–5320.
26. Tanaka K, Iwamoto S, Gon G, Nohara T, Iwamoto M, et al. (2000) Expression of survivin and its relationship to loss of apoptosis in breast carcinomas. *Clinical cancer research : an official journal of the American Association for Cancer Research* 6: 127–134.
27. Tamm I, Richter S, Oltersdorf D, Creutzig U, Harbott J, et al. (2004) High expression levels of x-linked inhibitor of apoptosis protein and survivin correlate with poor overall survival in childhood de novo acute myeloid leukemia. *Clinical cancer research : an official journal of the American Association for Cancer Research* 10: 3737–3744.
28. Li F, Altieri DC (1999) The cancer antiapoptosis mouse survivin gene: characterization of locus and transcriptional requirements of basal and cell cycle-dependent expression. *Cancer research* 59: 3143–3151.
29. Reed JC, Bischoff JR (2000) BIRing chromosomes through cell division— and survivin' the experience. *Cell* 102: 545–548.
30. Lens SM, Medema RH (2003) The survivin/Aurora B complex: its role in coordinating tension and attachment. *Cell cycle* 2: 507–510.
31. Hoffman WH, Biade S, Zilfou JT, Chen J, Murphy M (2002) Transcriptional repression of the anti-apoptotic survivin gene by wild type p53. *The Journal of biological chemistry* 277: 3247–3257.
32. Mirza A, McGuirk M, Hockenberry TN, Wu Q, Ashar H, et al. (2002) Human survivin is negatively regulated by wild-type p53 and participates in p53-dependent apoptotic pathway. *Oncogene* 21: 2613–2622.
33. Kaufmann SH, Desnoyers S, Ottaviano Y, Davidson NE, Poirier GG (1993) Specific proteolytic cleavage of poly(ADP-ribose) polymerase: an early marker of chemotherapy-induced apoptosis. *Cancer research* 53: 3976–3985.
34. Gu L, Zhu N, Zhang H, Durden DL, Feng Y, et al. (2009) Regulation of XIAP translation and induction by MDM2 following irradiation. *Cancer cell* 15: 363–375.
35. He J, Gu L, Zhang H, Zhou M (2011) Crosstalk between MYCN and MDM2-p53 signal pathways regulates tumor cell growth and apoptosis in neuroblastoma. *Cell cycle* 10: 2994–3002.
36. Gu L, Zhang H, He J, Li J, Huang M, et al. (2011) MDM2 regulates MYCN mRNA stabilization and translation in human neuroblastoma cells. *Oncogene*.
37. von Reitzenstein C, Kopitz J, Schuhmann V, Cantz M (2001) Differential functional relevance of a plasma membrane ganglioside sialidase in cholinergic and adrenergic neuroblastoma cell lines. *European journal of biochemistry/FEBS* 268: 326–333.
38. Tanaka N, Fukuzawa M (2008) MYCN downregulates integrin alpha1 to promote invasion of human neuroblastoma cells. *International journal of oncology* 33: 815–821.
39. Skehan P, Storeng R, Scudiero D, Monks A, McMahon J, et al. (1990) New colorimetric cytotoxicity assay for anticancer-drug screening. *Journal of the National Cancer Institute* 82: 1107–1112.
40. Aneja R, Zhou J, Zhou B, Chandra R, Joshi HC (2006) Treatment of hormone-refractory breast cancer: apoptosis and regression of human tumors implanted in mice. *Molecular cancer therapeutics* 5: 2366–2377.