

Analysis of input-output relationships of CPG elements and their contributions to rhythmic output

Terrence Michael Wright, *Emory University* Brian Mulloney, *University of California, Davis* <u>Ronald Calabrese</u>, *Emory University*

Journal Title: BMC Neuroscience Volume: Volume 13, Number Suppl 1 Publisher: BioMed Central | 2012-07-16, Pages P171-P171 Type of Work: Article | Final Publisher PDF Publisher DOI: 10.1186/1471-2202-13-S1-P171 Permanent URL: https://pid.emory.edu/ark:/25593/s5gpb

Final published version: http://dx.doi.org/10.1186/1471-2202-13-S1-P171

Copyright information:

©2012 Wright et al This is an Open Access work distributed under the terms of the Creative Commons Attribution 2.0 Generic License (<u>http://creativecommons.org/licenses/by/2.0/</u>).



Accessed January 20, 2022 2:38 AM EST

POSTER PRESENTATION



Open Access

Analysis of input-output relationships of CPG elements and their contributions to rhythmic output

Terrence Michael Wright Jr^{1,2*}, Brian Mulloney², Ronald L Calabrese¹

From Twenty First Annual Computational Neuroscience Meeting: CNS*2012 Decatur, GA, USA. 21-26 July 2012

We use the dynamic clamp technique [1] to explore how synaptic input patterns affect motor output (see Figure 1A). We show that leech motor neuron intrinsic properties make a contribution to their output phasing. Then, we show that leech motor neurons receiving the same complement of synaptic inputs can still be organized into a coordinated motor pattern given that a gradient of synaptic strengths exists and that the premotor interneurons fire at different times [2]. In both of these

cases, measuring motor neuron responses provides a direct assay for how premotor input patterns produce stereotyped motor output. We are currently extending this analysis to the crayfish swimmeret system, a system in which four segmental oscillators are interconnected by coordinating interneurons to produce a metachronal wave of swimmeret movements [3]. In this system, ascending (ASC) and descending (DSC) coordinating interneurons (see Figure 1B) encode salient features



segment 4 (A4) driving PS4 motor activity observed in the extracellular recordings (right panel; Return stroke [RS(4)] activity not shown). Coordinating interneurons ASC and DSC in A4 encode information that is delivered to an interneuron (C1) in segments 3 (ASC) and 5 (DSC), resulting in the stereotyped rear-to-front activity pattern observed across segments 5,4 and 3 (right panel).

* Correspondence: terrencemwright@msn.com

¹Department of Biology, Emory University, Atlanta, GA 30322, USA

Full list of author information is available at the end of the article



© 2012 Wright et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

about the activity pattern of their home segment and are exported (via spikes) to other segmental oscillators. Interestingly, when the system is driven across periods by different concentrations of neuromodulators, the number of spikes of a given coordinating interneuron remains constant although their duty cycles change. We are currently building a single-compartment, conductance-based model of the ASC and DSC coordinating neurons, with the goal of understanding how these coordinating neurons encode information in their home modules and how this encoding can be modulated when the swimmeret system is driven at different periods.

Author details

¹Department of Biology, Emory University, Atlanta, GA 30322, USA. ²Department of Neurobiology, Physiology and Behavior, University of California, Davis, Davis, CA, 95616, CA, USA.

Published: 16 July 2012

References

- Sharp AA, O'neil MB, Abbott LF, Marder E: Dynamic clamp: artificial conductances in biological neurons. *Trends Neurosci* 1993, 16(10):389-94.
- Wright TM, Calabrese RL: Patterns of presynaptic activity interact to produce motor output. J Neurosci 2011, 31(48):17555-71.
- Smarandache-Wellmann C, Mulloney B: Neurobiology of the crayfish swimmeret system. Prog Neurobiol 2012, 96(2):242-67.

doi:10.1186/1471-2202-13-S1-P171

Cite this article as: Wright *et al.*: **Analysis of input-output relationships of CPG elements and their contributions to rhythmic output.** *BMC Neuroscience* 2012 **13**(Suppl 1):P171.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

BioMed Central

Submit your manuscript at www.biomedcentral.com/submit