



P02.119. Cognitively-Based Compassion Training reduces peripheral inflammation in adolescents in foster care with high rates of early life adversity

Thaddeus Pace, *Emory University*
L Negi, *Emory University*
B Donaldson-Lavelle, *Emory University*
[Brendan Ozawa-de Silva](#), *Emory University*
Satyanarayan Reddy, *Emory University*
S Cole, *Research Design Associates, Inc.*
[Linda Craighead](#), *Emory University*
C Raison, *University of Arizona*

Journal Title: BMC Complementary and Alternative Medicine

Volume: Volume 12, Number Suppl 1

Publisher: BioMed Central | 2012-06-12, Pages P175-P175

Type of Work: Article | Final Publisher PDF

Publisher DOI: 10.1186/1472-6882-12-S1-P175

Permanent URL: <https://pid.emory.edu/ark:/25593/s5jfk>

Final published version: <http://dx.doi.org/10.1186/1472-6882-12-S1-P175>

Copyright information:

©2012 Pace et al

This is an Open Access work distributed under the terms of the Creative Commons Attribution 2.0 Generic License (<http://creativecommons.org/licenses/by/2.0/>).



POSTER PRESENTATION

Open Access

P02.119. Cognitively-Based Compassion Training reduces peripheral inflammation in adolescents in foster care with high rates of early life adversity

T Pace^{1*}, L Negi¹, B Donaldson-Lavelle¹, B Ozawa-de Silva¹, S Reddy¹, S Cole², L Craighead¹, C Raison³

From International Research Congress on Integrative Medicine and Health 2012
Portland, Oregon, USA. 15-18 May 2012

Purpose

Children exposed to early life adversity (ELA) demonstrate elevated circulating concentrations of health-relevant inflammatory biomarkers which persist into adulthood. Increased inflammation in individuals with ELA is believed to contribute to the increased risk for medical and psychiatric illnesses observed in these individuals. The objective of this study was to determine whether Cognitively-Based Compassion Training (CBCT) reduces salivary concentrations of C-reactive protein (CRP) in adolescents with high rates of ELA, and to evaluate the relationship between CBCT practice time and changes in CRP. CBCT is a meditation-based program designed to enhance compassion for self and others and to promote prosocial behavior. Based on prior findings, it was hypothesized that practice time during the study would be more strongly associated with reductions in CRP than would group assignment.

Methods

Seventy-one adolescents between the ages of 13 and 17 (31 females) in the Georgia Foster Care system were randomized to either six weeks of CBCT or a wait-list control condition. Saliva was obtained upon awakening prior to randomization and again 6 weeks later, as were self-report measures of depression and anxiety. Saliva was assayed for CRP using a high-sensitivity ELISA (Salimetrics, State College, PA). Trauma/neglect history was obtained from state records. Participants completed practice time diaries as a means of assessing amount of engagement with the CBCT program.

Results

No between group differences were observed in salivary CRP concentrations, self-reported depression or anxiety symptoms. Within the group randomized to CBCT, increased practice time was associated with reduced CRP from baseline to the six week assessment.

Conclusion

Engagement with CBCT positively impacts an inflammatory biomarker relevant to health in adolescents at high risk for poor adult health and social functioning as a result of significant early life adversity, including placement in foster care.

Author details

¹Emory University, Atlanta, USA. ²Research Design Associates, Inc., Yorktown Heights, USA. ³University of Arizona College of Medicine, Department of Psychiatry, Tucson, USA.

Published: 12 June 2012

doi:10.1186/1472-6882-12-S1-P175

Cite this article as: Pace et al.: P02.119. Cognitively-Based Compassion Training reduces peripheral inflammation in adolescents in foster care with high rates of early life adversity. *BMC Complementary and Alternative Medicine* 2012 12(Suppl 1):P175.

¹Emory University, Atlanta, USA

Full list of author information is available at the end of the article