



Interstage mortality after the Norwood procedure: Results of the multicenter Single Ventricle Reconstruction trial

Nancy S. Ghanayem, *Medical College of Wisconsin*
Kerstin R. Allen, *New England Research Institutes*
Sarah Tabbutt, *Children's Hospital of Philadelphia*
Andrew M. Atz, *Medical University of South Carolina*
[Martha Clabby](#), *Emory University*
David S. Cooper, *Congenital Heart Institute of Florida*
Pirooz Eghtesady, *Cincinnati Children's Medical Center*
Peter C. Frommelt, *Medical College of Wisconsin*
Peter J. Gruber, *Children's Hospital of Philadelphia*
Kevin D. Hill, *Duke University*

Only first 10 authors above; see publication for full author list.

Journal Title: Journal of Thoracic and Cardiovascular Surgery

Volume: Volume 144, Number 4

Publisher: Elsevier | 2012-10-01, Pages 896-906

Type of Work: Article | Post-print: After Peer Review

Publisher DOI: 10.1016/j.jtcvs.2012.05.020

Permanent URL: <https://pid.emory.edu/ark:/25593/s9dff>

Final published version: <http://dx.doi.org/10.1016/j.jtcvs.2012.05.020>

Copyright information:

© 2012 The American Association for Thoracic Surgery. Published by Elsevier Inc. All rights reserved.

This is an Open Access work distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



Published in final edited form as:

J Thorac Cardiovasc Surg. 2012 October ; 144(4): 896–906. doi:10.1016/j.jtcvs.2012.05.020.

Interstage mortality after the Norwood procedure: Results of the multicenter Single Ventricle Reconstruction trial

Nancy S. Ghanayem, MD^a, Kerstin R. Allen^b, Sarah Tabbutt, MD, PhD^c, Andrew M. Atz, MD^d, Martha L. Clabby, MD^e, David S. Cooper, MD^f, Pirooz Egthesady, MD^g, Peter C. Frommelt, MD^a, Peter J. Gruber, MD^c, Kevin D. Hill, MD^h, Jonathan R. Kaltman, MDⁱ, Peter C. Laussen, MBBS^j, Alan B. Lewis, MD^k, Karen J. Lurito, MD^l, L. LuAnn Minich, MD^m, Richard G. Ohye, MDⁿ, Julie V. Schonbeck^b, Steven M. Schwartz, MD^o, Rakesh K. Singh, MD^p, and Caren S. Goldberg, MDⁿ for the Pediatric Heart Network Investigators

^aChildren's Hospital of Wisconsin and Medical College of Wisconsin, Milwaukee, Wis

^bNew England Research Institutes, Watertown, Mass

^cChildren's Hospital of Philadelphia, Philadelphia, Pa

^dMedical University of South Carolina, Charleston, SC

^eEmory University, Atlanta, Ga

^fCongenital Heart Institute of Florida, St. Petersburg, Fla

^gCincinnati Children's Medical Center, Cincinnati, Ohio

^hDuke University, Durham, NC

ⁱNorth Carolina Consortium: National Heart, Lung, and Blood Institute, Bethesda, Md

^jChildren's Hospital Boston, Boston, Mass

^kChildren's Hospital Los Angeles, Los Angeles, Calif

^lEast Carolina University, Greenville, NC

^mPrimary Children's Medical Center and the University of Utah, Salt Lake City, Utah

ⁿUniversity of Michigan Medical School, Ann Arbor, Mich

^oHospital for Sick Children, Toronto, Ontario, Canada

^pColumbia University, New York, NY

Abstract

Copyright © 2012 by The American Association for Thoracic Surgery

Address for reprints: Nancy S. Ghanayem, MD, Medical College of Wisconsin, Children's Hospital of Wisconsin, Pediatrics, 9000 West Wisconsin Ave, PO Box 1997, Milwaukee, WI 53226 (nancyg@mcw.edu).

Kerstin R. Allen's current affiliation is Infinity Pharmaceuticals, Boston, Mass; Sarah Tabbutt's current affiliation is University of California San Francisco, San Francisco, Calif; Peter J. Gruber's current affiliation is Primary Children's Medical Center and University of Utah, Salt Lake City, Utah; David S. Cooper's and Pirooz Egthesady's current affiliation is Cincinnati Children's Medical Center, Cincinnati, Ohio.

Pediatric Heart Network Investigators are listed in Appendix 1.

Disclosures: Authors have nothing to disclose with regard to commercial support.

Objective—For infants with single ventricle malformations undergoing staged repair, interstage mortality is reported at 2% to 20%. The Single Ventricle Reconstruction trial randomized subjects with a single morphologic right ventricle undergoing a Norwood procedure to a modified Blalock–Taussig shunt (MBTS) or a right ventricle-to-pulmonary artery shunt (RVPAS). The aim of this analysis was to explore the associations of interstage mortality and shunt type, and demographic, anatomic, and perioperative factors.

Methods—Participants in the Single Ventricle Reconstruction trial who survived to discharge after the Norwood procedure were included (n = 426). Interstage mortality was defined as death postdischarge after the Norwood procedure and before the stage II procedure. Univariate analysis and multivariable logistic regression were performed adjusting for site.

Results—Overall interstage mortality was 50 of 426 (12%)—13 of 225 (6%) for RVPAS and 37 of 201 (18%) for MBTS (odds ratio [OR] for MBTS, 3.4; $P < .001$). When moderate to severe postoperative atrioventricular valve regurgitation (AVVR) was present, interstage mortality was similar between shunt types. Interstage mortality was independently associated with gestational age less than 37 weeks (OR, 3.9; $P = .008$), Hispanic ethnicity (OR, 2.6; $P = .04$), aortic atresia/mitral atresia (OR, 2.3; $P = .03$), greater number of post-Norwood complications (OR, 1.2; $P = .006$), census block poverty level ($P = .003$), and MBTS in subjects with no or mild postoperative AVVR (OR, 9.7; $P < .001$).

Conclusions—Interstage mortality remains high at 12% and is increased with the MBTS compared with the RVPAS if postoperative AVVR is absent or mild. Preterm delivery, anatomic, and socioeconomic factors are also important. Avoiding preterm delivery when possible and close surveillance after Norwood hospitalization for infants with identified risk factors may reduce interstage mortality.

Survivors of the Norwood procedure for hypoplastic left heart syndrome (HLHS) and other single right ventricle anomalies are left with the combination of an inefficient parallel circulation with volume load to the systemic ventricle, potential inferior pumping capability of a systemic right ventricle, and risk of compromise of the systemic to pulmonary artery shunt. These factors expose patients to a heightened risk for circulatory collapse. This inherently fragile physiology persists to the stage II procedure when the parallel circulation is eliminated through takedown of the systemic to pulmonary artery shunt and creation of a cavopulmonary connection. Mortality postdischarge after the Norwood hospitalization and before the stage II procedure (interstage period) is reported to be 2% to 20%.¹⁻⁴

Specific risks associated with interstage mortality identified in various investigations include a diminutive ascending aorta as seen in aortic atresia, the presence of a restrictive atrial communication, postoperative arch obstruction, obstructed shunt flow, pulmonary artery distortion, and atrioventricular valve regurgitation (AVVR).^{5,6} Age at surgery, postoperative arrhythmias, airway complications, feeding difficulties, and noncardiac disease processes such as gastroenteritis or upper respiratory infection also have been implicated.^{3,4} Compared with the right ventricle-to-pulmonary artery shunt (RVPAS), higher interstage mortality has been reported for the modified Blalock–Taussig shunt (MBTS).^{7,8}

The National Heart, Lung, and Blood Institute–sponsored Pediatric Heart Network Single Ventricle Reconstruction (SVR) trial includes the largest prospective cohort of infants with

HLHS or related single right ventricle anomalies with longitudinal follow-up after the Norwood procedure. The primary results of the SVR trial demonstrated improved transplant-free survival at 12 months in those randomized to receive an RVPAS at the Norwood procedure compared with those randomized to the MBTS. Although interstage survival was not specifically reported, the highest incidence of transplant or death occurred between 30 days after the Norwood procedure and the stage II procedure.⁹

The goal of this analysis was to determine anatomic, surgical, and additional patient-related risk factors for interstage mortality postdischarge after the Norwood procedure and before the stage II procedure in this unique cohort. We specifically hypothesized that infants palliated with an MBTS would be at increased risk of interstage death compared with those palliated with an RVPAS.

Materials and Methods

The Pediatric Heart Network SVR trial compared outcomes between subjects randomized to RVPAS and subjects randomized to MBTS at the time of the Norwood procedure. Details of the trial design have been reported.¹⁰ The institutional review board or research ethics board at each participating center approved the study protocol, and written informed consent was obtained from parents before trial enrollment.

Study Sample

Subjects randomized in the multicenter SVR trial who survived to discharge from the hospital after the Norwood procedure are included in this analysis. Twenty-two subjects surviving the Norwood procedure, but not discharged before the stage II procedure, were excluded from the primary analysis because of limitations in distinguishing Norwood operative mortality from interstage mortality for subjects with planned inpatient care to stage II procedure. Two subjects whose dates of stage II procedure could not be determined and 2 subjects whose stage II procedures were performed uncharacteristically late (ie, >14 months of age) were excluded from the primary analysis.

Study Design and Measurements

Shunt type was defined for this analysis as the shunt in place at the end of the Norwood procedure. The surgeon had the option to modify the shunt if anatomy was encountered that made the randomized shunt assignment not feasible. Other than random assignment of shunt, all participants were managed according to the standard practices at their clinical centers. A list of all variables that were recorded and analyzed is included in Appendix Table 2. In brief, before the Norwood procedure, a detailed preoperative medical history was recorded, including demographics, patient characteristics, and anatomic diagnosis. Operative variables included shunt type, pharmacologic strategies, perfusion method, and additional cardiac operations. The perfusion method during circulatory support was classified into 1 of 3 categories: deep hypothermic circulatory arrest (DHCA), regional cerebral perfusion with DHCA time of 10 minutes or less, or regional cerebral perfusion with DHCA time greater than 10 minutes. Postoperative data recorded for the Norwood hospitalization included procedures, serious adverse events such as cardiopulmonary resuscitation (CPR) and use of

extracorporeal membrane oxygenation (ECMO), number of previously described complications,¹¹ hospital length of stay, and feeding methods at discharge after the Norwood procedure.

Echocardiograms were obtained before and after the Norwood procedure. The echocardiograms were interpreted centrally at a core laboratory (Medical College of Wisconsin) to assess the degree of AVVR (none/mild vs moderate/severe), to measure right ventricular end-systolic volume (RVESV) and right ventricular end-diastolic volume (RVEDV), and to determine fractional area change and right ventricular ejection fraction (RVEF). The primary measure of right ventricular contractility used in the multivariable model was right ventricular fractional area change, because the other measures of systolic function including RVEF, RVESV, and RVEDV could not be obtained for 23% of the subjects. In the 330 subjects for whom both measurements were available, there was a high degree of correlation ($R = 0.89$) between the RVEF and the right ventricular fractional area change.

A subgroup of subjects consented for evaluation by a geneticist. Genetic evaluations performed for clinical indications were also recorded. Subjects were classified as to whether a specific genetic syndrome was identified and whether “other abnormalities” (ie, not identified with a syndrome) were present.

Socioeconomic status at the time of randomization was assigned using a US census-based score derived from 6 measures based on income, housing, and occupational-related features of the subject's census block tract, as well as a score measuring the percent below the poverty level in the subject's census block tract.¹²

Statistical Methods

Descriptive statistics presented include median with interquartile range for skewed variables, mean and standard deviation for other continuous variables, and frequency with percentage for categorical variables. Tertiles of continuous measures were fit to determine whether the association between predictor and outcome was nonlinear. Dichotomies were also explored.

Univariate logistic regression and logistic regression adjusted for site were used to obtain initial estimates of association between interstage survival and each candidate predictor. Predictors significant at P less than .20 at the univariate or site-adjusted level were included in the multivariable modeling. Multivariable models were developed using stepwise logistic regression with bootstrap bagging 1000 samples to determine the reliability of each individual predictor. The small number of events made the bootstrapping procedure unstable when site (14 degrees of freedom) was a candidate predictor; thus, the annual center volume of screened patients with a single right ventricle and planned Norwood procedure was included as a surrogate for site in bootstrapping models. A predictor was retained in the final site-adjusted multivariable model if bootstrapping reliability was greater than 45% and its P value was less than .05.

To determine whether some prespecified factors had differential associations dependent on shunt type, interaction terms were explored. Specifically, interactions of shunt type with

birth weight (<2.5 vs 2.5 kg), low gestational age (<37 weeks), ethnicity, obstructed pulmonary venous return, presence of aortic atresia, pre-Norwood and post-Norwood AVVR, and ECMO or CPR during the Norwood hospitalization were examined for their association with interstage death or transplant. After the multivariable model main effects were determined, the significant interactions were added and retained if significant at a *P* value less than .05.

In an exploratory analysis, the multivariable model was developed using the same approach, but after imputing missing values of RVEF, RVESV indexed to body surface area (BSA)^{1,3}, and RVEDV indexed to BSA^{1,3}. Simple regression imputation was used for RVEF, which had 23% of the data missing at random. Right ventricular fractional area change was used to predict RVEF. For 17 subjects (4%) who were missing both measurements and for subjects missing RVESV or RVEDV, mean imputation was used. These imputed predictors were included in the model selection process. All analyses were conducted using SAS version 9.2 (SAS Institute, Inc, Cary, NC).

Results

Study Sample

Of the 549 subjects who were randomized in the SVR trial and underwent the Norwood procedure, 97 died (88) or underwent heart transplantation (9) during the Norwood hospitalization, 22 remained as inpatients until the stage II procedure, and 430 were discharged before the stage II procedure (Figure 1). The 22 subjects who survived the Norwood procedure and were not discharged before the stage II procedure differed from those who were discharged before the stage II procedure in that hospitalized subjects had a higher rate of ECMO, CPR, moderate or greater AVVR, and postoperative complications (Appendix Table 3). Four subjects discharged were excluded from the analysis because of an uncharacteristically late stage II procedure (*n* = 2) or unknown timing of the stage II procedure (*n* = 2). The remaining 426 subjects were discharged after the Norwood procedure and formed the analytic cohort. No subject discharged after the Norwood procedure underwent heart transplantation before the stage II procedure.

Interstage Survival

Of the 426 interstage subjects, 376 (88%) survived to the stage II procedure. Interstage survival was higher in subjects who received the RVPAS (212/225; 94%) compared with those who received the MBTS (164/201; 82%). The odds ratio (OR) for interstage mortality for the MBTS versus the RVPAS was 3.4 (95% confidence interval, 1.8-6.6; *P* < .001) (Figure 2). Of the subjects who died during the interstage period, the mean time to death postdischarge after the Norwood procedure was earlier in the MBTS group (1.7 ± 1.6 months vs 2.8 ± 1.5 months postdischarge after the Norwood procedure, *P* < .001), with a mean age at death 3.0 ± 2.0 months in the MBTS group and 4.3 ± 1.7 months in the RVPAS group (*P* = .01). The stage II procedure was performed at similar ages for the 2 shunt types (5.5 ± 1.7 months vs 5.3 ± 1.6 months, *P* = .34).

Risk Factors for Interstage Mortality

Univariate analysis—Univariate analysis was performed to assess the association of each potential risk factor with interstage mortality (Table 1). In addition to the MBTS, younger gestational age, census tract block with 5.4% to 13% of inhabitants below federal poverty level, ascending aorta diameter less than 3 mm, longer Norwood hospital length of stay (median, 25; interquartile range, 17 to 44 days vs 23 days; interquartile range, 16-36), greater number of postoperative complications, and failure to feed orally before hospital discharge were associated with interstage mortality. The risk of interstage mortality was also higher in subjects without any oral feeding with a nasal enteral tube compared with a gastrostomy tube (OR, 5.0; 95% confidence interval, 1.5-17.1; $P = .01$). Moderate to severe postoperative AVVR, lower RVEF, and greater RVESV indexed to BSA measured on post-Norwood echocardiograms (performed 15 ± 10 days after the Norwood procedure) were each associated with interstage mortality. Site was not a significant predictor of interstage mortality ($P = .16$). Genetic syndrome and nonsyndromic comorbidities also were not associated with interstage mortality.

Of the prespecified factors, only the interaction of shunt type and postoperative AVVR was significantly associated with interstage mortality ($P = .009$). After adjustment for site, the influence of shunt type varied with degree of AVVR. In subjects with postoperative none/mild AVVR, interstage mortality was significantly higher for the MBTS group (MBTS 18% vs RVPAS 3% $P < .001$; OR, 6.7). However, for subjects with postoperative moderate/severe AVVR, the overall rate of interstage mortality was similar between the shunt types (MBTS 20% vs RVPAS 17% $P = .94$; OR, 1.0).

Multivariable analysis—Independent predictors of interstage mortality with adjustment for site determined through multivariable modeling (adjusted $R^2 = 0.35$) included gestational age less than 37 weeks (OR, 3.9; $P = .008$), Hispanic ethnicity (OR, 2.6; $P = .04$), census block poverty level ($P = .003$), presence of aortic and mitral atresia (OR, 2.3; $P = .03$), MBTS in subjects with postoperative none/mild AVVR (OR, 9.7; $P < .001$), and greater number of post-Norwood complications (OR, 1.2, per complication; $P = .006$). With respect to census block poverty levels, subjects in communities with 5.4% to 13% poverty had a greater risk of interstage mortality compared with subjects in the more affluent communities (OR, 6.7) and the poorest communities (OR, 2.5) (Table 2). The exploratory analysis that included imputed missing values of RVEF, RVESV, RVEDV, and RV fractional area change showed that these potential predictors were not independent risk factors of interstage mortality.

Discussion

This is the first multicenter prospective report of interstage mortality for infants with HLHS and other single right ventricle anomalies undergoing staged repair. The overall interstage mortality rate of 12% from this large cohort is similar to that of multiple prior single-center case series.^{1,3,4,7,8,13}

Subjects who had an MBTS were at higher risk for interstage mortality than those with RVPAS. With pulmonary blood flow occurring only during systole after palliation with the

RVPAS, a higher diastolic pressure and lower pulmonary to systemic flow ratio has been observed.^{14,15} Although we are unable to determine the mechanisms of interstage death in this population, it is possible that the higher diastolic pressure associated with RVPAS contributes to improved coronary and systemic perfusion, providing advantageous hemodynamic stability during periods of stress such as illness or feeding difficulties. Patients with an MBTS may have a higher risk for acute shunt thrombosis; however, the incidence of shunt thrombosis as a cause for death was not examined in this analysis.

Although only shunt type was randomized in the SVR trial, the sample size and prospective data collection allowed for evaluation of other potential predictors of interstage mortality. Previous reports have implicated anatomic diagnosis, residual or recurrent lesions, arrhythmias, subject specific characteristics, acquired illness, and, more recently, shunt type as risk factors for interstage death.^{1,3-8,13} Although Jonas and colleagues⁵ found the combination of aortic atresia with mitral stenosis to be associated with interstage mortality for infants with HLHS,⁵ other investigators reviewing single-center series have not found this anatomic variant to be a risk factor.^{4,16} In the SVR cohort, only aortic atresia combined with mitral atresia was independently associated with interstage mortality. Aortic atresia with mitral stenosis was not associated with worsened interstage survival for the cohort as a whole or for either shunt group.

Single-center comparisons of subjects with the RVPAS versus the MBTS have reported no observable differences in myocardial performance and ventricular volumes before the stage II procedure.^{17,18} Although measurable differences in RVEF and RVESV indexed for BSA were observed between shunt types before discharge after the Norwood procedure, these echocardiographic findings were not independent risk factors for interstage mortality. Previous studies have identified postoperative moderate/ severe AVVR as a risk factor for interstage mortality. In the current analysis, shunt type was a predictor of interstage mortality when there was mild or no postoperative AVVR. However, when postoperative AVVR was moderate or greater, there was no selective benefit of the RVPAS and both shunt groups had higher mortality rates compared with subjects with mild or no AVVR and an RVPAS. These data suggest that the higher interstage mortality risk for the MBTS is not a consequence of greater AVVR. Moderate to severe AVVR does increase the risk of interstage mortality, possibly secondary to the additional volume load and ineffective ventricular output that can be exacerbated by conditions that elevate systemic vascular resistance.

We found that gestational age less than 37 weeks was an independent risk factor for interstage mortality. Given the high correlation of gestational age with birth weight, it is not surprising that birth weight was not found to be a separate independent risk factor. In a series comparing outcomes after cardiac surgery for term and preterm infants, complications of prematurity and postoperative mortality were common in preterm infants, illustrating the added vulnerability of preterm subjects.¹⁹ Other studies have identified extracardiac anomalies as a risk factor for mortality in subjects with HLHS.^{20,21} Extracardiac anomalies were not associated with increased risk in our cohort, but this may be confounded by a greater tendency for subjects with extracardiac malformations to die before hospital discharge. Furthermore, although use of an RVPAS has been advocated in the presence of

high-risk characteristics,¹³ our analysis did not demonstrate a particular protective effect of the RVPAS for preterm or low-birth-weight infants or for infants with extracardiac anomalies. Again, when considering this finding, it is important to consider that inclusion in this cohort required survival of the Norwood procedure and discharge from the hospital. Thus, subjects with these risks may be less likely to survive to hospital discharge after the Norwood procedure. Delivery at term seems to be an important and likely modifiable risk factor. This observation is consistent with a large series in which neonates aged less than 39 weeks with critical congenital heart disease had greater morbidity and interstage mortality than infants delivered at more than 39 weeks.²²

Having a higher number of postoperative complications correlated with longer length of stay and was an independent risk factor for interstage mortality. Longer length of stay in other series of infants with serious congenital heart disease has been associated with negative outcomes, including early mortality and lower scores on neurodevelopmental testing.²³

Poor interstage growth velocity and malnutrition are common after the Norwood procedure^{24,25}; thus, alternative feeding methods may be necessary to ensure adequate nutritional support. However, the best approach to feeding children after the Norwood procedure remains a challenge and there is wide variation in practice patterns. Univariate analysis showed that the absence of oral feeding before discharge after the Norwood procedure was associated with interstage mortality, particularly if nasogastric tube feedings were used rather than a direct gastrostomy tube. Poor feeding and failure to thrive in infants with congenital heart disease, specifically single ventricle disease, may be secondary to several mechanisms, such as decompensated heart failure, airway abnormalities, or gastrointestinal dysmotility. Thus, inability to feed orally may be a surrogate for added vulnerability in the infant with Norwood physiology. These data suggest that for subjects who require enteral tube feeding with complete absence of oral feeding, discharge with a direct gastrostomy tube may be preferable over a nasal enteral tube. The mechanism for interstage death in those with a nasogastric tube may include aspiration due to stenting of the lower esophageal sphincter, impaired breathing due to nasopharyngeal obstruction, or vagal response from reflux or gagging leading to bradycardia-induced arrhythmias.

Interstage mortality was associated with 2 sociodemographic variables: living in a census block with 5.4% to 13% below the poverty level and Hispanic ethnicity. Although the finding that participants in census blocks with lower or higher poverty levels had a lower risk of interstage mortality is not intuitive, it is possible that access to care was more limited in communities with 5.4% to 13% of residents below the poverty level. Although determining access to care was not within the scope of this study, it may be valuable to identify high-risk families with reduced access to nearby care to develop a support network and optimize interstage surveillance. Home monitoring with daily oxygen saturations and weight measurements, and with increased communication, has been associated with reduced interstage mortality in some series^{2,4,26} and may be of particular value when access to pediatric cardiac care is not readily available. Practices for home monitoring were not standardized in the SVR trial, and its impact on interstage mortality was not evaluated in this analysis.

Although this is the largest prospectively followed cohort of children with HLHS and other single right ventricle anomalies studied for risk factors for interstage mortality, there are limitations of our investigation. There was wide variation in practice throughout the course of this study across the clinical sites, including inpatient management for select patients with complicated post-Norwood hospitalization. Many variables evaluated in this study may be linked, and thus the implications of some practices cannot be defined. Furthermore, some variables deemed important even in the multivariable models may be only surrogates for other unmeasured variables. By adjusting for site and using bootstrapping, we have tried to minimize bias and maximize reliability.

Conclusions

We found higher interstage mortality in subjects who received an MBTS. However, RVPAS was only protective for interstage mortality in those subjects with none/mild AVVR after the Norwood procedure. Additional independent risk factors for interstage mortality included gestational age less than 37 weeks, Hispanic ethnicity, census block with 5.4% to 13% of inhabitants below poverty level, aortic atresia/mitral atresia, and increased number of post-Norwood complications. On the basis of our findings, we recommend avoiding preterm delivery when possible and considering close surveillance of infants discharged after the Norwood procedure with these identified risk factors.

Acknowledgments

Supported by Grants HL068269, HL068270, HL068279, HL068281, HL068285, HL068288, HL068290, HL068292, and HL085057 from the National Heart, Lung, and Blood Institute. This work is solely the responsibility of the authors and does not necessarily represent the official views of the National Heart, Lung, and Blood Institute.

Appendix 1

Pediatric Heart Network Investigators

National Heart, Lung, and Blood Institute: Gail Pearson, Victoria Pemberton, Rae-Ellen Kavey,* Mario Stylianou, Marsha Mathis*

Network Chair: *University of Texas Southwestern Medical Center,* Lynn Mahony

Data Coordinating Center: *New England Research Institutes,* Lynn Sleeper (PI), Sharon Tennstedt (PI), Steven Colan, Lisa Virzi,* Patty Connell,* Victoria Muratov, Lisa Wruck,* Minmin Lu, Dianne Gallagher, Anne Devine,* Julie Schonbeck, Thomas Trivison,* David F. Teitel

Core Clinical Site Investigators: *Children's Hospital Boston*, Jane W. Newburger (PI), Peter Laussen, Pedro del Nido, Roger Breitbart, Jami Levine, Ellen McGrath, Carolyn Dunbar-Masterson, John E. Mayer, Jr, Frank Pigula, Emile A. Bacha, Francis Fynn-Thompson; *Children's Hospital of New York*, Wymun Lai (PI), Beth Printz, *Daphne Hsu, William Hellenbrand, Ismee Williams, Ashwin Prakash, *Seema Mital, *Ralph Mosca, *Darlene Servedio, *Rozelle Corda, Rosalind Korsin, Mary Nash; *Children's Hospital of Philadelphia*, Victoria L. Vetter (PI), Sarah Tabbutt, *J. William Gaynor (Study Co-Chair), Chitra Ravishankar, Thomas Spray, Meryl Cohen, Marisa Nolan, Stephanie Piacentino, Sandra DiLullo, *Nicole Mirarchi; *Cincinnati Children's Medical Center*, D. Woodrow Benson (PI), Catherine Dent Krawczeski, Lois Bogenschutz, Teresa Barnard, Michelle Hamstra, Rachel Griffiths, Kathryn Hogan, Steven Schwartz, *David Nelson, Pirooz Eghtesady; *North Carolina Consortium: Duke University, East Carolina University, Wake Forest University*, Page A. W. Anderson (PI) – deceased, Jennifer Li (PI), Wesley Covitz, Kari Crawford, *Michael Hines, James Jagers, *Theodore Koutlas, Charlie Sang, Jr, Lori Jo Sutton, Mingfen Xu; *Medical University of South Carolina*, J. Philip Saul (PI), Andrew Atz, Girish Shirali, Scott Bradley, Eric Graham, Teresa Atz, Patricia Infinger; *Primary Children's Medical Center and the University of Utah*, L. LuAnn Minich (PI), John A. Hawkins, Michael Puchalski, Richard V. Williams, Peter C. Kouretas, Linda M. Lambert, Marian E. Shearrow, Jun A. Porter; *Hospital for Sick Children*, Brian McCrindle (PI), Joel Kirsh, Chris Caldarone, Elizabeth Radojewski, Svetlana Khaikin, Susan McIntyre, Nancy Slater; *University of Michigan*, Caren S. Goldberg (PI), Richard G. Ohye (Study Chair), Cheryl Nowak; *Children's Hospital of Wisconsin and Medical College of Wisconsin*, Nancy S. Ghanayem (PI), James S. Tweddell, Kathleen A. Mussatto, Michele A. Frommelt, Peter C. Frommelt, Lisa Young-Borkowski

Auxiliary Sites: *Children's Hospital Los Angeles*, Alan Lewis (PI), Vaughn Starnes, Nancy Pike; *The Congenital Heart Institute of Florida*, Jeffrey P. Jacobs (PI), James A. Quintessenza, Paul J. Chai, David S. Cooper, J. Blaine John, James C. Huhta, Tina Merola, Tracey Griffith; *Emory University*, William Mahle (PI), Kirk Kanter, Joel Bond, *Jeryl Huckaby; *Nemours Cardiac Center*, Christian Pizarro (PI), Carol Prospero; Julie Simons, Gina Baffa, Wolfgang A. Radtke; *University of Texas Southwestern Medical Center*, Ilana Zeltzer (PI), Tia Tortoriello, *Deborah McElroy, Deborah Town

Angiography Core Laboratory: *Duke University*, John Rhodes, J. Curt Fudge

Echocardiography Core Laboratories: *Children's Hospital of Wisconsin*, Peter Frommelt; *Children's Hospital Boston*, Gerald Marx

Genetics Core Laboratory: *Children's Hospital of Philadelphia*, Catherine Stolle

Protocol Review Committee: Michael Artman (Chair), Erle Austin, Timothy Feltes, Julie Johnson, Thomas Klitzner, Jeffrey Krischer, G. Paul Matherne

Data and Safety Monitoring Board: John Kugler (Chair), Rae-Ellen Kavey* (Executive Secretary), David J. Driscoll, Mark Galantowicz, Sally A. Hunsberger, Thomas J. Knight, Holly Taylor, Catherine L. Webb*

PI, Principal investigator.

* No longer at the institution listed.

Appendix Table 2

Candidate predictor list

Site
Shunt type
Baseline characteristics
Gestational age (wk)
Preterm (gestational age < 37 wk)
Birth weight (kg)
Low birth weight (<2500 g)
Race (white, black, other)
Hispanic
Anatomic diagnosis (HLHS, single right ventricle with systemic outflow, straddling mitral valve with left ventricle hypoplasia and outflow obstruction, other)
HLHS (yes/no)
Heterotaxy (yes/no)
Aortic atresia/mitral atresia (yes/no)
Aortic atresia/mitral stenosis (yes/no)

Aortic stenosis/mitral stenosis (yes/no)
 Obstructed pulmonary venous return (yes/no)
 Age at Norwood (d)
 US Census Socioeconomic Index Score
 US Census % below poverty level
 Norwood hospitalization
 Total bypass time (min)
 Perfusion strategy during Norwood: DHCA only, RCP only or RCP with DHCA no > 10 min, RCP and DHCA time > 10 min)
 ECMO (yes/no)
 Ultrafiltration during (yes/no)
 Aortic atresia (yes/no)
 Steroids (yes/no)
 Trasylol (yes/no)
 Alpha-blockade (yes/no)
 CPR post-Norwood during hospital stay (yes/no)
 Oxygen saturation at discharge post-Norwood (yes/no)
 Norwood length of stay
 No. of unintended cardiovascular procedures
 Balloon dilation, stent placement, or surgical revision of the neo-aorta (yes/no)
 Pulmonary artery reconstruction (yes/no)
 Balloon dilation or stenting of the shunt or branch pulmonary arteries (yes/no)
 Shunt revision or crossover (yes/no)
 Atrioventricular valve repair (yes/no)
 Diaphragm plication (yes/no)
 Gastrostomy tube (yes/no)
 Pericardial window (yes/no)
 Shunt revision (no crossover)
 Thoracostomy tube (yes/no)
 No. of complications pre-discharge, post-Norwood
 Any serious adverse event (yes/no)
 Any oral feeding at discharge (yes/no)
 Only oral feeding at discharge (yes/no)
 Feeding category (gastrostomy or gastrostomy-jejunal tube, nasal gastric or nasal jejunal tube, only oral)
 Feeding category (no oral, G or GJ tube; no oral, NG or NJ tube; oral, G or GJ tube; oral, NG or NJ tube; only oral)
 No. of discharge medications
 Echocardiography pre-Norwood
 RVEDV indexed to BSA^{1,3}
 RVESV indexed to BSA^{1,3}
 RVEF
 Atrioventricular valve insufficiency (moderate to severe AVVR) at pre-Norwood
 Ascending aorta diameter (mm)
 Echocardiography post-Norwood
 Echo Nakata Index

Aortic valve insufficiency (not patent, no regurgitation, regurgitation)
 Atrioventricular valve insufficiency at post-Norwood (none/mild vs moderate/severe).
 Right ventricular end-diastolic volume indexed to BSA^{1,3}
 Right ventricular end-systolic volume indexed to BSA^{1,3}
 RVEF
 Right ventricular percent area change
 Reversal of pulmonary vein flow (yes/no)
 Genetic evaluations (yes at any visit)
 Genetic syndrome (yes/no)
 Any nonsyndromic abnormality (yes/no)
 Genotype (E3/E3, E4/E4, E2/E3, E2/E4, E3/E4)
 Site volume and surgeon Norwood volume
 Surgeon Norwood experience, based on all screened subjects (continuous and 4-level categories)
 Site volume, based on annual center volume of Norwood procedures (continuous and 4-level categories)

HLHS, Hypoplastic left heart syndrome; *DHCA*, deep hypothermic circulatory arrest; *RCP*, regional cerebral perfusion; *ECMO*, extracorporeal membrane oxygenation; *CPR*, cardiopulmonary resuscitation; *G*, gastrostomy, *GJ*, gastric-jejunal; *NG*, naso-gastric; *NJ*, nasojejunal; *RVEDV*, right ventricular end-diastolic volume; *BSA*, body surface area; *RVESV*, right ventricular end-systolic volume; *RVEF*, right ventricular ejection fraction; *AVVR*, atrioventricular valve regurgitation.

Appendix Table 3

Univariate comparisons of subjects who were discharged after Norwood procedure with those who were not discharged and survived to stage II

Characteristic	Discharged (N = 426)	Not discharged and survived (N = 22)	P value
Shunt			.52
RVPAS	225 (53)	10 (46)	
MBTS	201 (47)	12 (55)	
Norwood hospitalization Norwood length of stay (d)			
Median (IQR)	23 (16-36)	115 (67-147)	<.001
Total bypass time (min)			
Median (IQR)	130.0 (101.0-163.0)	152.0 (122.0-184.0)	.03
Placed on ECMO			<.001
No	402 (94)	13 (59)	
Yes	24 (5.6)	9 (41)	
Ultrafiltration during CPB			.02
No	161 (38)	3 (14)	
Yes	265 (62)	19 (86)	
Steroids			.005
No	40 (9.4)	7 (32)	
Yes	386 (91)	15 (68)	

Characteristic	Discharged (N = 426)	Not discharged and survived (N = 22)	P value
CPR post-Norwood during hospital stay			<.001
No	397 (93)	10 (45)	
Yes	29 (6.8)	12 (55)	
No. of unintended cardiovascular procedures			<.001
0	360 (85)	12 (55)	
1	56 (13)	6 (27)	
2	10 (2.3)	4 (18)	
Balloon dilation or stenting of the shunt or branch pulmonary arteries			.02
No	408 (96)	18 (82)	
Yes	18 (4.2)	4 (18)	
Shunt revision or crossover			.04
No	378 (89)	16 (73)	
Yes	48 (11)	6 (27)	
Shunt revision without crossover			.002
No	411 (96)	17 (77)	
Yes	15 (3.5)	5 (23)	
PA reconstruction			.05
No	426 (100)	21 (95)	
Yes	0 (0.0)	1 (4.5)	
Gastrostomy tube			.002
No	357 (84)	12 (55)	
Yes	69 (16)	10 (45)	
Thoracostomy tube			.03
No	404 (95)	18 (82)	
Yes	22 (5)	4 (18)	
Complications and serious adverse events			
No. of complications			
Median (IQR)	2.0 (1.0-3.0)	6.0 (3.0-15.0)	<.001
No. of SAEs			
Median (IQR)	0.0 (0.0-0.0)	1.0 (0.0-2.0)	<.001
Any SAE			<.001
0	362 (85)	9 (41)	
1	64 (15)	13 (59)	
Feeding at discharge Any oral feeding?			
No	91 (21)	18 (82)	
Yes	335 (79)	4 (18)	
Only oral feeding?			<.001
No	286 (67)	22 (100)	
Yes	140 (33)	0 (0)	
Feeding category (3 df)			<.001
Any oral, other tube	0 (0.0)	1 (5)	

Characteristic	Discharged (N = 426)	Not discharged and survived (N = 22)	P value
Any oral, G or GJ tube	76 (18)	12 (55)	
Any oral, NG or NJ tube	210 (49)	9 (41)	
Only oral	140 (33)	0 (0)	
Feeding category (4 df)			<.001
No oral, other tube	0 (0)	1 (5)	
No oral, G or GJ tube	55 (13)	10 (45)	
No oral, NG or NJ tube	36 (8)	7 (32)	
Oral and G or GJ tube	21 (5)	2 (9)	
Oral and NG or NJ tube	174 (41)	2 (9)	
Only oral	140 (33)	0 (0)	
Echocardiography pre-Norwood			
AVVR			.04
None/mild	368 (88)	16 (73)	
Moderate/severe	49 (12)	6 (27)	
post-Norwood			
Akata Index			
Median (IQR)	123.3 (95.5-159.0)	157.1 (120.4- 223.6)	.006
AVVR			.003
None/mild	329 (78)	10 (48)	
Moderate/severe	94 (22)	11 (52)	
Death during stage II hospitalization among subjects surviving to stage II			
No	367 (98)	14 (64)	<.001
Yes	9 (2)	8 (36)	

With the exception of shunt type, only differences in characteristics between discharged and nondischarged subjects with $P < .05$ are presented. Fisher exact test was used to compare subject groups for variables that are dichotomous or categorical. The Wilcoxon rank-sum test was used to compare subject groups for continuous variables that are skewed. Student 2-sample t test was used for continuous variables that are not skewed. *RVPAS*, Right ventricle-to-pulmonary artery shunt; *MBTS*, modified Blalock-Taussig shunt; *IQR*, interquartile range; *ECMO*, extracorporeal membrane oxygenation; *CPB*, cardiopulmonary bypass; *CPR*, cardiopulmonary resuscitation; *PA*, pulmonary artery; *SAE*, serious adverse event; *G*, gastrostomy; *GJ*, gastric-jejunal; *NG*, nasogastric; *NJ*, nasojejunal; *AVVR*, atrioventricular valve regurgitation.

References

1. Simsic JM, Bradley SM, Stroud MR, Atz AM. Risk factors for interstage death after the Norwood procedure. *Pediatr Cardiol*. 2005; 26:400–3. [PubMed: 16374690]
2. Ghanayem NS, Tweddell JS, Hoffman GM, Mussatto K, Jaquiss RD. Optimal timing of the second stage of palliation for hypoplastic left heart syndrome facilitated through home monitoring, and the results of early cavopulmonary anastomosis. *Cardiol Young*. 2006; 16(Suppl I):60–5.
3. Hehir DA, Dominguez TE, Ballweg JA, Ravishankar C, Marino BS, Bird GL, et al. Risk factors for interstage death after stage I reconstruction of hypoplastic left heart syndrome and variants. *J Thorac Cardiovasc Surg*. 2008; 136:94–9. [PubMed: 18603060]
4. Furck AK, Uebing A, Hansen JH, Scheewe J, Jung O, Fischer G, et al. Outcome of the Norwood operation in patients with hypoplastic left heart syndrome: a 12-year single-center survey. *J Thorac Cardiovasc Surg*. 2010; 139:359–65. [PubMed: 19879598]

5. Jonas RA, Hansen DD, Cook N, Wessel D. Anatomic subtype and survival after reconstructive operation for hypoplastic left heart syndrome. *J Thorac Cardiovasc Surg.* 1994; 107:1121–8. [PubMed: 7512677]
6. Bartram U, Grunenfelder J, Van Praagh R. Causes of death after the modified Norwood procedure: a study of 122 postmortem cases. *Ann Thorac Surg.* 1997; 64:1795–802. [PubMed: 9436575]
7. Tabbutt S, Dominguez TA, Ravishankar C, Marino BS, Gruber PJ, Wernovsky G, et al. Contemporary comparison of outcomes following use of the right ventricular to pulmonary artery conduit vs the modified Blalock Taussig shunt as part of the modified Norwood procedure. *Ann Thorac Surg.* 2005; 80:1582–90. [PubMed: 16242421]
8. Cua CL, Thiagarajan RR, Gauvreau K, Lai L, Costello JM, Wessel DL, et al. Early postoperative outcomes in a series of infants with hypoplastic left heart syndrome undergoing stage 1 palliation operation with either modified Blalock-Taussig shunt or right ventricle to pulmonary artery conduit. *Pediatr Crit Care Med.* 2006; 7:238–44. [PubMed: 16474256]
9. Ohye RG, Sleeper LA, Mahony L, Newburger JW, Pearson GD, Goldberg CS, et al. Comparison of shunt types in the Norwood procedure for single ventricle lesions. *N Engl J Med.* 2010; 362:1980–92. [PubMed: 20505177]
10. Ohye RG, Gaynor JW, Ghanayem NS, Goldberg CS, Laussen PC, Frommelt PC, et al. Design and rationale of a randomized trial comparing the Blalock-Taussig and right ventricle-pulmonary artery shunts in the Norwood procedure. *J Thorac Cardiovasc Surg.* 2008; 136:968–75. [PubMed: 18954638]
11. Virzi L, Pemberton V, Ohye RG, Tabbutt S, Lu M, Atz TC, et al. Reporting adverse events in a surgical trial for complex congenital heart disease: The Pediatric Heart Network experience. *J Thorac Cardiovasc Surg.* 2011; 142:531–7. [PubMed: 21397260]
12. Singh TP, Gauvreau K, Bastardi HJ, Blume ED, Mayer JE. Socioeconomic position and graft failure in pediatric heart transplant recipients. *Circ Heart Fail.* 2009; 2:160–5. [PubMed: 19808335]
13. Sano S, Huang SC, Kasahara S, Yoshizumi K, Kotani Y, Ishino K. Risk factors for mortality after the Norwood procedure using the right ventricle to pulmonary artery shunt. *Ann Thorac Surg.* 2009; 87:178–86. [PubMed: 19101293]
14. Bradley SM, Simsic JM, McQuinn TC, Habib DM, Shirali GS, Atz AM. Hemo-dynamic status after the Norwood procedure: a comparison of right ventricle-to-pulmonary artery connection versus modified Blalock-Taussig shunt. *Ann Thorac Surg.* 2004; 78:933–41. [PubMed: 15337023]
15. Ghanayem NS, Jaquiss RD, Cava JR, Frommelt PC, Mussatto KA, Hoffman GM, et al. Right ventricle-to-pulmonary artery conduit versus Blalock-Taussig shunt: a hemodynamic comparison. *Ann Thorac Surg.* 2006; 82:1603–10. [PubMed: 17062213]
16. Polimenakos AC, Sathanandam SK, Hussayni TS, El Zein CF, Roberson DA, Ilbawi MN. Hypoplastic left heart syndrome and aortic atresia-mitral stenosis variant. Role of myocardial protection strategy and impact of ventriculo-coronary connections after stage 1 palliation. *Pediatr Cardiol.* 2011; 32:929–39. [PubMed: 21644013]
17. Frommelt PC, Sheridan DC, Mussatto KM, Hoffman GM, Ghanayem NS, Frommelt MA, et al. Effect of shunt type on echocardiographic indices after initial palliation for hypoplastic left heart syndrome: Blalock-Taussig shunt versus right ventricle-pulmonary artery conduit. *J Am Soc Echocardiogr.* 2007; 20:1364–73. [PubMed: 17604955]
18. Graham EM, Atz AM, Bradley SM, Scheurer MA, Bandisode VM, Laudito A, et al. Does a ventriculotomy have deleterious effects following palliation in the Norwood procedure using a shunt placed from the right ventricle to the pulmonary arteries? *Cardiol Young.* 2007; 17:145–50. [PubMed: 17244384]
19. Natarajan G, Anne SR, Aggarwal S. Outcomes of congenital heart disease in late pre-term infants: double jeopardy? *Acta Paediatr.* 2011; 100:1104–7. [PubMed: 21362036]
20. Gaynor JW, Mahle WT, Cohen MI, Ittenbach RF, DeCampi WM, Steven JW, et al. Risk factors for mortality after the Norwood procedure. *Eur J Cardiothorac Surg.* 2002; 22:82–9. [PubMed: 12103378]

21. Stasik CN, Gelehrter S, Goldberg CS, Bove EL, Devaney EJ, Ohye RG. Current outcomes and risk factors for the Norwood procedure. *J Thorac Cardiovasc Surg.* 2006; 131:412–7. [PubMed: 16434272]
22. Costello JM, Polito A, Brown DW, McElrath TF, Graham DA, Thiagarajan RR, et al. Delivery before 39 weeks is associated with adverse outcomes in neonates with cardiac disease. *Pediatrics.* 2010; 126:277–84. [PubMed: 20603261]
23. Newburger JW, Wypij D, Bellinger DC, du Plessis AJ, Kuban KC, Rappaport LA, et al. Length of stay after infant heart surgery is related to cognitive outcome at age 8 years. *J Pediatr.* 2003; 143:67–73. [PubMed: 12915826]
24. Kelleher DK, Laussen P, Teixeira-Pinto A, Duggan C. Growth and correlates of nutritional status among infants with hypoplastic left heart syndrome (HLHS) after stage 1 Norwood procedure. *Nutrition.* 2006; 22:237–44. [PubMed: 16500550]
25. Anderson JB, Beekman RH 3rd, Border WL, Kalkwarf HJ, Khoury PR, Uzark K, et al. Lower weight-for-age z score adversely affects hospital length of stay after the bidirectional Glenn procedure in 100 infants with a single ventricle. *J Thorac Cardiovasc Surg.* 2009; 138:397–404. [PubMed: 19619784]
26. Ghanayem NS, Hoffman GM, Mussatto KA, Cava JR, Frommelt PC, Rudd NA, et al. Home surveillance program prevents interstage mortality after the Norwood procedure. *J Thorac Cardiovasc Surg.* 2003; 126:1367–77. [PubMed: 14666008]

Abbreviations and Acronyms

AVVR	atrioventricular valve regurgitation
BSA	body surface area
CPR	cardiopulmonary resuscitation
DHCA	deep hypothermic circulatory arrest
ECMO	extracorporeal membrane oxygenation
HLHS	hypoplastic left heart syndrome
MBTS	modified Blalock–Taussig shunt
OR	odds ratio
RVEDV	right ventricular end-diastolic volume
RVEF	right ventricular ejection fraction
RVESV	right ventricular end-systolic volume
RVPAS	right ventricle-to-pulmonary artery shunt
SVR	Single Ventricle Reconstruction




Figure 1.

Subject participation status from randomization through the interstage period. A total of 430 subjects were discharged after the Norwood procedure. Four of these subjects were excluded because of an unusually late or missing date of the stage II procedure.




Figure 2. Kaplan-Meier plot of interstage survival by shunt type: Interstage survival, defined as percentage alive postdischarge after the Norwood procedure and before the stage II surgery, was significantly better for subjects with an RVPAS than for subjects with an MBTS. *MBTS*, Modified Blalock–Taussig shunt; *RVPAS*, right ventricle-to-pulmonary artery shunt.

Table 1

Associations of interstage death, site adjusted

Variables	n	Responses	Interstage survivals (N = 376)		Interstage deaths (N = 50)		Site adjusted	
			df	OR	P	OR	P	
Site	426	-	-	-	-	-	14	
Shunt	426		376		50		1	<.001
		RVPAS	212 (56)		13 (26)			
		MBTS	164 (44)		37 (74)			3.44
Demographic characteristics								
Hispanic	421		371		50		1	.07
		No	305 (82)		33 (66)			-
		Yes	66 (18)		17 (34)			1.99
US census % below poverty level								
Tertiles	409		364		45		2	.006
		<5.42%	131 (36)		5 (11)			0.38
		5.42%-13%	113 (31)		23 (51)			1.73
		13%	120 (33)		17 (38)			-
US census socioeconomic score	409		0.32 ± 5.1		-1.4 ± 4.4		1	0.95
Baseline characteristics								
Gestational age (wk)	426		38.3 ± 1.5		37.7 ± 1.9		1	0.80
Preterm (gestational age < 37 wk)	426		376		50		1	.01
		37 wk	344 (91)		40 (80)			-
		<37 wk	32 (9)		10 (20)			2.81
Birth weight (kg)	426		3.2 (0.5)		3.0 (0.5)		1	0.57
AA/MA	426		376		50		1	.20
		No	235 (62)		26 (52)			-
		Yes	141 (38)		24 (48)			1.49
Norwood hospitalization								
Age at Norwood (d)	426		5.6 ± 3.7		6.5 ± 5.1		1	1.06
Norwood length of stay (d)	426		28.9 ± 19.0		36.2 ± 33.3		1	1.01
Perfusion strategy	422		373		49		2	.19

Variables	n	Responses	Interstage survivals (N = 376)		Interstage deaths (N = 50)		Site adjusted	
							OR	P
DHCA only			211 (57)		28 (57)		0.34	
RCP only or RCP/DHCA time < 10 min			91 (24)		8 (16)		0.38	
RCP and DHCA time > 10 min			71 (19)		13 (27)		-	
Complications and serious adverse events during Norwood hospitalization								
No. of complications post-Norwood	426		2.08 ± 2.1		3.58 ± 3.4	1	1.22	<.001
No. of medications at discharge	426		4.9 ± 1.7		5.6 ± 2.2	1	1.15	.098
Feeding status at discharge post-Norwood								
Any oral feeding	426		376		50	1	-	<.001
		No	71 (19)		20 (40)		-	
		Yes	305 (81)		30 (60)		0.28	
Feeding category (4 df)	426		376		50	4	-	<.001
No oral, G or GJ tube			47 (13)		8 (16)		1.68	
No oral, NG or NJ tube			24 (6)		12 (24)		8.40	
Oral and G or GJ tube			19 (5)		2 (4)		1.52	
Oral and NG or NJ tube			159 (42)		15 (30)		0.88	
Only oral			127 (34)		13 (26)		-	
Echocardiography pre-Norwood								
AVVR	417		370		47	1	-	.004
		None/mild	333 (90)		35 (74)		-	
		Moderate/severe	37 (10)		12 (26)		3.15	
Ascending aorta diameter (R301), cm pre-Norwood	414		368		46	1	-	.008
Median cutoff	414		172 (47)		30 (65)		2.44	
		<0.3 cm	196 (53)		16 (35)		-	
		0.3 cm						
Echocardiography post-Norwood								
RV end-diastolic volume/BSA ^{1.3} (mL/m ^{2.6})	330		91.2 ± 25.2		98.3 ± 32.8	1	1.01	.14
RV end-systolic volume/BSA ^{1.3} (mL/m ^{2.6})	330		48.7 ± 17.0		56.6 ± 23.2	1	1.02	.03
RVEF (%)	330		47.3 ± 8.0		43.2 ± 7.3	1	0.94	.01
RV fractional area change (%)	409		37.0 ± 7.7		34.9 ± 6.8	1	0.97	.12
AVVR grade	423		374		49	1	-	.06

Table 2
Multivariable model of predictors of interstage mortality with site adjustment: N = 399
(adjusted $R^2 = 0.35$)

Predictor	Died/heart transplant?		Odds ratio (95% CI)	Model P value
	No	Yes		
Site	–	–	–	.823
AVVR grade by shunt			–	.03
None/mild				
MBTS	117 (84)	23 (16)	9.7 (3.1-30.3)	Contrast <.001
RVPAS	165 (97)	5 (2.9)	–	
Moderate/severe				
MBTS	33 (79)	9 (21)	1.4 (0.4-5.0)	Contrast .57
RVPAS	39 (83)	8 (17)		
Low gestational age (<37 wk)				
Yes	30 (75)	10 (25)	3.9 (1.4-10.8)	.008
No	324 (90)	35 (10)		
Hispanic				
Yes	63 (79)	17 (21)	2.6 (1.1-6.5)	.04
No	291 (91)	28 (8.7)		
AA/MA				
Yes	134 (85)	23 (15)	2.3 (1.1-5.0)	.03
No	220 (91)	22 (9.1)		
No. of complications post-Norwood and pre-discharge				
Mean ± SD	2.0 ± 2.0	3.2 ± 3.3	1.2 (1.1-1.4)	.006
US Census percentage below poverty level tertiles				.003
<5.42	128 (96)	5 (4)	0.38 (0.12-1.3)	.15 (0.05-0.48)*
5.42-13.0	109 (83)	23 (17)	2.5 (1.02-6.0)	–
13.0	117 (87)	17 (13)	–	

Site 19 was excluded from this analysis because all 3 subjects from this site survived and render the model unstable. *CI*, Confidence interval; *AVVR*, atrioventricular valve repair; *MBTS*, modified Blalock–Taussig shunt; *RVPAS*, right ventricle-to-pulmonary artery shunt; *AA/MA*, aortic atresia/mitral atresia; *SD*, standard deviation.

* Odds of mortality/transplant among subjects in the < 5.42 tertile compared with subjects in the 5.42-13.0 tertile.