



Exploring the relationships between neuronal parameters and network activity

Anca Doloc Mihi, *Emory University*
[Ronald Calabrese](#), *Emory University*

Journal Title: BMC Neuroscience
Volume: Volume 13, Number Suppl 1
Publisher: BioMed Central | 2012-07-16, Pages P172-P172
Type of Work: Article | Final Publisher PDF
Publisher DOI: 10.1186/1471-2202-13-S1-P172
Permanent URL: <http://pid.emory.edu/ark:/25593/cw2kq>

Final published version: <http://dx.doi.org/10.1186%2F1471-2202-13-S1-P172>

Copyright information:

©2012 Doloc-Mihu and Calabrese; licensee BioMed Central Ltd.
This is an Open Access work distributed under the terms of the Creative Commons Attribution 2.0 Generic License (<http://creativecommons.org/licenses/by/2.0/>).



Accessed December 9, 2021 4:27 AM EST

POSTER PRESENTATION

Open Access

Exploring the relationships between neuronal parameters and network activity

Anca Doloc-Mihu*, Ronald L Calabrese

From Twenty First Annual Computational Neuroscience Meeting: CNS*2012
Decatur, GA, USA. 21-26 July 2012

This study constitutes our next step toward a full investigation of how intrinsic membrane and synaptic parameters affect the electrical activity of a half-center oscillator (HCO) model and how different parameter regimes influence stability and modulatability of the HCO model's output. In our previous study [3], we constructed an efficient relational database table with the resulting characteristics of Hill et al.'s [2] HCO single-compartment conductance-based model. This model consists of two reciprocally inhibitory neurons and replicates the electrical activity of the oscillator interneurons of the leech heart-beat CPG under a variety of experimental conditions.

We systematically [4] explored the parameter space of this HCO model by varying a set of eight selected parameters (maximal conductances of intrinsic and synaptic currents) in all combinations possible, resulting in 10,485,760 simulated model instances (10,321,920 HCO instances and 163,840 corresponding isolated neuron instances). We classified within a group the simulated instances showing similar electrical activity. The HCOs were split into ten groups labeled: *spiking*, *silent*, *asymmetric activity*, *plateau*, *irregular spikes*, *asymmetric bursting*, *one burst*, *irregular period*, *unbalanced*, and *functional* (1,202,139 instances); the isolated neuron instances were split into seven groups: *spiking*, *silent*, *bistable*, *plateau*, *irregular spikes*, *irregular period*, and *regular bursting* [3].

By querying the database we started analyzing the relationships between the neuronal parameters of the model instances within the same group and across groups. Model instances whose neurons are synaptically isolated are called isolated neuron instances. We found that *regularly bursting* (isolated) neuron instances compose only a small minority of *functional* HCOs (424 instances); the vast majority was composed of *spiking* neurons (32,568

instances). Also, our data suggest that *regular bursting* isolated neurons form *functional* HCOs that are more robust to variations in synaptic parameters than those formed by *spiking* isolated neurons. Approximately 8.2% (99,066 instances) of the functional HCOs have bursting characteristics that are similar to those of the living animal [1].

We used dimensional stacking (the NDVis tool) to visualize the distribution of each group of model instances versus the neuronal parameters. While each group has its own distribution within the parameter space, the groups of bistable and regular bursting isolated neuron instances separate large zones of spiking and silent model instances.

To explore in depth the interaction of parameters that lead to the different activity groups we have identified, we used the Principal Component Analysis (PCA) method. When we applied the PCA to the group of regular bursting isolated neuron instances, we found that only 4 (out of 6) components are important for this group (sum of variances > 95%) and that the 1st component has the greatest impact (variance > 60%). This component is mostly influenced by the maximal conductances of NaP, K₂ and Leak, and the E_{Leak} (linear combination of the parameters). Then, we applied PCA to our group of functional HCOs showing realistic leech bursting characteristics and we found that the first 6 (out of 8) are the most important principal components for this group (sum of variances > 96%). However, unlike in the regular bursting group, none of these principal components is dominant. This suggests that this group has complex non-linear relationships between its parameters and to explore them we need to use more sophisticated computational methods than the (linear) PCA.

* Correspondence: adolocm@emory.edu
Department of Biology, Emory University, Atlanta, GA 30322, USA

Acknowledgements

Work supported by the National Institute Health Grant NS024072 to R.L. Calabrese.

Published: 16 July 2012

References

1. Cymbalyuk GS, Gaudry Q, Masino MA, Calabrese RL: **Bursting in leech heart interneurons: cell-autonomous and network-based mechanisms.** *J Neurosci* 2002, **22**:10580-10592.
2. Hill AAV, Lu J, Masino MA, Olsen H, Calabrese RL: **A model of a segmental oscillator in the leech heartbeat neuronal network.** *J Comp. Neurosci* 2001, **10**:281-302.
3. Doloc-Mihu A, Calabrese RL: **Bringing up the rear: new premotor interneurons add regional complexity to a segmentally distributed motor pattern.** *J Biol Physics* 2011, **37**:263-283.
4. Prinz AA, Billimoria CP, Marder E: **Alternative to hand-tuning conductance-based models: construction and analysis of databases of model neurons.** *J Neurophysiol* 2003, **90**:3998-4015.

doi:10.1186/1471-2202-13-S1-P172

Cite this article as: Doloc-Mihu and Calabrese: Exploring the relationships between neuronal parameters and network activity. *BMC Neuroscience* 2012 **13**(Suppl 1):P172.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

