



EMORY
LIBRARIES &
INFORMATION
TECHNOLOGY

OpenEmory

Commonalities and Differences in Correlates of Depressive Symptoms in Men and Women with Heart Failure

Jo-Ann Eastwood, *University of California at Los Angeles*
Debra K. Moser, *University of Kentucky*
Barbara J. Riegel, *University of Pennsylvania*
Nancy M. Albert, *Cleveland Clinic*
Susan Pressler, *University of Michigan*
Misook L. Chung, *University of Kentucky*
[Sandra B Dunbar](#), *Emory University*
Jia-Rong Wu, *University of North Carolina*
Terry A. Lennie, *University of Kentucky*

Journal Title: European Journal of Cardiovascular Nursing
Volume: Volume 11, Number 3
Publisher: SAGE Publications (UK and US) | 2012-09-01, Pages 356-365
Type of Work: Article | Post-print: After Peer Review
Publisher DOI: 10.1177/1474515112438010
Permanent URL: <https://pid.emory.edu/ark:/25593/sq6v9>

Final published version: <http://dx.doi.org/10.1177/1474515112438010>

Copyright information:

© The European Society of Cardiology 2012.

Accessed October 22, 2019 5:05 AM EDT



Published in final edited form as:

Eur J Cardiovasc Nurs. 2012 September ; 11(3): 356–365. doi:10.1177/1474515112438010.

Commonalities and Differences in Correlates of Depressive Symptoms in Men and Women with Heart Failure

Jo-Ann Eastwood, PhD¹, Debra K. Moser, DNSc², Barbara J. Riegel, DNSc³, Nancy M. Albert, PhD⁴, Susan Pressler, PhD⁵, Misook L. Chung, PhD², Sandra Dunbar, DNS⁶, Jia-Rong Wu, PhD⁷, and Terry A. Lennie, PhD²

¹University of California School of Nursing

²College of Nursing, University of Kentucky

³School of Nursing, University of Pennsylvania

⁴Nursing Research and Innovation, Cleveland Clinic

⁵University of Michigan School of Nursing

⁶Emory University

⁷University of North Carolina at Chapel Hill, School of Nursing

Heart failure (HF) is a major cause of excess morbidity and mortality, and a leading contributor to healthcare expenditures in the U.S. with estimated direct and indirect costs of greater than 37.2 billion dollars per year.¹ Direct costs are mainly attributed to repeat hospitalizations (75%), and on average are approximately €10,000 per patient per hospitalization in Europe.² Depressive symptoms are common in outpatients with HF with a prevalence of 13–42%,^{3,4–6} and the presence of depressive symptoms is linked to poorer clinical outcomes in patients with HF.^{3,7–9} Patients with HF and depressive symptoms commonly suffer from poorer quality of life, declining functional status, greater symptom burden, poorer adherence, more frequent rehospitalizations, and worse survival.^{3,8,10–12} Given the common finding that women experience higher levels of depression than men,¹³ a major step to understanding depression in patients with HF is determining how gender contributes to the depression experience.

Gender contributes to differences in clinical presentation and outcomes in patients with HF,¹⁴ yet most research on depressed patients with HF does not examine gender differences despite the increased prevalence of depressive symptoms in women.¹³ Consequently, it is impossible to determine if women and men with HF differ on important characteristics such as demographics, behavioral, clinical, and psychosocial variables that may influence depression. Gender differences among patients with heart failure must be considered when developing targeted, preventive treatment strategies in order to address healthcare costs and

Address for correspondence: Jo-Ann Eastwood PhD, RN, CCNS, ACNP-BC, FAHA, Assistant Professor, UCLA School of Nursing, University of California, Los Angeles, 700 Tiverton Blvd., Factor Building-4-254, Los Angeles, CA. 90095-6918, jeastwoo@sonnet.ucla.edu, 310-206-3443.

The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institute of Nursing Research or the National Institutes of Health

societal burden related to the increasing numbers of men and women with HF.¹⁵ Therefore the aims of this study were to 1) compare the prevalence and severity of depressive symptoms between men and women enrolled in a large HF registry and 2) to determine gender differences in predictors of depressive symptoms from among relevant demographic (i.e., age, ethnicity, marital status, education, and financial status), behavioral (i.e., smoking and exercise), clinical (i.e., HF etiology, functional status, functional capacity, comorbidity burden, and body mass index), and psychosocial (i.e., anxiety, perceived control, and health perceptions) factors.

REVIEW OF RELEVANT LITERATURE

Heart failure (HF) patients with depressive symptoms have poorer clinical outcomes,^{3, 7, 8, 10, 11, 16–19} lower quality of life and substantially higher health care costs than HF patients without such symptoms.^{3, 8, 11, 12} A recent meta analysis demonstrated that patients with HF who have depressive symptoms are more than twice as likely to die or experience a cardiac event compared to HF patients without depressive symptoms.³ Although depression is more prevalent in women overall and in cardiac disease in general,²⁰ investigators recently reported finding more male than female depression sufferers in a large cohort of patients with HF.²¹ However, little is known about the role of gender in comorbid HF and depression. Given the importance of depression in HF, further investigation is needed to ascertain how gender interacts with HF and depressive symptoms to influence clinical outcomes.

Previous studies have reported demographic factors which are related to HF and depression. The incidence of depression is inversely related to age.²² Patients with HF manifest greater depressive symptoms when they are younger.^{7, 8, 23, 24} Other sociodemographic risk factors for depression among patients with HF include low educational attainment, being unmarried or living alone, and being unemployed or having a low income.^{17, 25–27} In comorbid depression and HF, ethnicity prevalence data have been mixed, with one large study reporting a higher rate of depression history in whites¹⁶ and a smaller study finding that non-Hispanic Blacks were more likely to be depressed than others.²⁸

Biological and behavioral factors are likely linked to both depression and HF. For example, higher NYHA functional class is associated with higher prevalence of depression³ and may be a reflection of higher illness burden from HF that contributes to depression.¹⁵ Unhealthy behaviors such as alcohol and tobacco use, poor nutrition, sedentary lifestyle and poor adherence to medical regimens are associated with depression in HF patients.^{29, 30}

Patients with chronic illnesses such as HF experience increased mental distress.^{31, 32} Symptoms of anxiety are highly prevalent in HF patients.^{28, 33, 34} Anxiety symptoms overlap significantly with depressive symptoms and increase the risk of mortality and functional disability as well as health care utilization and cost.^{8, 35, 36} In addition to anxiety, perceptions of health status, functional capacity, and perceived control influence the incidence and severity of depressive symptoms^{37–39} and are linked to poorer outcomes.^{40, 41} Higher levels of perceived control are linked with fewer anxious and depressive symptoms and a higher likelihood of patient involvement in self-care management, which is related to

improved outcomes in HF patients.^{28, 38, 42} HF patients who indicate low perceived health status are more prone to develop depressive symptoms.¹⁷

Previous studies of depression and HF have included measures of sociodemographic, clinical, behavioral and psychosocial risk factors.^{7, 17, 29, 43–45} A comprehensive multivariate examination has not, however, been conducted and few studies have reported comparisons by gender.^{7, 10}

METHODS

Data from the HF Quality of Life Collaborative,^{46–48} an ongoing HF health-related quality of life registry, were used to meet the study aims. A total of 622 patients in the registry had complete data on all variables.

Adult patients over age 18 were enrolled from cardiology clinics associated with academic medical centers in the Midwest, Northeast, South and Southeast of the United States. Patients were included if they had a documented diagnosis from a cardiologist of chronic HF from either preserved or non-preserved ejection fraction and were community dwelling (i.e., not residing in an extended care or skilled nursing facility). Patients were excluded if they had acute myocardial infarction or stroke within the prior 3 months or were cognitively impaired as determined by inability to answer questions from the researcher, complete questionnaires, or provide informed consent. All participants provided written informed consent, and all participating sites obtained institutional review board approval.

Measures

Demographic variables—Data on age, gender, ethnicity, education level, marital status, and financial status were collected by patient self-reported questionnaires. Because participants commonly are reluctant to provide income data, we used an accepted alternative approach in which patients were asked to respond to the question, “Considering your household income, would you say you...” with the following three options: 1) are comfortable; 2) have enough to make ends meet; or 3) do not have enough to make ends meet?” This method reflects the impact of a participant’s socioeconomic status on daily life.⁴⁹

Behavioral and clinical variables—Data on smoking and exercise were collected using patient questionnaires. Body mass index (BMI) was calculated from height and weight available by medical record review. Etiology of HF was determined from medical record review, and was indicated as ischemic or non-ischemic (includes largely idiopathic and hypertensive HF).

NYHA functional class was determined by careful patient interview. NYHA is a subjective indicator of functional status.⁵⁰ Based on patients’ report of how able they are to perform their usual activities they are assigned a NYHA classification of I (ordinary physical activity causes no symptoms of fatigue, dyspnea, angina or palpitations), II (symptoms with ordinary physical activity that slightly limit physical activity), III (symptoms occur with less than ordinary physical activity and markedly limit activity) or IV (symptoms occur even at rest).

Reproducibility both among different raters (inter-rater reliability) and across the same rater (intra-rater reliability) was insured by training raters and testing them in sample patients until inter-rater agreement is 100%.

Comorbidity burden was assessed using the interview format of the Charlson Comorbidity Index.⁵¹ Higher scores on the Charlson Comorbidity Index indicate greater comorbidity burden. Most conditions are scored with 1 point although some (e.g., hemiplegia, cirrhosis) are assigned >1 point. Scores can range from 0 to 34 but because each subject has HF they will have a score >1. Responses are summed, weighted, and indexed into one of 3 categories (low, moderate, or high) according to the published method. Validity was demonstrated when comorbidity category predicted mortality, complications, health care resource use, length of hospital stay, and discharge disposition.^{52, 53} Comorbidities were documented during the patient and family interview at the time of patient enrollment, and confirmed by reviewing the medical record.

The Duke Activity Status Index (DASI) was used to measure self-reported functional capacity with higher scores indicating higher functional capacity.⁵⁴ The DASI consists of 12 activity items. Each item has four (1-can do without limitation to 4 – unable to do) response options. Each item is scored with a zero given to responses of 2, 3, or 4 all of which indicate there is some degree of limitation performing the activity. For a response of 1 (can do without difficulty), the item is given a weighted score based on the MET level associated with the given activity. The total score is calculated by summing all the item scores, and can range from 0 to 58.2. The reliability⁵⁵ and the validity^{55, 56} of the DASI have been supported in previous studies. The Cronbach's α for this scale in the current study was 0.85.

Psychosocial variables—The severity of depressive symptoms was measured using the Patient Health Questionnaire-9 (PHQ-9).^{57, 58} The PHQ-9 is widely used and has been demonstrated to be valid across diverse groups of individuals including patients with HF.⁵⁹ The items on the PHQ-9 are derived from DSM-IV criteria for diagnosing depression.⁶⁰ This instrument consists of 9 items each rated by the patient using the range of 0 (*not at all*) to 3 (*nearly every day*). The scores range from 0–27 with higher scores indicating more severe depressive symptoms. A score of greater than or equal to 10 is defined as having moderate depressive symptoms,^{57, 58} and this is the cutpoint we used in this study to define the presence of depressive symptoms.

Anxiety: Anxiety was measured using the anxiety scale of the Brief Symptom Inventory (BSI).^{61, 62} The anxiety scale of the BSI consists of six items that are rated by patients on a 5-point Likert type scale that ranges from 0 (no distress related to the item) to 4 (extreme distress related to the item). The score is calculated by averaging the ratings of the items with a possible range from 0 to 4. Acceptable reliability and validity have been demonstrated for this instrument in psychiatric and medical patients, including those with HF.^{61–63}

Perceived Control: Perceived control was measured by the Control Attitudes Scale (CAS).⁶⁴ The CAS consists of a total of 4-items that are ranked by the respondent on a 7-point scale according to the degree to which they agree or disagree with a given statement;

ratings on two of the items are reversed before scoring. The total score can range from 4 to 28 with higher scores indicating greater perceived control. The instrument has well-established reliability and validity in a variety of cardiac populations, including HF.^{38, 65}

Health Perception: Health perception was measured by a single item from the Medical Outcomes Study Short Form –36. Patients rated their health on a Likert type scale 1–5, (1 equals excellent, and 5 equals poor).

Data Analysis

To describe sample characteristics, we compared each demographic, behavioral, clinical and psychosocial characteristic based on gender and presence of depressive symptoms. Comparisons were done using two-way analysis of variance (ANOVA), or chi-square, as appropriate. For ANOVAs, posthoc group comparisons were conducted using the Tukey statistic.

To address Specific Aim 1, chi-square was used to determine the prevalence of depressive symptoms in men and women by comparing the proportion of men and women who met criteria for depressive symptoms. Independent t-tests were used to determine the severity of depressive symptoms in men and women by comparing the means of scores on the PHQ-9 between men and women. Analysis of variance (ANOVA) was used to determine main effects of gender and/or depressive symptoms.

To address Specific Aim 2, regression analyses were used to identify multivariate predictors of depressive symptoms in men and women. Separate multiple regressions were run for men and women. Variables were entered in three blocks: demographic, behavioral and clinical, and psychosocial variables to determine which factors were independently associated with depression scores. At each step, variables were forced into the model in order to provide simultaneous control within the step.

RESULTS

Sample Characteristics

Data from a total of 622 patients, 436 men and 186 women, are included.

Sample characteristics are compared among patients with and without depressive symptoms stratified by gender in Table 1. Interactions were examined first, and if no interaction was present, main effects were examined. There were no main effects for gender or depressive symptoms in the characteristics of ethnicity, marital status, smoking history, or body mass index. The only interaction in gender by depressive symptoms was present for the variable of financial status. Women who reported not having enough to make ends meet were more likely to have higher levels of depressive symptoms than women who reported being financially comfortable or having enough to make ends meet. There was no association between levels of depressive symptoms and financial status in men.

There were differences based on depressive symptoms (i.e., main effect of depressive symptoms) in age, years of education, exercise behavior, NYHA class, comorbidity burden,

health perceptions, anxiety and perceived control (Table 1). With regard to age, both men and women with depressive symptoms were younger compared to those without depressive symptoms (Table 1, $p = 0.02$ main effect for depressive symptoms). Regardless of gender, those with depressive symptoms had attained fewer years of education than those without depressive symptoms ($p = 0.015$, Table 1). In both men ($p = 0.001$) and women ($p = 0.015$), those with depressive symptoms reported performing less exercise than those without depressive symptoms (Table 1). In both men and women ($p = 0.001$ for both), there was a greater proportion of people with depressive symptoms in the worse NYHA functional classes (Table 1). Regardless of gender, those who had depressive symptoms reported a greater comorbidity burden ($p = 0.001$) than those who did not have depressive symptoms (Table 1). In both men and women, health perceptions were worse ($p = 0.001$), anxiety levels were higher ($p = 0.001$), and levels of perceived control were lower ($p = 0.001$) in those with depressive symptoms than in those without depressive symptoms (Table 1).

There were gender differences (i.e., main effect of gender), but no difference based on depressive symptoms in etiology of heart failure (Table 1). More women than men had a non-ischemic etiology (hypertension) for their HF ($p = 0.05$).

There was both a main effect of gender ($p = 0.045$) and a main effect of depressive symptoms ($p = 0.001$) on functional capacity as measured by the DASI (Table 1). Women reported lower functional capacity than men. Individuals of either gender who had depressive symptoms reported lower functional capacity than those without depressive symptoms.

Specific Aim 1

A total of 134 men (28%) met the criterion for depressive symptoms, while 65 women (35%) met the criteria for depressive symptoms ($p = 0.00$). Mean scores on the PHQ-9 were significantly different between men and women (7.02 ± 6.0 versus 8.63 ± 6.3 , respectively, $p = 0.003$), with women having higher mean scores on the PHQ-9, representing greater severity of depressive symptoms than men

Specific Aim 2: Gender-Specific Predictors of Depression

Separate multivariate regression analyses were used to identify predictors of depressive symptoms in men (Table 2) and women (Table 3). In men, demographic variables entered in the first step explained 7% of the variance in depressive symptoms. The R^2 increased to 29% in the second step with the addition of the behavioral and clinical variables. The R^2 increased by an additional 19% in the third step with the addition of the psychosocial variables. The final model explained 49% of the variance in depressive symptoms in men. Six variables independently predicted higher levels of depressive symptoms in men: lower financial status, worse NYHA class, poorer functional status, poor ratings of perceived health, increased anxiety and low perceived control.

In women, demographic variables entered in the first step of the regression model explained 12% variance in depressive symptoms. The R^2 increased by 17% to 27% in the second step with the addition of the behavioral and clinical variables. The R^2 increased by an additional 25% in the third step with the addition of the psychosocial variables. The final model

explained 52% of the variance in depressive symptoms in women. Three variables independently contributed to the prediction of higher levels of depressive symptoms in women: higher BMI, higher anxiety and lower perceived control.

DISCUSSION

The present study is among the first to report differences in behavioral and clinical correlates of depressive symptoms and report commonalities of psychosocial correlates of depressive symptoms in a diverse multicenter outpatient sample of women and men. In the current study, depressive symptoms were predicted by socioeconomic status, NYHA Classification, functional status, current health perception, perceived control and anxiety in men. In women, predictors of depressive symptoms were BMI, perceived control and anxiety. Only perceived control and anxiety were predictors in both men and women. In HF patients their perceptions of their ability to control their symptoms, illness and lives was an important correlate of depressive symptoms. A sense of control is known to be important in reducing anxiety and depression. This relationship has also been found to be true in many cohorts of cardiac patients. In randomized trials, where HF patients have been given control of therapeutic strategies such as home-based exercise programs, daily weights and self-management, investigators have shown favorable effects on QOL. The finding that perceived control is an important correlate of depressive symptoms has implications for the development of more nursing interventions. Reducing morbidity and mortality in HF patients with depressive symptoms should be the goal for optimal outcomes and perhaps men and women need different standards of care. HF patients require a high degree of self-management thus programs focused on increasing autonomy and encouraging patient control in decision making regarding care should be designed. We determined that women with HF reported substantially higher levels of depressive symptoms than men. Few studies of HF patients report significant gender differences in depression. Friedman reports no gender differences in depression among older adults with HF.⁶⁶ When comparing men and women Heo et al found high levels of depression and anxiety, but no significant gender differences.^{67, 68} Vaccarino found a trend toward greater depressive symptoms in women.¹⁹ Sample sizes in the previous studies were relatively small, ranging from 98 to 381,^{19, 66-68} and it may be that the small sample sizes in prior research contribute to the lack of gender differences. This speculation is suggested by the findings from a large study of over 48,000 hospitalized HF patients in which women and whites were more likely to have a history of depression when compared with men and minority groups.¹⁶

In conducting separate multivariable models for each gender, we demonstrated that there are different models of factors associated with the presence of depressive symptoms in women and men with HF. In both men and women with HF, models that included demographic, clinical, behavioral and psychosocial variables explained about half the variance in depressive symptoms. In both men and women, the addition of psychosocial variables explained the largest amount of variance in depressive symptoms. The strong predictive value of psychosocial variables over clinical and demographic variables has been demonstrated previously.^{64, 69} Only perceived control and anxiety were correlates in both men and women. In women, the relationship of anxiety and perceived control to depressive

symptoms was more powerful than similar relationships in men, with depressive symptoms explaining greater than 50% of the model variance in women.

For both men and women with HF, their perceptions of their ability to control their symptoms, illness and lives was an important correlate of depressive symptoms. A sense of control is known to be important in reducing anxiety and depression in patients experiencing cardiac events.^{70, 71} This relationship has also been found to be true in many cohorts of cardiac patients, including those recovering from myocardial infarction and coronary artery bypass graft surgery.^{64, 72} In randomized trials, where HF patients have been given control of therapeutic strategies such as home-based exercise programs, daily weights and self-management of diuretic therapy, investigators have demonstrated improved outcomes.^{38, 73} The finding that perceived control is an important correlate of depressive symptoms has implications for the development of more nursing interventions.

For both men and women in the current study, anxiety was a correlate of depressive symptoms. Anxiety, along with other negative emotions, is known to be predictive of CHD⁷⁴ and stroke.⁷⁵ Researchers have reported that both men and women who struggle with the emotional and life-altering effects of HF have heightened levels of anxiety.⁴² Further, anxiety is highly prevalent in HF patients, with reports as high as 63%, similar to rates in lung disease or cancer.⁷⁶ Of note, anxiety at any level in the HF population has negative consequences.^{33, 34} The current study confirms the strong association of anxiety and depressive symptoms in both men and women. While an initial study indicated that a stress-management intervention may reduce depressive symptoms, along with other negative physical and emotional symptoms, in HF patients,⁷⁷ future studies may need to identify gender-specific interventions to reduce anxiety in men and women with HF in order to positively affect comorbid depression.

In men, depressive symptoms were associated with financial status, functional status (NYHA classification), functional capacity (DASI scores), current health perception, perceived control and anxiety. The NYHA classification provides a measurement of perceptions of disease - related restrictions in physical activity.⁷⁸ In this study, NYHA classification contributed independently to the presence of depressive symptoms for men, but not for women with HF. Physical restrictions that affect daily activities have been reported as more bothersome for males which may affect their perception of self-worth.⁷⁹ Additionally, it is known that MI patients who have depressive symptoms are less likely to exercise.⁸⁰ Further, the relationship between depressive symptoms is believed to be bidirectional, so that both contribute synergistically to a downward spiral characterized by Whooley et al as a “mutually reinforcing” phenomena.⁸⁰ It is possible that this “mutually reinforcing” relationship of depressive symptoms and lack of exercise is mediated by other factors, such as increased negative health perceptions and lower levels of perceived control over health, among non-exercisers. Further study is needed to elucidate these relationships in the context of HF and inform targeted interventions aimed at reducing depressive symptoms and increasing exercise in HF patients.

Health perception was a significant predictor of depressive symptoms for men only. A possible explanation for this male-only finding may be that men may place more value than

women on role-related self-worth. They describe the physical limitations, social restrictions in their work and leisure activities caused by HF as hindering them from taking part in activities that had been a natural part of their lives.⁸¹ For men recovering from MI, in particular, functional status appears to have great importance and has been shown to have significant impact on self rating of health.⁸² Faced with HF symptoms, men with HF may perceive their health as unrealistically low, underestimate their actual functional potential, and believe that they have lost core social roles as providers. Such negative thinking would be consistent with depressive symptoms.⁷ While further study to support our findings is needed, interventions for men with HF that are designed to focus on attaining realistic perceptions of health warrant exploration.

Body mass index was a correlate of depressive symptoms in women with HF only. Weight is an important issue in women that can transcend their feelings about body image and societal expectations.⁸³ Being overweight or obese may potentially increase feelings of anxiety and low perceived control in women in this study. Primary care providers need to address BMI and develop individualized strategies for weight reduction and management to deter the escalating and detrimental effects of high BMI on depressive symptoms in women. Future studies should include exercise, nutrition and psychological interventions to reduce the incidence and progression of depressive symptoms in women and be longitudinal in design.

Implications for Nursing

Reducing hospital readmissions, morbidity and mortality in HF patients with depressive symptoms should be the goal for optimal outcomes and perhaps men and women need different standards of care.⁷⁹ Advanced Practice Nurses (APNs), who possess advanced assessment and disease management skills, are in a unique position to assess and evaluate depressive symptoms in HF patients and to employ educational and counseling programs to reduce depressive symptoms in HF patients. Documentation of mental health screening is essential in HF patients in order for eligible patients to receive necessary mental health treatment.⁸⁴ APNs are adept at pulling care teams together to institute patient –centered care, promoting care integration and coordination, and improving access to care. Such activities hold great potential to identify and treat depressive symptoms in men and women with HF. HF patients require a high degree of self-management thus programs focused on increasing autonomy and encouraging patient control in decision making regarding care should be designed. Educating patients and families to identify worsening symptoms and to adjust and revise treatments is an example of increasing control and potentially decreasing hospitalizations and economic burdens on the family and on society.

Previous studies have found little gender difference in predictors of symptoms of depression.^{66, 85} Important findings from the present study should be used to develop different management models for men and women. Nursing will play a critical role in adequately identifying and managing HF patients at risk for depression, and decreasing the personal and societal burden of depression in this growing population.

Strengths and Limitations—The strengths of our investigation included a relatively large sample size, multicenter enrollment of subjects, and the availability of a broad spectrum of covariates of interest for multivariable analyses by gender. In most previous studies interpretation of the findings on gender differences has been seriously hampered due to power problems. In addition the number of variables was restricted and inconsistent across designs. Limitations of the current study include a use of a convenience sample, which limits the generalizability. Patients who have depressive symptoms may be less likely to participate in research. Data for the study were collected by self-report questionnaires that may not reflect patients' true feelings. The sample was predominately Caucasian and English-speaking, so caution should be used in generalizing results to those patients who do not speak English and are not Caucasian.

Conclusions

Heart failure is a debilitating and chronic clinical syndrome which is compromised by the presence of depressive symptoms. Studies of correlates of depressive symptoms among patients with HF are important because symptomatic depression is a widespread and debilitating illness. The present study demonstrates that there are differences and commonalities in correlates of depressive symptoms in men and women from a relatively large, heterogeneous sample of patients with HF from diverse parts of the country.

We found distinct differences in demographic, clinical, behavioral and psychosocial predictors of symptoms of depression for men and women. Identification of correlates of depressive symptoms and of gender differences is helpful for the development of gender-based risk profiles, and determination of targets of gender-based interventions to reduce depressive symptoms. In the present study the psychological factors of perceived control and anxiety were independent correlates of depressive symptoms in both men and women. Although anxiety and perceived control were also strong correlates of depressive symptoms for women with HF, they seemed to stem from important relationships rather than physical achievement. Evidence from this study provides justification to support gender specific standards of care⁷⁹ and gender specific treatments and management to address the problem of depressive symptoms, which have been associated with increased use of health care resources and increased costs.⁷

Acknowledgments

Funding: The project described was supported by grant numbers R01 NR008567 and R01 NR009280 from NIH, the National Institute of Nursing Research, and by a Center grant, 1P20NR010679 from NIH, National Institute of Nursing Research. It was also supported by NRSA Clinical Scholars Training in Cardiovascular Science (grant number: T32 HL091812).

References

1. Lloyd-Jones D, Adams R, Carnethon M, et al. Heart disease and stroke statistics--2009 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation*. Jan 27; 2009 119(3):e21–181. [PubMed: 19075105]
2. Bundkirchen A, Schwinger R. Epidemiology and economic burden of chronic heart failure. *European Heart Journal Supplements*. 2004; 6(Supplement D):D57–D60.

3. Rutledge T, Reis VA, Linke SE, Greenberg BH, Mills PJ. Depression in heart failure a meta-analytic review of prevalence, intervention effects, and associations with clinical outcomes. *J Am Coll Cardiol.* Oct 17; 2006 48(8):1527–1537. [PubMed: 17045884]
4. Chriss PM, Shepesh J, Carlson B, Riegel B. Predictors of successful heart failure self-care maintenance in the first three months after hospitalization. *Heart Lung.* Nov-Dec;2004 33(6):345–353. [PubMed: 15597287]
5. Sayers SL, Riegel B, Pawlowski S, Coyne JC, Samaha FF. Social support and self-care of patients with heart failure. *Ann Behav Med.* Feb; 2008 35(1):70–79. [PubMed: 18347906]
6. McMurray JJ, Stewart S. Epidemiology, aetiology, and prognosis of heart failure. *Heart.* May; 2000 83(5):596–602. [PubMed: 10768918]
7. Gottlieb SS, Khatta M, Friedmann E, et al. The influence of age, gender, and race on the prevalence of depression in heart failure patients. *J Am Coll Cardiol.* May 5; 2004 43(9):1542–1549. [PubMed: 15120809]
8. Jiang W, Alexander J, Christopher E, et al. Relationship of depression to increased risk of mortality and rehospitalization in patients with congestive heart failure. *Arch Intern Med.* Aug 13–27; 2001 161(15):1849–1856. [PubMed: 11493126]
9. Macchia A, Monte S, Pellegrini F, et al. Depression worsens outcomes in elderly patients with heart failure: an analysis of 48,117 patients in a community setting. *Eur J Heart Fail.* Jul; 2008 10(7):714–721. [PubMed: 18565789]
10. Murberg TA, Bru E, Aarsland T, Svebak S. Functional status and depression among men and women with congestive heart failure. *Int J Psychiatry Med.* 1998; 28(3):273–291. [PubMed: 9844832]
11. Sullivan M, Levy WC, Russo JE, Spertus JA. Depression and health status in patients with advanced heart failure: a prospective study in tertiary care. *J Card Fail.* Oct; 2004 10(5):390–396. [PubMed: 15470649]
12. Sullivan M, Simon G, Spertus J, Russo J. Depression-related costs in heart failure care. *Arch Intern Med.* Sep 9; 2002 162(16):1860–1866. [PubMed: 12196084]
13. Thomas SA, Chapa DW, Friedmann E, et al. Depression in patients with heart failure: prevalence, pathophysiological mechanisms, and treatment. *Crit Care Nurse.* Apr; 2008 28(2):40–55. [PubMed: 18378727]
14. Roger VL, Weston SA, Redfield MM, et al. Trends in heart failure incidence and survival in a community-based population. *Jama.* Jul 21; 2004 292(3):344–350. [PubMed: 15265849]
15. Fallor H, Stork S, Schowalter M, et al. Depression and survival in chronic heart failure: does gender play a role? *Eur J Heart Fail.* Oct; 2007 9(10):1018–1023. [PubMed: 17660002]
16. Albert NM, Fonarow GC, Abraham WT, et al. Depression and clinical outcomes in heart failure: an OPTIMIZE-HF analysis. *Am J Med.* Apr; 2009 122(4):366–373. [PubMed: 19332232]
17. Havranek EP, Spertus JA, Masoudi FA, Jones PG, Rumsfeld JS. Predictors of the onset of depressive symptoms in patients with heart failure. *J Am Coll Cardiol.* Dec 21; 2004 44(12):2333–2338. [PubMed: 15607395]
18. Junger J, Schellberg D, Muller-Tasch T, et al. Depression increasingly predicts mortality in the course of congestive heart failure. *Eur J Heart Fail.* Mar 2; 2005 7(2):261–267. [PubMed: 15701476]
19. Vaccarino V, Kasl SV, Abramson J, Krumholz HM. Depressive symptoms and risk of functional decline and death in patients with heart failure. *J Am Coll Cardiol.* Jul; 2001 38(1):199–205. [PubMed: 11451275]
20. Kessler RC. Epidemiology of women and depression. *J Affect Disord.* Mar; 2003 74(1):5–13. [PubMed: 12646294]
21. Gottlieb SS, Kop WJ, Ellis SJ, et al. Relation of depression to severity of illness in heart failure (from Heart Failure And a Controlled Trial Investigating Outcomes of Exercise Training [HF-ACTION]). *Am J Cardiol.* May 1; 2009 103(9):1285–1289. [PubMed: 19406273]
22. Kessler D, Sharp D, Lewis G. Screening for depression in primary care. *Br J Gen Pract.* Sep; 2005 55(518):659–660. [PubMed: 16176729]

23. Rohyans LM, Pressler SJ. Depressive symptoms and heart failure: examining the sociodemographic variables. *Clin Nurse Spec.* May-Jun;2009 23(3):138–144. [PubMed: 19395890]
24. Freedland KE, Rich MW, Skala JA, Carney RM, Davila-Roman VG, Jaffe AS. Prevalence of depression in hospitalized patients with congestive heart failure. *Psychosom Med.* Jan-Feb;2003 65(1):119–128. [PubMed: 12554823]
25. Ladwig KH, Lehmacher W, Roth R, Breithardt G, Budde T, Borggrefe M. Factors which provoke post-infarction depression: results from the post-infarction late potential study (PILP). *J Psychosom Res.* Dec; 1992 36(8):723–729. [PubMed: 1432862]
26. Brummett BH, Babyak MA, Barefoot JC, et al. Social support and hostility as predictors of depressive symptoms in cardiac patients one month after hospitalization: a prospective study. *Psychosom Med.* Nov-Dec;1998 60(6):707–713. [PubMed: 9847029]
27. Anthony JC, Petronis KR. Suspected risk factors for depression among adults 18–44 years old. *Epidemiology.* Mar; 1991 2(2):123–132. [PubMed: 1932309]
28. Evangelista LS, Ter-Galstanyan A, Moughrabi S, Moser DK. Anxiety and depression in ethnic minorities with chronic heart failure. *J Card Fail.* Sep; 2009 15(7):572–579. [PubMed: 19700133]
29. Johansson P, Dahlstrom U, Brostrom A. Consequences and predictors of depression in patients with chronic heart failure: implications for nursing care and future research. *Prog Cardiovasc Nurs.* Fall;2006 21(4):202–211. [PubMed: 17170596]
30. Horwich TB, Fonarow GC, Hamilton MA, MacLellan WR, Woo MA, Tillisch JH. The relationship between obesity and mortality in patients with heart failure. *J Am Coll Cardiol.* Sep; 2001 38(3): 789–795. [PubMed: 11527635]
31. Katon W, Lin EH, Kroenke K. The association of depression and anxiety with medical symptom burden in patients with chronic medical illness. *Gen Hosp Psychiatry.* Mar-Apr;2007 29(2):147–155. [PubMed: 17336664]
32. Koike AK, Unutzer J, Wells KB. Improving the care for depression in patients with comorbid medical illness. *Am J Psychiatry.* Oct; 2002 159(10):1738–1745. [PubMed: 12359681]
33. De Jong MJ, Moser DK, An K, Chung ML. Anxiety is not manifested by elevated heart rate and blood pressure in acutely ill cardiac patients. *Eur J Cardiovasc Nurs.* Sep; 2004 3(3):247–253. [PubMed: 15350235]
34. Friedmann E, Thomas SA, Liu F, Morton PG, Chapa D, Gottlieb SS. Relationship of depression, anxiety, and social isolation to chronic heart failure outpatient mortality. *Am Heart J.* Nov; 2006 152(5):940, e941–948. [PubMed: 17070164]
35. Katon WJ. Clinical and health services relationships between major depression, depressive symptoms, and general medical illness. *Biol Psychiatry.* Aug 1; 2003 54(3):216–226. [PubMed: 12893098]
36. Rumsfeld JS, Havranek E, Masoudi FA, et al. Depressive symptoms are the strongest predictors of short-term declines in health status in patients with heart failure. *J Am Coll Cardiol.* Nov 19; 2003 42(10):1811–1817. [PubMed: 14642693]
37. Carels RA. The association between disease severity, functional status, depression and daily quality of life in congestive heart failure patients. *Qual Life Res.* Feb; 2004 13(1):63–72. [PubMed: 15058788]
38. Dracup K, Westlake C, Erickson VS, Moser DK, Caldwell ML, Hamilton MA. Perceived control reduces emotional stress in patients with heart failure. *J Heart Lung Transplant.* Jan; 2003 22(1): 90–93. [PubMed: 12531418]
39. Vollman MW, Lamontagne LL, Hepworth JT. Coping and depressive symptoms in adults living with heart failure. *J Cardiovasc Nurs.* Mar-Apr;2007 22(2):125–130. [PubMed: 17318038]
40. Scherer M, Himmel W, Stanske B, et al. Psychological distress in primary care patients with heart failure: a longitudinal study. *Br J Gen Pract.* Oct; 2007 57(543):801–807. [PubMed: 17925137]
41. Jiang W, Kuchibhatla M, Cuffe MS, et al. Prognostic value of anxiety and depression in patients with chronic heart failure. *Circulation.* Nov 30; 2004 110(22):3452–3456. [PubMed: 15557372]
42. Konstam V, Moser DK, De Jong MJ. Depression and anxiety in heart failure. *J Card Fail.* Aug; 2005 11(6):455–463. [PubMed: 16105637]

43. Haworth JE, Moniz-Cook E, Clark AL, Wang M, Waddington R, Cleland JG. Prevalence and predictors of anxiety and depression in a sample of chronic heart failure patients with left ventricular systolic dysfunction. *Eur J Heart Fail.* Aug; 2005 7(5):803–808. [PubMed: 16054436]
44. Friedman MM, Griffin JA. Relationship of physical symptoms and physical functioning to depression in patients with heart failure. *Heart Lung.* Mar-Apr;2001 30(2):98–104. [PubMed: 11248712]
45. Turvey CL, Schultz K, Arndt S, Wallace RB, Herzog R. Prevalence and correlates of depressive symptoms in a community sample of people suffering from heart failure. *J Am Geriatr Soc.* Dec; 2002 50(12):2003–2008. [PubMed: 12473012]
46. Riegel B, Moser DK, Carlson B, et al. Gender differences in quality of life are minimal in patients with heart failure. *Journal of Cardiac Failure.* Feb; 2003 9(1):42–48. [PubMed: 12612872]
47. Riegel B, Moser DK, Glaser D, et al. The Minnesota Living With Heart Failure Questionnaire: Sensitivity to differences and responsiveness to intervention intensity in a clinical population. *Nursing Research.* Jul-Aug;2002 51(4):209–218. [PubMed: 12131233]
48. Riegel B, Moser DK, Rayens MK, et al. Ethnic differences in quality of life in persons with heart failure. *Journal of Cardiac Failure.* Feb; 2008 14(1):41–47. [PubMed: 18226772]
49. Wu JR, Moser DK, Chung ML, Lennie TA. Predictors of medication adherence using a multidimensional adherence model in patients with heart failure. *J Card Fail.* Sep; 2008 14(7): 603–614. [PubMed: 18722327]
50. Mills RM Jr, Haught WH. Evaluation of heart failure patients: objective parameters to assess functional capacity. *Clin Cardiol.* Jun; 1996 19(6):455–460. [PubMed: 8790948]
51. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *Journal of Chronic Diseases.* 1987; 40(5):373–383. [PubMed: 3558716]
52. Peterson JC, Charlson ME, Williams-Russo P, et al. New postoperative depressive symptoms and long-term cardiac outcomes after coronary artery bypass surgery. *Am J Geriatr Psychiatry.* Mar-Apr;2002 10(2):192–198. [PubMed: 11925280]
53. Katz JN, Chang LC, Sangha O, Fassel AH, Bates DW. Can comorbidity be measured by questionnaire rather than medical record review? *Med Care.* Jan; 1996 34(1):73–84. [PubMed: 8551813]
54. Hlatky MA, Boineau RE, Higginbotham MB, et al. A brief self-administered questionnaire to determine functional capacity (the Duke Activity Status Index). *American Journal of Cardiology.* Sep 15; 1989 64(10):651–654. [PubMed: 2782256]
55. Alonso J, Permanyer-Miralda G, Cascant P, Brotons C, Prieto L, Soler-Soler J. Measuring functional status of chronic coronary patients. Reliability, validity and responsiveness to clinical change of the reduced version of the Duke Activity Status Index (DASI). *Eur Heart J.* Mar; 1997 18(3):414–419. [PubMed: 9076377]
56. Parissis JT, Nikolaou M, Birmpa D, et al. Clinical and prognostic value of Duke's Activity Status Index along with plasma B-type natriuretic peptide levels in chronic heart failure secondary to ischemic or idiopathic dilated cardiomyopathy. *Am J Cardiol.* Jan 1; 2009 103(1):73–75. [PubMed: 19101233]
57. Spitzer RL, Kroenke K, Williams JB. Validation and utility of a self-report version of PRIME-MD: the PHQ primary care study. *Primary Care Evaluation of Mental Disorders. Patient Health Questionnaire.* *JAMA.* Nov 10; 1999 282(18):1737–1744. [PubMed: 10568646]
58. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *Journal of General Internal Medicine.* Sep; 2001 16(9):606–613. [PubMed: 11556941]
59. Huang FY, Chung H, Kroenke K, Delucchi KL, Spitzer RL. Using the Patient Health Questionnaire-9 to measure depression among racially and ethnically diverse primary care patients. *Journal of General Internal Medicine.* Jun; 2006 21(6):547–552. [PubMed: 16808734]
60. Association AP. *Diagnostic and Statistical Manual of Mental Disorders.* 4. Washington, DC: American Psychiatric Association; 2000. Text Revision
61. Derogatis LR, Melisaratos N. The Brief Symptom Inventory: an introductory report. *Psychol Med.* Aug; 1983 13(3):595–605. [PubMed: 6622612]

62. Derogatis, LP. BSI. Brief Symptom Inventory. Administration, scoring, and procedure manual. Minneapolis: National Computer Systems, Inc; 1993.
63. Abu Ruz MA, Lennie TA, Riegel B, et al. Anxiety can be measured quickly and reliably in patients hospitalized for acute myocardial infarction (abstract). *Circulation*. 2005; 112:II-392.
64. Moser DK, Dracup K. Psychosocial recovery from a cardiac event: the influence of perceived control. *Heart Lung*. Jul-Aug;1995 24(4):273–280. [PubMed: 7591794]
65. Moser DK, Dracup K. Impact of cardiopulmonary resuscitation training on perceived control in spouses of recovering cardiac patients. *Res Nurs Health*. Aug; 2000 23(4):270–278. [PubMed: 10940952]
66. Friedman MM. Gender differences in the health related quality of life of older adults with heart failure. *Heart Lung*. Sep-Oct;2003 32(5):320–327. [PubMed: 14528189]
67. Heo S, Moser DK, Widener J. Gender differences in the effects of physical and emotional symptoms on health-related quality of life in patients with heart failure. *Eur J Cardiovasc Nurs*. Jun; 2007 6(2):146–152. [PubMed: 16919502]
68. Heo S, Moser DK, Lennie TA, Riegel B, Chung ML. Gender differences in and factors related to self-care behaviors: a cross-sectional, correlational study of patients with heart failure. *Int J Nurs Stud*. Dec; 2008 45(12):1807–1815. [PubMed: 18674762]
69. Dracup K, Walden JA, Stevenson LW, Brecht ML. Quality of life in patients with advanced heart failure. *J Heart Lung Transplant*. Mar-Apr;1992 11(2 Pt 1):273–279. [PubMed: 1576133]
70. Ziemann KM, Dracup K. Patient-nurse contracts in critical care: a controlled trial. *Prog Cardiovasc Nurs*. Jul-Sep;1990 5(3):98–103. [PubMed: 2267246]
71. Miller SM. Controllability and human stress: method, evidence and theory. *Behav Res Ther*. 1979; 17(4):287–304. [PubMed: 486033]
72. Moser DK, Riegel B, McKinley S, et al. The control attitudes scale-revised: psychometric evaluation in three groups of patients with cardiac illness. *Nurs Res*. Jan-Feb;2009 58(1):42–51. [PubMed: 19092554]
73. Rich MW, Beckham V, Wittenberg C, Leven CL, Freedland KE, Carney RM. A multidisciplinary intervention to prevent the readmission of elderly patients with congestive heart failure. *N Engl J Med*. Nov 2; 1995 333(18):1190–1195. [PubMed: 7565975]
74. Kubzansky LD, Cole SR, Kawachi I, Vokonas P, Sparrow D. Shared and unique contributions of anger, anxiety, and depression to coronary heart disease: a prospective study in the normative aging study. *Ann Behav Med*. Feb; 2006 31(1):21–29. [PubMed: 16472035]
75. Morrison V, Pollard B, Johnston M, MacWalter R. Anxiety and depression 3 years following stroke: demographic, clinical, and psychological predictors. *J Psychosom Res*. Oct; 2005 59(4): 209–213. [PubMed: 16223623]
76. Riedinger MS, Dracup KA, Brecht ML. Quality of life in women with heart failure, normative groups, and patients with other chronic conditions. *Am J Crit Care*. May; 2002 11(3):211–219. [PubMed: 12022484]
77. Luskin F, Reitz M, Newell K, Quinn TG, Haskell W. A controlled pilot study of stress management training of elderly patients with congestive heart failure. *Prev Cardiol*. Fall;2002 5(4):168–172. [PubMed: 12417824]
78. Norozi K, Wessel A, Buchhorn R, et al. Is the Ability index superior to the NYHA classification for assessing heart failure?: comparison of two classification scales in adolescents and adults with operated congenital heart defects. *Clin Res Cardiol*. Aug; 2007 96(8):542–547. [PubMed: 17593319]
79. Stromberg A, Martensson J. Gender differences in patients with heart failure. *Eur J Cardiovasc Nurs*. Apr; 2003 2(1):7–18. [PubMed: 14622644]
80. Whooley MA, de Jonge P, Vittinghoff E, et al. Depressive symptoms, health behaviors, and risk of cardiovascular events in patients with coronary heart disease. *Jama*. Nov 26; 2008 300(20):2379–2388. [PubMed: 19033588]
81. Martensson J, Karlsson JE, Fridlund B. Male patients with congestive heart failure and their conception of the life situation. *J Adv Nurs*. Mar; 1997 25(3):579–586. [PubMed: 9080286]

82. Ziegelstein RC, Fauerbach JA, Stevens SS, Romanelli J, Richter DP, Bush DE. Patients with depression are less likely to follow recommendations to reduce cardiac risk during recovery from a myocardial infarction. *Arch Intern Med.* Jun 26; 2000 160(12):1818–1823. [PubMed: 10871976]
83. Halm M, Penque S, Doll N, Behrs M. Women and cardiac rehabilitation: referral and compliance patterns. *J Cardiovasc Nurs.* Apr; 1999 13(3):83–92. [PubMed: 10098008]
84. Cully JA, Jimenez DE, Ledoux TA, Deswal A. Recognition and Treatment of Depression and Anxiety Symptoms in Heart Failure. *Prim Care Companion J Clin Psychiatry.* 2009; 11(3):103–109. [PubMed: 19617942]
85. Zambroski CH, Moser DK, Bhat G, Ziegler C. Impact of symptom prevalence and symptom burden on quality of life in patients with heart failure. *Eur J Cardiovasc Nurs.* Sep; 2005 4(3):198–206. [PubMed: 15916924]

Table 1
Sample Characteristics Compared by Gender and by Depressive Symptom Status

| | Male (N=436) | | Female (N=186) | | p value |
|--|------------------------|---------------------|------------------------|---------------------|------------------------------------|
| | No Depressive Symptoms | Depressive Symptoms | No Depressive Symptoms | Depressive Symptoms | |
| Demographic Characteristics | | | | | |
| Age (Mean ± SD) | 62.49 ± 12.79 | 57.77 ± 11.53 | 61.33 ± 13.41 | 60.60 ± 12.38 | .021 [‡] |
| Ethnicity, N (%) | | | | | |
| African American | 43 (14) | 22 (18) | 35 (29) | 14 (22) | NS |
| Caucasian | 265 (85) | 102 (82) | 82 (68) | 48 (74) | |
| All others | 4 (1) | 0 (0) | 4 (3) | 3 (5) | |
| Marital Status, N (%) | | | | | |
| Single/Divorced/Widowed (compared to married) | 101 (32) | 51 (41) | 81 (67) | 37(57) | NS |
| Years Education, (Mean ± SD) | 13.7 ± 3.6 | 12.0 ± 3.0 | 13.3 ± 2.9 | 12.6 ± 2.0 | .015 [‡] |
| Financial Status, N (%) | | | | | |
| Comfortable | 117 (38) | 34 (28) | 37 (37) | 12 (19) | .001 [*] |
| Enough to make ends meet | 151 (49) | 49 (40) | 58 (48) | 53 (51) | |
| Not enough to make ends meet | 42 (14) | 40 (33) | 26 (22) | 20 (31) | |
| Behavioral and Clinical Characteristics | | | | | |
| Exercise (during past week), N (%) | | | | | |
| None | 82 (26) | 58 (47) | 40 (33) | 34 (52) | .001 for men; .015 for women ** |
| <30 min per week | 42 (14) | 22 (18) | 21 (17) | 12 (19) | |
| 30–60 min per week | 49 (16) | 13 (11) | 26 (22) | 6 (9) | |
| 1–3 hours per week | 77 (25) | 15 (12) | 18 (15) | 11 (17) | |
| >3 hours per week | 62 (20) | 15 (12) | 16 (13) | 2 (3) | |
| Smoking, N (%) | | | | | |
| Current or quit < 1 year | 73 (23) | 42 (33) | 28 (23) | 19 (29) | NS |

| | Male (N=436) | | Female (N=186) | | p value |
|--|------------------------|---------------------|------------------------|---------------------|---|
| | No Depressive Symptoms | Depressive Symptoms | No Depressive Symptoms | Depressive Symptoms | |
| Former quit > 1 year | 152 (49) | 49 (40) | 43 (36) | 27 (42) | |
| Never | 87 (28) | 33 (27) | 50 (41) | 19 (29) | |
| Body Mass Index, (Mean ± SD) | 30.27 ± 7.22 | 29.95 ± 7.40 | 30.38 ± 8.18 | 32.34 ± 8.92 | NS |
| Charlson Comorbidity Index Scores (Mean ± SD) | 3.27 ± 2.03 | 3.75 ± 2.33 | 3.09 ± 1.98 | 4.34 ± 2.72 | .001 [†] |
| Etiology, N (%) | | | | | .05 [§] |
| Ischemic | 188 (59) | 66 (59) | 47 (33) | 33 (43) | |
| Non-ischemic (idiopathic, hypertensive, alcoholic) | 132 (41) | 53 (47) | 95 (67) | 43 (57) | |
| NYHA Functional Class, N (%) | | | | | .001 for men and women** |
| I & II | 164 (53) | 28 (23) | 49 (41) | 10 (15) | |
| III & IV | 148 (47) | 96 (77) | 72 (60) | 55 (85) | |
| Functional Capacity (DASI scores) (Mean ± SD) | 16.97 ± 14.00 | 7.00 ± 8.01 | 12.86 ± 11.56 | 6.62 ± 7.46 | .001 for depressive symptoms; .045 for gender ^{††} |
| Psychosocial Characteristics | | | | | |
| Health Perception (now), N (%): | | | | | |
| Excellent | 7 (2) | 2 (2) | 2 (2) | 1 (2) | .001*** |
| Very good | 40 (13) | 4 (3) | 16 (13) | 2 (3) | |
| Good | 113 (36) | 14 (11) | 43 (36) | 16 (25) | |
| Fair | 105 (34) | 58 (47) | 43 (36) | 16 (25) | |
| Poor | 47 (15) | 46 (37) | 17 (14) | 30 (46) | |
| Perceived control (CAS total scores), (Mean ± SD) | 17.69 ± 5.13 | 14.09 ± 5.54 | 16.90 ± 5.82 | 14.31 ± 6.14 | .001 [†] |
| Anxiety (BSI scores), (Mean ± SD) | .48 ± .59 | 1.53 ± 1.09 | .61 ± .62 | 1.58 ± 1.11 | .001 [†] |

Notes: CAS=Control Attitudes Scale; BSI= Brief Symptom Inventory; DASI= Duke Activity Status Index; PHQ-9=Patient Health Questionnaire-9;

* Differences by gender and depressive symptoms by chi-square

[†] Main effect of gender by ANOVA;

‡ Main effect of depressive symptoms by ANOVA

§ Differences by gender only using chi-square

** Differences by depressive symptoms only using chi-square

NS= No significant differences

Table 2

Predictors of depressive symptoms in men (N= 436)

| Predictor variables | Standardized β^* | p-value* |
|-----------------------------------|------------------------|----------|
| Age | -0.003 | 0.937 |
| Ethnicity | 0.026 | 0.485 |
| Marital status | -0.013 | 0.731 |
| Education | -0.022 | 0.546 |
| Financial status | 0.087 | 0.027 |
| Exercise | -0.065 | 0.074 |
| Smoking | -0.014 | 0.709 |
| Body mass index | -0.019 | 0.603 |
| Charlson Comorbidity Index scores | 0.017 | 0.665 |
| Etiology | 0.075 | 0.061 |
| NYHA class | 0.138 | 0.001 |
| Functional capacity (DASI) | -0.159 | <0.001 |
| Health perception | 0.086 | 0.043 |
| Perceived control | -0.120 | 0.002 |
| Anxiety | 0.423 | <0.001 |

Notes: NYHA= New York Heart Association; DASI Duke Activity Status Index;

* Standardized β s and p-values shown are for the final model

Table 3

Predictors of depressive symptoms in women (N= 186)

| Predictor variables | Standardized β * | p-value* |
|-----------------------------------|------------------------|----------|
| Age | -0.003 | 0.954 |
| Ethnicity | 0.084 | 0.142 |
| Marital status | 0.066 | 0.234 |
| Education | -0.098 | 0.069 |
| Financial status | 0.085 | 0.119 |
| Exercise | -0.097 | 0.086 |
| Smoking | -0.022 | 0.693 |
| Body mass index | 0.118 | 0.037 |
| Charlson Comorbidity Index scores | 0.114 | 0.059 |
| Etiology | 0.013 | 0.830 |
| NYHA class | 0.108 | 0.073 |
| Functional capacity (DASI) | -0.030 | 0.636 |
| Health perception | -0.001 | 0.983 |
| Perceived control | -0.147 | 0.013 |
| Anxiety | 0.539 | <0.001 |

Notes: NYHA= New York Heart Association; DASI Duke Activity Status Index;

* Standardized β s and p-values shown are for the final model