



**EMORY**  
LIBRARIES &  
INFORMATION  
TECHNOLOGY

**OpenEmory**

# **Statins Reduce Epicardial Adipose Tissue Attenuation Independent of Lipid Lowering: A Potential Pleiotropic Effect**

[Paolo Raggi](#), *Emory University*

[Varuna Gadiyaram](#), *Emory University*

[Chao Zhang](#), *Emory University*

[Zhengjia Chen](#), *Emory University*

[Gary Lopaschuk](#), *University of Alberta*

[Arthur Stillman](#), *Emory University*

---

**Journal Title:** Journal of the American Heart Association

**Volume:** Volume 8, Number 12

**Publisher:** Wiley Open Access: Creative Commons Attribution  
Non-Commercial | 2019-06-18, Pages e013104-e013104

**Type of Work:** Article | Final Publisher PDF

**Publisher DOI:** 10.1161/JAHA.119.013104

**Permanent URL:** <https://pid.emory.edu/ark:/25593/txczp>

---

Final published version: <http://dx.doi.org/10.1161/JAHA.119.013104>

## **Copyright information:**

© 2019 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley.

This is an Open Access work distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



Accessed December 15, 2019 2:52 PM EST

# Statins Reduce Epicardial Adipose Tissue Attenuation Independent of Lipid Lowering: A Potential Pleiotropic Effect

Paolo Raggi, MD, PhD; Varuna Gadiyaram, MD; Chao Zhang, PhD; Zhengjia Chen, PhD; Gary Lopaschuk, PhD; Arthur E. Stillman, MD, PhD

**Background**—High epicardial adipose tissue (EAT) attenuation (Hounsfield units [HUs]) on computed tomography is considered a marker of inflammation and is associated with an increased risk of cardiovascular events. Statins reduce the volume of EAT, but it is unknown whether they affect EAT HUs.

**Methods and Results**—We reviewed the chest computed tomographic scans of 420 postmenopausal women randomized to either 80 mg of atorvastatin or 40 mg of pravastatin daily and rescanned after 1 year to measure change in coronary artery calcium score. EAT HUs were measured near the proximal right coronary artery and remote from any area of coronary artery calcium. Computed tomographic images were also queried for subcutaneous adipose tissue (SubQ) attenuation (HUs) change over time. The mean patients' age was  $65 \pm 6$  years. The baseline EAT HU value was higher than the SubQ HU value ( $-89.4 \pm 24.0$  HU versus  $-123.3 \pm 30.4$  HU;  $P < 0.001$ ). The EAT HU value decreased significantly in the entire cohort ( $-5.4 \pm 29.7$  HU [ $-6\%$  change];  $P < 0.001$ ), but equally in the patients given atorvastatin and pravastatin ( $-6.35 \pm 31$  HU and  $-4.55 \pm 28$  HU;  $P = 0.55$ ). EAT HU change was not associated with change in total cholesterol, low-density lipoprotein cholesterol, coronary artery calcium, and EAT volume (all  $P =$  not significant). Change in high-density lipoprotein cholesterol was marginally associated with EAT HU change ( $P = 0.07$ ). Statin treatment did not induce a change in SubQ HUs.

**Conclusions**—Statins induced a decrease in EAT HUs over time, independent of intensity of low-density lipoprotein cholesterol lowering. The positive effect on EAT and the neutral effect on SubQ suggest that statins induced a decrease in metabolic activity in EAT by reduction in cellularity, vascularity, or inflammation. The clinical significance of the observed change in EAT HUs remains to be demonstrated. (*J Am Heart Assoc.* 2019;8:e013104. DOI: 10.1161/JAHA.119.013104.)

**Key Words:** computed tomography • epicardial fat • low-density lipoprotein • statin therapy

Epicardial adipose tissue (EAT) is visceral fat surrounding and in direct contact with the coronary artery adventitia, and it is believed to be implicated in the development of atherosclerosis. In their initial publication, Mazurek et al<sup>1</sup> demonstrated that in patients scheduled to undergo elective coronary artery bypass surgery, EAT is highly inflamed. However, the subcutaneous adipose tissue (SubQ) of the same patients did not show any or only minimal signs of

inflammation. Whether EAT contributes directly to the development of atherosclerosis via paracrine effects or via systemic mechanisms remains to be clarified. Nonetheless, there is accumulating evidence that EAT is more plentiful in patients at risk of atherosclerosis or in those with preclinical atherosclerosis<sup>2-4</sup> and adverse outcomes.<sup>5</sup> Recent investigations correlated a higher computed tomography (CT) attenuation of EAT with coronary artery plaque characteristics and adverse outcomes.<sup>6-8</sup> In one publication, EAT attenuation was a better predictor of myocardial infarction than EAT volume.<sup>8</sup> A higher CT attenuation may indicate the presence of inflammatory changes and neovascularization in the context of EAT or a different type of adipose tissue in the pericoronary space. If inflammation was the source of such difference in radiodensity, its reversal may be desirable. Statins have been shown to induce a volumetric reduction of EAT, and more aggressive low-density lipoprotein cholesterol (LDL-C) lowering induced a more marked reduction of EAT volume.<sup>9</sup> Whether statins affect the attenuation, hence potentially the inflammatory state of EAT, is unknown. We tested this hypothesis in a subanalysis of a double-blind, randomized, clinical trial designed to investigate the effect of 2 different

From the Mazankowski Alberta Heart Institute and Departments of Medicine (P.R.) and Pediatrics (G.L.), University of Alberta, Edmonton, Alberta, Canada; Division of Cardiology (V.G.) and Department of Biostatistics and Bioinformatics (C.Z., Z.C.), Emory University, Atlanta, GA; and Division of Cardiothoracic Imaging, Department of Radiology and Imaging Sciences, Emory University, Atlanta, GA (A.E.S.).

**Correspondence to:** Paolo Raggi, MD, PhD, Mazankowski Alberta Heart Institute, University of Alberta, 11220 83rd Ave, Ste 5A9-014, Edmonton, Alberta, Canada T6G 2B7. E-mail: raggi@ualberta.ca

Received April 29, 2019; accepted May 23, 2019.

© 2019 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

## Clinical Perspective

### What Is New?

- New evidence suggests that high epicardial adipose tissue attenuation on computed tomography imaging is a marker of inflammation and may promote development of atherosclerosis; however, brown adipose tissue is also present in this fat depot and it has a higher radiological attenuation.

### What Are the Clinical Implications?

- We demonstrated that statins reduce the attenuation of epicardial adipose tissue independent of reduction in low-density lipoprotein levels.
- This may be caused by an anti-inflammatory and a positive pleiotropic effect of statins, or it could reflect inhibition of brown adipose tissue physiological features by these drugs; in this case, it could represent a negative influence of statins.

statins to slow progression of coronary artery calcium (CAC).<sup>10</sup>

## Methods

The data that support the findings of this study are available from the corresponding author on reasonable request.

## Study Population

Details of the study protocol have been published before and are summarized in brief here. The purpose of the BELLES (Beyond Endorsed Lipid Lowering With Electron Beam Tomography Scanning) trial<sup>10</sup> was to investigate whether CAC progression can be slowed more effectively with aggressive lipid lowering treatment with 80 mg daily of atorvastatin than with a more moderate LDL lowering attained with 40 mg daily of pravastatin. Of the 615 enrolled postmenopausal women with dyslipidemia, 475 underwent both a baseline and a follow-up CT scan at a 1-year interval and formed the core of the original report. In this subanalysis, we reviewed all the stored CT scans of sufficient quality to allow a detailed analysis of EAT attenuation. We were able to review 2 CT scans for each of 420 postmenopausal women.

The clinical characteristics of the 55 patients excluded from these analyses, because of poor quality CT images, were identical to those of the patients included. Postmenopausal state was defined as either surgically induced or spontaneous loss of menstrual cycle for >6 months before enrollment. Dyslipidemia was defined as a fasting serum level of LDL-C >160 mg/dL for women with a 10-year Framingham risk

score <10% or LDL-C >130 mg/dL for women with known coronary heart disease or a 10-year Framingham risk score >10%. All women had to have a minimum baseline CAC score of 30 to be included in the study; therefore, by definition, all patients had subclinical coronary atherosclerosis.

During the screening visit, patients were interviewed about their medical history; and a detailed analysis of their coronary artery disease risk, inclusive of laboratory testing, was performed.

After meeting the inclusion criteria, patients became eligible for the baseline chest CT scan and were randomized in a double-blind approach to either atorvastatin, 80 mg/d, and matching pravastatin placebo or pravastatin, 40 mg/d, and matching atorvastatin placebo. The second CT scan was performed after 12 months from randomization. The study was approved by the ethics review committee of each participating institution, and the subjects gave informed consent to participate.


## CT Imaging for CAC and EAT Assessment

Imatron (GE Imatron, San Francisco, CA) C-150 electron beam CT scanners were used to perform the baseline and follow-up scans for all patients using a standard imaging protocol.<sup>10</sup> Scanning started at the bronchial carina and extended to the diaphragm for a total of 36 to 40 slices. The CAC score was calculated as a volume score, as previously described.<sup>11</sup>

The EAT volume and attenuation were measured on the same axial images used for CAC scoring. The EAT volume was calculated using the Volume Analysis software of a Leonardo workstation (Siemens, Erlangen, Germany), as previously described.<sup>9</sup> The EAT volume was assessed after manually tracing the epicardium in each successive axial slice in the craniocaudal direction, using a threshold of −190 to −30 Hounsfield units (HUs). The fat voxels from each slice were then summed to obtain the total EAT volume (in mL).

The EAT attenuation was measured in HUs in the vicinity of the proximal right coronary artery (Figure 1). A region of interest was drawn in the EAT located in the space separating the right atrium from the right ventricular outflow tract. The region of interest was drawn in an area not contiguous with CAC and not affected by streak or motion artifacts. The location and surface area of the region of interest were kept identical for the baseline and follow-up scans. Similar procedures were followed to measure SubQ HUs. In this case, the region of interest was drawn in an area of thoracic subcutaneous tissue, mostly along the midaxillary line (Figure 1).

The same experienced investigator (G.V.) performed all measurements of adipose tissue attenuation. The same investigator also performed repeated measurements of attenuation on a random sample of 100 CT scans selected



**Figure 1.** Example of regions of interest used to measure epicardial adipose tissue (EAT) and subcutaneous adipose tissue (SubQ) attenuation.

in a blinded manner. The intrareader variability of the measurements was <1%, demonstrating a high degree of reproducibility.

### Statistical Analysis

Mean and SD were calculated for normally distributed numerical variables; medians were calculated for nonnormally distributed variables. Frequency and percentages were used to describe categorical variables. The variables CAC, EAT volume, EAT HUs, and SubQ HUs were measured at baseline and at 1 year from randomization in all patients. The percentage change for each parameter was calculated as follows:  $[(\text{final}-\text{baseline})/\text{baseline}]*100\%$ . One-way ANOVA test and Kruskal-Wallis test were used to compare numerical variables in univariable analyses, as appropriate. The paired *t*-test was used to compare baseline and follow-up CAC, EAT volume, EAT HUs, and SubQ HUs. The Pearson correlation coefficient was used to test the association between percentage EAT HU change and percentage change in CAC score, EAT volume, SubQ HUs, total cholesterol, high-density lipoprotein cholesterol (HDL-C), LDL-C, and total triglycerides. Differences in categorical variables between treatment arms were compared by means of  $\chi^2$  test or Fisher's exact test, where appropriate. A multivariate linear regression model was fit by a backward variable selection method with an  $\alpha=0.20$  removal criterion, to estimate the association of EAT HU change in the whole group, the atorvastatin group, and the pravastatin group with age, race, diabetes mellitus, hypertension, smoking, baseline use of hormone replacement therapy, body mass index, and history of cardiovascular disease (angina, coronary artery angioplasty or bypass, or peripheral vascular disease). Scatterplots were drawn to show the relationship between percentage EAT HU change and percentage change in LDL-C and HDL-C in the entire cohort. The

same analyses performed for EAT HU change were applied to assess change in SubQ HU. The significance level was set at  $P<0.05$ . The SAS statistical package, version 9.4 (SAS Institute, Inc, Cary, NC), was used for data management and analysis.

### Results

The clinical characteristics of the 420 patients enrolled in this study are shown in Table 1. The randomization was effective, and there were no significant clinical or demographic differences between patients in the 2 treatment arms. The characteristics of the 55 patients excluded because of poor image quality were identical to the 420 patients included. The LDL-C and triglycerides lowering of atorvastatin were superior to those of pravastatin ( $48\pm 18\%$  versus  $25\pm 18\%$  [ $P<0.001$ ] and  $22.8\pm 33\%$  versus  $9.7\pm 30\%$  [ $P<0.001$ ], respectively). There was no significant difference in the HDL-C increasing effect between the 2 statins. We previously demonstrated that both statins reduced the EAT volume from baseline, although atorvastatin was more effective than pravastatin.<sup>9</sup> Similarly, statins reduced (made more negative) the EAT attenuation ( $-89.4\pm 24$  HUs at baseline versus  $-94.7\pm 21$  HUs at follow-up; average change for the entire cohort,  $-5.4\pm 29.7$  HUs [ $-6\%$  change];  $P<0.001$ ). However, the 2 statins were equally effective and there was no significant difference in the EAT attenuation change induced by treatment (atorvastatin versus pravastatin,  $-6.35\pm 31$  HUs versus  $-4.55\pm 28$  HUs;  $P=0.55$ ). Among the clinical categorical variables, only diabetes mellitus was marginally correlated with change in EAT attenuation ( $P=0.067$ ). Among the continuous variables, percentage change in EAT attenuation was marginally correlated with percentage change in HDL-C ( $P=0.073$ ; Table 2), but no other variable. Figures 2 and 3 show the correlation between change in EAT HUs and change in LDL-C and HDL-C for the entire cohort.

The multivariable model for the entire cohort did not produce any variable significantly associated at the 0.05 level with change in EAT attenuation. The only 2 variables marginally associated with EAT attenuation change were age and percentage change in HDL-C ( $P=0.084$  and  $P=0.086$ , respectively). Although there were no significant predictors of change in EAT attenuation in the pravastatin group, in the atorvastatin group, history of diabetes mellitus ( $\beta=18.8$ ;  $P=0.039$ ), age ( $\beta=-0.94$ ;  $P=0.045$ ), and change in HDL-C ( $\beta=-0.64$ ;  $P=0.003$ ) were predictive of EAT change.

The subcutaneous tissue of the 420 patients included in this subanalysis had a lower attenuation at baseline than EAT ( $-123\pm 30$  versus  $-89\pm 24$  HUs;  $P<0.001$ ). This observation supports the notion that the tissue composition of the 2 fat compartments is substantially different. Treatment with

**Table 1.** Patients' Baseline Clinical Characteristics by Treatment Arm

Covariate	Statistics	Level	Treatment		P Value
			Atorvastatin (N=194)	Pravastatin (N=226)	
Age, y	Mean±SD		65.2±6.5	65.5±6.0	0.548
Race	No. (%)	White	175 (90)	209 (93)	0.606
	No. (%)	Black	10 (5)	8 (3.6)	
	No. (%)	Other	9 (4.6)	8 (3.6)	
HRT	No. (%)	Yes	46 (24)	51 (23)	0.781
Hypertension	No. (%)	Yes	75 (39)	98 (43)	0.329
Diabetes mellitus	No. (%)	Yes	26 (13.4)	35 (15.5)	0.546
Prior MI	No. (%)	Yes	5 (2.6)	4 (1.8)	0.738
Prior CABG	No. (%)	Yes	1 (0.5)	2 (0.9)	1.000
Prior angina	No. (%)	Yes	14 (7.2)	15 (6.4)	0.815
Prior PVD	No. (%)	Yes	11 (6)	18 (8)	0.355
Smoking	No. (%)	Never	80 (41)	95 (42)	0.748
	No. (%)	Ex/current	114 (59)	131 (58)	
BMI, kg/cm <sup>2</sup>	Mean±SD		28.6±5.5	29.2±5.8	0.194
Total cholesterol, mmol/L	Mean±SD		6.95±0.95	6.88±0.98	0.491
HDL-C, mmol/L	Mean±SD		1.47±0.36	1.5±0.36	0.357
LDL-C, mmol/L	Mean±SD		4.55±0.88	4.47±0.88	0.332
Total triglycerides, mmol/L	Mean±SD		2.1±1.0	2.0±1.1	0.419
EAT volume, mL	Mean±SD		111.3±46.8	110.3±43.2	0.950
EAT HUs	Mean±SD		−88.07±26	−90.73±22	0.097
SubQ HUs	Mean±SD		−121.8±32	−124.7±28	0.275
CAC score	Mean±SD		262.9±386	348.9±549	0.071

BMI indicates body mass index; CABG, coronary artery bypass grafting; CAC, coronary artery calcium; EAT, epicardial adipose tissue; HDL-C, high-density lipoprotein cholesterol; HRT, hormone replacement therapy; HU, Hounsfield unit; LDL-C, low-density lipoprotein cholesterol; MI, myocardial infarction; PVD, peripheral vascular disease; SubQ, subcutaneous adipose tissue.

statins did not affect the subcutaneous tissue attenuation after 1 year from randomization (average for the entire cohort,  $-123.3\pm 30$  HUs at baseline versus  $-123\pm 21$  HUs at follow-up;  $P=0.865$ ).

## Discussion

In this subanalysis of a randomized clinical trial, we report, for the first time, a change in EAT attenuation under treatment with statins that was independent of their effect on serum lipid levels, with the exception of a borderline association between EAT HU decrease and HDL-C increase. The SubQ in the same patients had a lower radiodensity at baseline and was not affected by treatment with statins.

The interaction of EAT with the cardiovascular system in health and disease is complex. EAT can be both a friend (providing thermogenic and nourishing support to the coronary arteries and myocardium) and a foe. The influence of

obesity,<sup>12</sup> systemic inflammation, hyperlipidemia,<sup>13</sup> and smoking<sup>14</sup> can promote a phenotypic change in the EAT adipocytes and induce a secretome responsible for the production of proinflammatory cytokines that replace the anti-inflammatory adiponectin. In this new state, EAT and visceral fat are believed to have both paracrine and autocrine activities that may promote atherosclerosis and myocardial fibrosis.<sup>13</sup>

The thickness and volume of EAT have been associated with presence and severity of coronary artery disease<sup>15,16</sup> and adverse outcomes.<sup>17–19</sup> Perivascular adipose tissue accumulation is associated with plaque size and adventitial inflammation.<sup>20,21</sup> More recently, EAT attenuation has been associated with a high prevalence of traditional risk factors and the presence of high-risk atherosclerotic plaques.<sup>6–8</sup> Murphy et al<sup>22</sup> and Antonopoulos et al<sup>23</sup> showed that higher visceral fat density was associated with smaller and less differentiated adipocytes. In addition, higher visceral adipose tissue attenuation and pericoronary artery adipose tissue

**Table 2.** Continuous Variables Associated With Change in EAT Attenuation in the Entire Patient Cohort


Change	Pearson CC	Pearson P Value
EAT volume	-0.003	0.954
SubQ HUs	-0.004	0.941
Total cholesterol	-0.063	0.241
HDL-C	-0.096	0.073
LDL-C	-0.042	0.433
Triglycerides	-0.034	0.526
CAC	0.029	0.570

All parameters are expressed as percentage change. CAC indicates coronary artery calcium; CC, correlation coefficient; EAT, epicardial adipose tissue; HDL-C, high-density lipoprotein cholesterol; HU, Hounsfield unit; LDL-C, low-density lipoprotein cholesterol; SubQ, subcutaneous adipose tissue.


attenuation have been associated with an increased risk of all-cause and cardiovascular mortality.<sup>22,24</sup> A higher fat attenuation has long been interpreted as a surrogate marker of inflammation, after an initial report that visceral fat showed an increased radiodensity in patients with pancreatitis.<sup>25</sup> Accordingly, recently, Antonopoulos et al<sup>23</sup> reported that

higher pericoronary artery attenuation, measured with a novel CT method, reflects a state of inflammation in the context of the vessel wall.

Applying this novel imaging method in a derivation and validation study, Oikonomou et al<sup>24</sup> confirmed that this index of vascular inflammation is predictive of all-cause and cardiac mortality, after adjustment for several clinical characteristics and risk factors. Both in experimental animals<sup>26,27</sup> and in human observations,<sup>28</sup> statins have been shown to have a direct anti-inflammatory effect on adipose tissue. Parisi et al<sup>28</sup> showed a direct correlation between EAT thickness and inflammasome expression within the EAT and further demonstrated an in vitro anti-inflammatory effect of atorvastatin on cultured EAT adipocytes. Atorvastatin had no effect on subcutaneous tissue, in line with our observations in the present study. We previously showed that statins reduced the volume of EAT<sup>9</sup> and suggested that a reduction of inflammation and/or partial disappearance of vasa vasorum in the perivascular fat surrounding the coronary arteries<sup>29</sup> may have contributed to our findings. If interpreted in this light, the current results can be seen as a positive pleiotropic action of statins on EAT. In fact, a reduction in tissue attenuation could reflect a reduction in EAT inflammation, hence an



**Figure 2.** Percentage change in epicardial adipose tissue attenuation vs percentage low-density lipoprotein cholesterol (LDL-C) change for the entire study population. EAT indicates epicardial adipose tissue; HU, Hounsfield unit.



**Figure 3.** Percentage change in epicardial adipose tissue attenuation vs percentage high-density lipoprotein cholesterol (HDL-C) change for the entire study population. EAT indicates epicardial adipose tissue; HU, Hounsfield unit.

antiatherosclerotic action of statins. However, there is at least one other explanation for our findings that needs careful consideration. Imaging and histological studies suggest that the perivascular fat of epicardial coronary arteries is composed of brown adipose tissue (BAT) with dispersed beige cells (white adipose tissue [WAT] cells that express a BAT phenotype).<sup>30–32</sup> BAT is essential for the production of nonshivering thermogenesis via the action of several enzymes, the most important of which is UCP-1 (uncoupling protein-1). By burning large quantities of free fatty acids and glucose, BAT contributes to better cardiometabolic health overall, improved glucose homeostasis, and insulin sensitivity.<sup>33,34</sup> Indeed, its stimulation with a  $\beta$ -3 adrenergic receptor agonist in mice stymied the development of experimental atherosclerosis.<sup>35</sup> The adipocytes found in BAT contain more mitochondria and are smaller than those of WAT. This is likely due to the fact that they have smaller triacylglycerol stores because BAT cells oxidize rather than store fatty acids. Furthermore, activation of BAT induces a several fold increase in blood flow through it, for the purpose of delivering free fatty acids and glucose for consumption. The end result is a higher radiological attenuation of BAT than WAT,<sup>36,37</sup> both at rest and especially after stimulation. A surprising recent study suggested that statins may reduce the expression of genes

necessary for thermogenesis within BAT and that the mechanism may reside in inhibition of the geranylgeranyl pyrophosphate cascade.<sup>38</sup> Balaz et al<sup>38</sup> reported that hydroxy methyl-glutaryl coenzyme A synthase 2 is highly expressed in human BAT and is correlated with expression of UCP-1. Blockade of the hydroxy methyl-glutaryl coenzyme A synthase 2 with statins, both in murine and in in vitro adipose tissue experiments, led to a marked decrease in the expression of UCP-1, hence a reduction in the thermogenetic potential of BAT. In in vivo murine experiments, Balaz et al<sup>38</sup> further demonstrated that fluvastatin and simvastatin inhibited “browning of WAT,” essentially inhibiting a phenotypic transformation of WAT adipocytes into BAT adipocytes. Of interest, the negative influence of statins on UCP-1 expression could be rescued with geranylgeranyl pyrophosphate, suggesting a non-cholesterol-related action of statins on BAT. In a retrospective analysis of 738 patients submitted to multiple positron emission tomography/CT scans with <sup>18</sup>F-fluorodeoxyglucose over a year’s time, Balaz et al<sup>38</sup> further showed a lower BAT activity in patients receiving statins compared with statin-naïve patients. Finally, in 16 volunteers with high baseline BAT activity, 2 weeks of fluvastatin therapy effectively reduced total cholesterol and LDL-C levels, but it also blocked UCP-1 expression, whereas there was no difference

in expression of inflammatory markers. Taken together, these data suggest that what we observed in our analyses potentially was a loss of activity of BAT that resulted in reduction in the CT attenuation of EAT. Although the deactivation of BAT may be a negative collateral effect of statins, these drugs obviously have a proven record of cardiovascular safety. Whether the inhibition of BAT and browning of WAT are a potential other pathway leading to the development of diabetes mellitus in patients receiving statins remains to be studied in detail. In obesogenic states, even BAT can become inflamed.<sup>39</sup> In addition, experiments in apolipoprotein E knockout mice showed that stimulation of BAT with cold induced a striking increase in hepatic production of hydroxy methyl-glutaryl coenzyme A reductase and serum LDL-C, intermediate-density lipoprotein cholesterol, and very low density lipoprotein-cholesterol.<sup>40</sup> The end result was promotion of atherosclerosis development, apparently via an adiponectin-sensitive mechanism. Hence, it is possible that BAT itself may become proatherogenic in some disease states, such as hypercholesterolemia and obesity.<sup>41</sup> Finally, the separation of WAT, beige adipose tissue, and BAT into separate depots may be somewhat artificial; and it has been proposed that adipocytes of all types are likely present in every compartment.<sup>42</sup>

The SubQ in our patients had a lower attenuation at baseline and did not respond to treatment with statins compared with EAT. These observations support the notion that subcutaneous fat is mainly constituted of WAT, and it may not be as inflamed as EAT. In their original report, Mazurek et al<sup>1</sup> noted that the SubQ of patients referred for elective coronary artery bypass surgery was not inflamed, whereas the EAT expressed a large number of proinflammatory mRNAs and was densely infiltrated with monocytes and lymphocytes. Our observation appears to be consistent with theirs.

A few limitations of the current report should be highlighted. This was a subanalysis of a previously randomized clinical trial in postmenopausal women, with hypercholesterolemia and CAC<sup>10</sup>; hence, our findings cannot be automatically extended to other patient groups. In the parent trial, we did not collect serum markers of inflammation. Other investigators reported an anti-inflammatory action of statins on EAT, but without histological confirmation our observations remain hypothesis generating. Patients were randomized to 2 different statins, but there was no placebo arm. We measured EAT and SubQ attenuation in a relatively small region of interest, although we made it as large as possible and we also ensured that the areas chosen were free from artifacts. Selecting smaller regions of interest in the pericoronary space would have increased the SD and the error in the measurement.

This study also had a few positive connotations. The randomization followed strict criteria, and the 2 groups were

perfectly matched. All imaging studies were performed with the same CT scanner brand and model, thus avoiding variability caused by different types of image reconstruction and x-ray tube voltage.

In summary, we showed that moderate to aggressive treatment with statins reduced the EAT attenuation by  $\approx 6\%$  after 1 year of treatment in a cohort of postmenopausal women with subclinical atherosclerosis and hypercholesterolemia. The SubQ fat attenuation was not affected by statin treatment. Because a higher EAT attenuation has been shown to be associated with an unfavorable outcome, a decrease in attenuation with treatment may represent another beneficial pleiotropic action of statins. On the other hand, whether the changes we described are suggestive of a potential adverse effect of these popular drugs, that in rare cases induce development of diabetes mellitus, will require further clarification.

## Disclosures

None.

## References

- Mazurek T, Zhang L, Zalewski A, Mannion JD, Diehl JT, Arafat H, Sarov-Blat L, O'Brien S, Keiper EA, Johnson AG, Martin J, Goldstein BJ, Shi Y. Human epicardial adipose tissue is a source of inflammatory mediators. *Circulation*. 2003;108:2460–2466.
- Alexopoulos N, McLean D, Janik M, Arepalli C, Stillman AE, Raggi P. Epicardial adipose tissue and coronary artery plaque characteristics. *Atherosclerosis*. 2010;210:150–154.
- Iwasaki K, Urabe N, Kitagawa A, Nagao T. The association of epicardial fat volume with coronary characteristics and clinical outcome. *Int J Cardiovasc Imaging*. 2018;34:301–309.
- Nerlekar N, Brown AJ, Muthalaly RG, Talman A, Hettige T, Cameron JD, Wong DTL. Association of epicardial adipose tissue and high-risk plaque characteristics: a systematic review and meta-analysis. *J Am Heart Assoc*. 2017;6:e006379. DOI: 10.1161/JAHA.117.006379.
- Mahabadi AA, Berg MH, Lehmann N, Kälsch H, Bauer M, Kara K, Dragano N, Moebus S, Jöckel KH, Erbel R, Möhlenkamp S. Association of epicardial fat with cardiovascular risk factors and incident myocardial infarction in the general population: the Heinz Nixdorf Recall Study. *J Am Coll Cardiol*. 2013;61:1388–1395.
- Marwan M, Hell M, Schuhbäck A, Gauss S, Bittner D, Pflederer T, Achenbach S. CT attenuation of pericoronary adipose tissue in normal versus atherosclerotic coronary segments as defined by intravascular ultrasound. *J Comput Assist Tomogr*. 2017;41:762–767.
- Goeller M, Achenbach S, Marwan M, Doris MK, Cadet S, Commandeur F, Chen X, Slomka PJ, Gransar H, Cao JJ, Wong ND, Albrecht MH, Rozanski A, Tamarappoo BK, Berman DS, Dey D. Epicardial adipose tissue density and volume are related to subclinical atherosclerosis, inflammation and major adverse cardiac events in asymptomatic subjects. *J Cardiovasc Comput Tomogr*. 2018;12:67–73.
- Mahabadi AA, Balcer B, Dykun I, Forsting M, Schlosser T, Heusch G, Rassaf T. Cardiac computed tomography-derived epicardial fat volume and attenuation independently distinguish patients with and without myocardial infarction. *PLoS One*. 2017;12:e0183514.
- Alexopoulos N, Melek BH, Arepalli CD, Hartlage GR, Chen Z, Kim S, Stillman AE, Raggi P. Effect of intensive versus moderate lipid-lowering therapy on epicardial adipose tissue in hyperlipidemic post-menopausal women: a substudy of the BELLES trial (Beyond Endorsed Lipid Lowering with EBT Scanning). *J Am Coll Cardiol*. 2013;61:1956–1961.
- Raggi P, Davidson M, Callister TQ, Welty FK, Bachmann GA, Hecht H, Rumberger JA. Aggressive versus moderate lipid-lowering therapy in hypercholesterolemic postmenopausal women: Beyond Endorsed Lipid Lowering with EBT Scanning (BELLES). *Circulation*. 2005;112:563–571.

11. Callister TQ, Cooil B, Raya SP, Lippolis NJ, Russo DJ, Raggi P. Coronary artery disease: improved reproducibility of calcium scoring with an electron-beam CT volumetric method. *Radiology*. 1998;208:807–814.
12. Hajer GR, van Haefthen TW, Visseren FL. Adipose tissue dysfunction in obesity, diabetes, and vascular diseases. *Eur Heart J*. 2008;29:2959–2971.
13. Packer M. Epicardial adipose tissue may mediate deleterious effects of obesity and inflammation on the myocardium. *J Am Coll Cardiol*. 2018;71:2360–2372.
14. Mach L, Bedanova H, Soucek M, Karpisek M, Nemecek P, Orban M. Tobacco smoking and cytokine levels in human epicardial adipose tissue: impact of smoking cessation. *Atherosclerosis*. 2016;255:37–42.
15. Djaberi R, Schuijff JD, van Werkhoven JM, Nucifora G, Jukema JW, Bax JJ. Relation of epicardial adipose tissue to coronary atherosclerosis. *Am J Cardiol*. 2008;102:1602–1607.
16. Ding J, Kritchevsky SB, Harris TB, Burke GL, Detrano RC, Szklo M, Jeffrey Carr J; Multi-Ethnic Study of Atherosclerosis. The association of pericardial fat with calcified coronary plaque. *Obesity (Silver Spring)*. 2008;16:1914–1919.
17. Ding J, Hsu FC, Harris TB, Liu Y, Kritchevsky SB, Szklo M, Ouyang P, Espeland MA, Lohman KK, Criqui MH, Allison M, Bluemke DA, Carr JJ. The association of pericardial fat with incident coronary heart disease: the Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Clin Nutr*. 2009;90:499–504.
18. Raggi P, Zona S, Scaglioni R, Stentarelli C, Ligabue G, Besutti G, Menozzi M, Santoro A, Malagoli A, Bellasi A, Guaraldi G. Epicardial adipose tissue and coronary artery calcium predict incident myocardial infarction and death in HIV-infected patients. *J Cardiovasc Comput Tomogr*. 2015;9:553–558.
19. D'Marco LG, Bellasi A, Kim S, Chen Z, Block GA, Raggi P. Epicardial adipose tissue predicts mortality in incident hemodialysis patients: a substudy of the Renegal in New Dialysis trial. *Nephrol Dial Transplant*. 2013;28:2586–2595.
20. Konishi M, Sugiyama S, Sato Y, Oshima S, Sugamura K, Nozaki T, Ohba K, Matsubara J, Sumida H, Nagayoshi Y, Sakamoto K, Utsunomiya D, Awai K, Jinnouchi H, Matsuzawa Y, Yamashita Y, Asada Y, Kimura K, Umemura S, Ogawa H. Pericardial fat inflammation correlates with coronary artery disease. *Atherosclerosis*. 2010;213:649–655.
21. Verhagen SN, Vink A, van der Graaf Y, Visseren FL. Coronary perivascular adipose tissue characteristics are related to atherosclerotic plaque size and composition: a post-mortem study. *Atherosclerosis*. 2012;225:99–104.
22. Murphy RA, Register TC, Shively CA, Carr JJ, Ge Y, Heilbrun ME, Cummings SR, Koster A, Nevitt MC, Satterfield S, Tykvasky FA, Strotmeyer ES, Newman AB, Simonsick EM, Scherzinger A, Goodpaster BH, Launer LJ, Eiriksdottir G, Sigurdsson S, Sigurdsson G, Gudnason V, Lang TF, Kritchevsky SB, Harris TB. Adipose tissue density, a novel biomarker predicting mortality risk in older adults. *J Gerontol A Biol Sci Med Sci*. 2014;69:109–117.
23. Antonopoulos AS, Sanna F, Sabharwal N, Thomas S, Oikonomou EK, Herdman L, Margaritis M, Shirodaria C, Kampoli AM, Akoumianakis I, Petrou M, Sayeed R, Krasopoulos G, Psarros C, Ciccone P, Brophy CM, Digby J, Kelion A, Uberoi R, Anthony S, Alexopoulos N, Tousoulis D, Achenbach S, Neubauer S, Channon KM, Antoniades C. Detecting human coronary inflammation by imaging perivascular fat. *Sci Transl Med*. 2017;9:eaal2658.
24. Oikonomou EK, Marwan M, Desai MY, Mancio J, Alashi A, Hutt Centeno E, Thomas S, Herdman L, Kotanidis CP, Thomas KE, Griffin BP, Flamm SD, Antonopoulos AS, Shirodaria C, Sabharwal N, Deanfield J, Neubauer S, Hopewell JC, Channon KM, Achenbach S, Antoniades C. Non-invasive detection of coronary inflammation using computed tomography and prediction of residual cardiovascular risk (the CRISP CT study): a post-hoc analysis of prospective outcome data. *Lancet*. 2018;392:929–939.
25. Thornton E, Mendiratta-Lala M, Siewert B, Eisenberg RL. Patterns of fat stranding. *AJR Am J Roentgenol*. 2011;197:W1–W14.
26. Abe M, Matsuda M, Kobayashi H, Miyata Y, Nakayama Y, Komuro R, Fukuhara A, Shimomura I. Effects of statins on adipose tissue inflammation: their inhibitory effect on MyD88-independent IRF3/IFN-beta pathway in macrophages. *Arterioscler Thromb Vasc Biol*. 2008;28:871–877.
27. Yamada Y, Takeuchi S, Yoneda M, Ito S, Sano Y, Nagasawa K, Matsuura N, Uchinaka A, Murohara T, Nagata K. Atorvastatin reduces cardiac and adipose tissue inflammation in rats with metabolic syndrome. *Int J Cardiol*. 2017;240:332–338.
28. Parisi V, Petraglia L, D'Esposito V, Cabaro S, Rengo G, Caruso A, Grimaldi MG, Baldascino F, De Bellis A, Vitale D, Formisano R, Ferro A, Paolillo S, Davin L, Lancellotti P, Formisano P, Perrone Filardi P, Ferrara N, Leosco D. Statin therapy modulates thickness and inflammatory profile of human epicardial adipose tissue. *Int J Cardiol*. 2019;274:326–330.
29. Dong L, Kerwin WS, Chen H, Chu B, Underhill HR, Neradilek MB, Hatsukami TS, Yuan C, Zhao XQ. Carotid artery atherosclerosis: effect of intensive lipid therapy on the vasa vasorum-evaluation by using dynamic contrast-enhanced MR imaging. *Radiology*. 2011;260:224–231.
30. Sacks HS, Fain JN, Bahouth SW, Ojha S, Frontini A, Budge H, Cinti S, Symonds ME. Adult epicardial fat exhibits beige features. *J Clin Endocrinol Metab*. 2013;98:E1448–E1455.
31. Sacks HS, Fain JN, Holman B, Cheema P, Chary A, Parks F, Karas J, Optican R, Bahouth SW, Garrett E, Wolf RY, Carter RA, Robbins T, Wolford D, Samaha J. Uncoupling protein-1 and related messenger ribonucleic acids in human epicardial and other adipose tissues: epicardial fat functioning as brown fat. *J Clin Endocrinol Metab*. 2009;94:3611–3615.
32. Chechi K, Voisine P, Mathieu P, Laplante M, Bonnet S, Picard F, Joubert P, Richard D. Functional characterization of the Ucp1-associated oxidative phenotype of human epicardial adipose tissue. *Sci Rep*. 2017;7:15566.
33. Stanford KI, Middelbeek RJ, Townsend KL, An D, Nygaard EB, Hitchcox KM, Markan KR, Nakano K, Hirshman MF, Tseng YH, Goodyear LJ. Brown adipose tissue regulates glucose homeostasis and insulin sensitivity. *J Clin Invest*. 2013;123:215–223.
34. Chondronikola M, Volpi E, Børshiem E, Porter C, Annamalai P, Enerbäck S, Lidell ME, Saraf MK, Labbe SM, Hurren NM, Yfanti C, Chao T, Andersen CR, Cesani F, Hawkins H, Sidossis LS. Brown adipose tissue improves whole-body glucose homeostasis and insulin sensitivity in humans. *Diabetes*. 2014;63:4089–4099.
35. Berbée JF, Boon MR, Khedoe PP, Bartelt A, Schlein C, Worthmann A, Koopman S, Hoeke G, Mol IM, John C, Jung C, Vazirpanah N, Brouwers LP, Gordts PL, Esko JD, Hiemstra PS, Havekes LM, Scheja L, Heeren J, Rensen PC. Brown fat activation reduces hypercholesterolaemia and protects from atherosclerosis development. *Nat Commun*. 2015;6:6356.
36. Hu HH, Chung SA, Nayak KS, Jackson HA, Gilsanz V. Differential CT attenuation of metabolically active and inactive adipose tissues: preliminary findings. *J Comput Assist Tomogr*. 2011;35:65–71.
37. Baba S, Jacene HA, Engles JM, Honda H, Wahl RL. CT Hounsfield units of brown adipose tissue increase with activation: preclinical and clinical studies. *J Nucl Med*. 2010;51:246–250.
38. Balaz M, Becker AS, Balazova L, Straub L, Müller J, Gashi G, Maushart CI, Sun W, Dong H, Moser C, Horvath C, Efthymiou V, Rachamin Y, Modica S, Zellweger C, Bacanovic S, Stefanicka P, Varga L, Ukropcova B, Profant M, Opitz L, Amri EZ, Akula MK, Bergo M, Ukropec J, Falk C, Zamboni N, Betz MJ, Burger IA, Wolfrum C. Inhibition of mevalonate pathway prevents adipocyte browning in mice and men by affecting protein prenylation. *Cell Metab*. 2019;29:901–916.e8.
39. Villarroya F, Cereijo R, Gavaldà-Navarro A, Villarroya J, Giral M. Inflammation of brown/beige adipose tissues in obesity and metabolic disease. *J Intern Med*. 2018;284:492–504.
40. Dong M, Yang X, Lim S, Cao Z, Honek J, Lu H, Zhang C, Seki T, Hosaka K, Wahlberg E, Yang J, Zhang L, Länne T, Sun B, Li X, Liu Y, Zhang Y, Cao Y. Cold exposure promotes atherosclerotic plaque growth and instability via UCP1-dependent lipolysis. *Cell Metab*. 2013;18:118–129.
41. Lohmann C, Schäfer N, von Lukowicz T, Sokrates Stein MA, Borén J, Rützi S, Wahl W, Donath MY, Lüscher TF, Matter CM. Atherosclerotic mice exhibit systemic inflammation in periadventitial and visceral adipose tissue, liver, and pancreatic islets. *Atherosclerosis*. 2009;207:360–367.
42. Cinti S. The adipose organ. *Prostaglandins Leukot Essent Fatty Acids*. 2005;73:9–15.