



EMORY
LIBRARIES &
INFORMATION
TECHNOLOGY

OpenEmory

COVID-19 and Shock: A Cautionary Tale for Elderly Patients From a Pooled Analysis

R Desai, *Atlanta VA Med Ctr*
S Singh, *University of Amsterdam*
T Parekh, *George Mason University*
Sonali Sachdeva, *Lady Hardinge Medical College*
[Lekshmi Kumar](#), *Emory University*
[Rajesh Sachdeva](#), *Emory University*
[Gautam Kumar](#), *Emory University*

Journal Title: Annals of Emergence Medicine

Volume: Volume 75, Number 6

Publisher: Mosby-Elsevier | 2020-06-01, Pages 789-791

Type of Work: Article | Final Publisher PDF

Publisher DOI: 10.1016/j.annemergmed.2020.04.014

Permanent URL: <https://pid.emory.edu/ark:/25593/vjd75>

Final published version: <http://dx.doi.org/10.1016/j.annemergmed.2020.04.014>

Copyright information:

© 2020 by the American College of Emergency Physicians.

Accessed September 25, 2023 12:10 AM EDT



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Even in the early years of the 21st century, lessons can be learned in regard to adverse events of mass vaccination. The association of myopericarditis after smallpox (variola major) vaccination in preparation for a possible smallpox event from December 2002 to March 2003 noted an incidence of myopericarditis of 1 per 12,819 primary vaccines within 30 days.⁴ This was associated with an odds ratio of 3.6 (95% confidence interval 3.33 to 4.11) compared with that for unvaccinated individuals.

With the current and future pandemics caused by emerging respiratory pathogens, it would be wise to evaluate interventions with possible consequences.⁵ Attempting to modify hypothetical risks can lead to further risks, which may lead to collateral toxicity that was originally unforeseen at the initial intervention. Evidence-based medicine (whenever possible) with an appreciation of medical history should guide our risk-benefit decisional evaluations in the near future.

Jerrold B. Leikin, MD
NorthShore University HealthSystem—OMEGA
Glenview, IL

<https://doi.org/10.1016/j.annemergmed.2020.04.013>

Funding and support: By *Annals* policy, all authors are required to disclose any and all commercial, financial, and other relationships in any way related to the subject of this article as per ICMJE conflict of interest guidelines (see www.icmje.org). The author has stated that no such relationships exist.

1. Starko KM. Salicylates and pandemic influenza mortality 1918-1919 pharmacology, pathology and historic evidence. *Clin Infect Dis*. 2009;49:1405-1410.
2. Evans D, Cauchemez S, Hayden FG. "Prepandemic" influenza for novel influenza viruses, "swine flu" vaccine, Guillain-Barre syndrome, and detection of rare severe adverse events. *J Infect Dis*. 2009;200:321-328.
3. Starko KM, Ray CG, Dominguez BS, et al. Reye's syndrome and salicylate use. *Pediatrics*. 1980;66:859-864.
4. Halsell JS, Riddle JR, Atwood JE, et al. Myopericarditis following smallpox vaccination among vaccinia-naïve US military personnel. *JAMA*. 2003;289:3183-3289.
5. McFee RB. Selected epidemics and emerging pathogens. *Dis Mon*. 2017;63:235-278.

COVID-19 and Shock: A Cautionary Tale for Elderly Patients From a Pooled Analysis



To the Editor:

The shocking scale of coronavirus disease 2019 (COVID-19) infections is worrisome, with more than 1 million

confirmed cases and greater than 50,000 reported deaths across the globe by the end of March 2020. The unprecedented challenges brought by the COVID-19 pandemic have overwhelmed the health care system, strained health care workers, and raised a dire need to collect, analyze, and interpret real-time data to expedite understanding the etiopathogenesis, risk factors, and prognosis of COVID-19 and ways to curtail overall mortality rate.

Although the awareness of the natural progression of COVID-19 is increasing, with cardiovascular risk factors and older age being frequently identified as major risk factors for poor survival in COVID-19 patients, our knowledge remains limited on the pooled prevalence of shock and its effect on predicting mortality in COVID-19 infection. Although the predominant complication of COVID-19 is acute respiratory illness that could lead to acute respiratory distress syndrome, COVID-19 patients with cardiovascular complications and sepsis have a heightened risk of developing shock with potential in-hospital mortality during the disease course. Acute cardiac injury has been reported to range from 12% to 16.7% in COVID-19 patients.

To our knowledge, this is the first pooled estimate of the prevalence of shock in COVID-19 patients with age-based variation (mean age <50 versus >50 years). Random-effects models were obtained to perform a meta-analysis, and I^2 statistics were used to measure interstudy heterogeneity. After a thorough review of the literature from PubMed, Scopus, and Google Scholar, data were collected from 5 studies (4/5 septic shock) reported from China until March 2020.¹⁻⁵ Among 1,578 COVID-19 patients, the pooled prevalence of shock was 6.3% (95% confidence interval 4.0% to 16.8%) and $I^2=96%$, with the cohort older than 50 years (mean age) showing significantly higher prevalence of shock (9.5%; 95% confidence interval 2.4% to 20.5%) compared with the younger cohort (mean age <50 years) (3.3%; 95% confidence interval 0% to 11.4%) (Figure).

In this meta-analysis, the elderly population had a higher burden of shock compared with the younger cohort. The predilection of elderly patients toward sepsis and a higher burden of cardiovascular diseases owing to higher comorbidities could be a major reason for the observed age-related disparity in the prevalence of shock. This warrants stricter precautions and social distancing for the geriatric population.

A potential link for shock in COVID-19 could be due to elevated cytokine levels. During a viral illness, the up-regulation of cytokines such as interferon gamma and interleukin (IL)-10 potentiates the vasodilation leading to shock. Similarly, IL-1 β , 6, and 8; monocyte chemoattractant protein-1; and plasminogen activator inhibitor-1 levels are increased in the acute phase of sepsis,




Figure. Random-effects pooled prevalence of shock among COVID-19 patients stratified by mean age.

reflecting endothelial injury. Viral infections could trigger a hyperinflammatory state, such as cytokine-storm syndrome, as featured in COVID-19 patients, which could contribute to developing shock with a potential for multiorgan dysfunction.

Concisely, this meta-analysis highlights the worse effect of COVID-19 in an older age group (9.5% versus 3.3%) compared with the younger cohort. More data are required on the prevalence of shock, its predictors, and their effect on the survival of elderly COVID-19 patients to effectively maneuver supportive resuscitation measures on time. Furthermore, elderly survivors requiring prolonged nursing care and ventilator support might add to the magnitude of complications and economic liability during this global health care crisis.

Rupak Desai, MBBS
Division of Cardiology
Atlanta VA Medical Center
Decatur, GA

Sandeep Singh, MBBS
Division of Epidemiology, Biostatistics, and Bioinformatics
Amsterdam University Medical Center
Amsterdam, Netherlands

Tarang Parekh, MBBS, MSc
Department of Health Administration and Policy
George Mason University
Fairfax, VA

Sonali Sachdeva, MBBS
Department of Medicine
Lady Hardinge Medical College
New Delhi, India

Lekshmi Kumar, MD, MPH
Department of Emergency Medicine
Emory University School of Medicine
Atlanta, GA

Rajesh Sachdeva, MD
Division of Cardiology
Atlanta VA Medical Center
Decatur, GA
Division of Cardiology
Medical College of Georgia
Augusta, GA
Division of Cardiology
Morehouse School of Medicine
Atlanta, GA

Gautam Kumar, MD
Division of Cardiology
Atlanta VA Medical Center
Decatur, GA
Division of Cardiology
Emory University School of Medicine
Atlanta, GA

<https://doi.org/10.1016/j.annemergmed.2020.04.014>

Funding and support: By *Annals* policy, all authors are required to disclose any and all commercial, financial, and other relationships in any way related to the subject of this article as per ICMJE conflict of interest guidelines (see www.icmje.org). The authors have stated that no such relationships exist.

1. Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*. 2020;395:497-506.
2. Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. *JAMA*. 2020;323:1061-1069.
3. Chen T, Wu D, Chen H, et al. Clinical characteristics of 113 deceased patients with coronavirus disease 2019: retrospective study. *BMJ*. 2020;368:m1091.
4. Guan WJ, Ni ZY, Hu Y, et al. Clinical characteristics of coronavirus disease 2019 in China. *N Engl J Med*. 2020;382:1708-1720.
5. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395:1054-1062.

Fighting COVID-19 Hypoxia With One Hand Tied Behind Our Back: Blanket Prohibition of High-Flow Oxygen and Noninvasive Positive End-Expiratory Pressure in US Hospitals



To the Editor:

Before the coronavirus disease 2019 (COVID-19) pandemic, patients with hypoxia who were failing low-flow oxygen through nasal cannula were treated with noninvasive positive-pressure ventilation or high-flow nasal cannula oxygen, aimed at delivering higher concentration and flow of oxygen to match patient demand, decreasing anatomic dead space by preventing rebreathing, and recruiting alveoli by using positive end-expiratory pressure. Many emergency physicians began their fight against COVID-19 with neither of these options because of concerns about virus aerosolization exposure to staff and other patients. Instead, early intubation has been the suggested option for a patient failing nasal cannula. Often this is the correct answer, but not always.

There is a cost to staff and patients from overuse of intubation. It is an extremely high-aerosol-generating event. A systematic review of aerosol-generating procedures and their risk of transmission of severe acute respiratory syndrome to health care workers found intubation to have an odds ratio of 6.6 compared with 2.2 for noninvasive positive-pressure ventilation.¹ Also, many hospitals face the risk of running out of ventilators and ICU beds. Once intubated, COVID-19 patients tend to continue receiving mechanical ventilation for greater than 1 week, with poor outcomes. In vitro studies have demonstrated less airflow dispersal from high-flow nasal

cannula or continuous positive airway pressure (CPAP) using sealed masks than from nasal cannula.²

Furthermore, data coming from overseas indicate an important role for noninvasive positive-pressure ventilation and high-flow nasal cannula in managing COVID-19 patients. Retrospective data from China demonstrate that approximately 21% of patients required high-flow nasal cannula and 14% required noninvasive positive-pressure ventilation.³ Among admitted patients in Italy, approximately 30% required ventilation support beyond oxygen therapy. Of patients given ventilation support, 89% were assisted with noninvasive positive-pressure ventilation compared with 12% receiving invasive ventilation.⁴ The *Handbook of COVID-19 Prevention and Treatment* recommends the use of high-flow nasal cannula for hypoxic patients not tolerating nasal cannula.⁵

We advocate early planning with colleagues from critical care, respiratory therapy, and nursing to develop protocols that mitigate risk associated with high-flow nasal cannula and noninvasive positive-pressure ventilation rather than prohibiting the use of these critical alternatives to intubation. Ideally, these interventions should be performed in negative-airflow rooms, but most emergency departments have a limited number of these. We are recommending use of these interventions in closed isolation rooms with staff in full airborne personal protective equipment. High-flow nasal cannula should be provided with a surgical mask over the patient's mouth, or high-flow oxygen could be provided with a nonvented sealed CPAP mask attached to a self-inflating bag plus viral filter with dual oxygen source (see <https://emcrit.org/emcrit/covid-airway-management/>). CPAP should be provided through a helmeted setup or nonvented CPAP masks with viral filter attached to the expiratory port. Policies can be rapidly adapted as more data emerge in regard to COVID-19, but it is already clear that we must find safe and creative ways to expand, not limit, our armamentarium during this pandemic.

Joseph Colla, MD

Adam Rodos, MD

University of Illinois Department of Emergency Medicine
Chicago, IL

Hannah Seyller, BS

University of Illinois at Chicago College of Medicine
Chicago, IL

Scott Weingart, MD

Emergency Critical Care

Department of Emergency Medicine