



EMORY
LIBRARIES &
INFORMATION
TECHNOLOGY

OpenEmory

BMI- and age-related associations with mortality from COVID-19

[Thomas Austin](#), *Emory University*
[Bruce Miller](#), *Emory University*

Journal Title: ANAESTHESIA

Publisher: WILEY | 2021-04-11

Type of Work: Article | Final Publisher PDF

Publisher DOI: 10.1111/anae.15478

Permanent URL: <https://pid.emory.edu/ark:/25593/vkzh0>

Final published version: <http://dx.doi.org/10.1111/anae.15478>

Copyright information:

© 2021 Association of Anaesthetists

Accessed November 29, 2022 10:44 PM EST

Correspondence

BMI- and age-related associations with mortality from COVID-19

We congratulate the COVIDSurg and GlobalSurg Collaboratives for their recent article [1], which addresses a fundamental surgical concern during the COVID-19 pandemic. Although the authors rightfully decided on a host of probable confounding factors to include in their regression modelling a priori, we believe an important variable was omitted from the analysis that requires adjustment, namely BMI. Since higher BMIs are associated with a greater risk of contracting COVID-19 [2] and COVID-19-related mortality [3], this variable could potentially complicate the results of this investigation and conclusions based on its findings. Thus, the regression models should account for BMI in order to better ascertain the correlation between timing of surgery after COVID-19 and mortality.

As paediatric anaesthetists, we are also concerned as to whether the results of this study can be generalised to our paediatric practice, because COVID-19-related death occurs at a far lower rate in those aged 0–17 y relative to those who are older [4]. We believe that performing a sensitivity analysis that includes only patients aged < 18 y would be beneficial to paediatric anaesthetists and surgeons in our continual pursuit of optimal patient care and safety.

T. M. Austin 

B. E. Miller

Emory University, Children's Healthcare of Atlanta,
Atlanta, GA, USA

Email: thomas.austin@emory.edu

No competing interests declared.

References

1. COVIDSurg Collaborative and GlobalSurg Collaborative. Timing of surgery following SARS-CoV-2 infection: an international prospective cohort study. *Anaesthesia* 2021. Epub 9 March. <https://doi.org/10.1111/anae.15458>
2. Jung CY, Park H, Kim DW, et al. Association between body mass index and risk of coronavirus disease 2019 (COVID-19): a nationwide case-control study in South Korea. *Clinical Infectious Diseases* 2020. Epub 25 August. <https://doi.org/10.1093/cid/ciaa1257>.
3. Hendren NS, de Lemos JA, Ayers C, et al. Association of body mass index and age with morbidity and mortality in patients hospitalized with COVID-19: results from the American Heart Association COVID-19 Cardiovascular Disease Registry. *Circulation* 2021; **143**: 135–44.
4. Centers for Disease Control and Prevention. Risk for COVID-19 infection, hospitalization, and death by age group. 2021. <https://www.cdc.gov/coronavirus/2019-ncov/covid-data/investigations-discovery/hospitalization-death-by-age.html#footnote01> (accessed 14/03/2021).

doi:10.1111/anae.15478