



EMORY
LIBRARIES &
INFORMATION
TECHNOLOGY

OpenEmory

Contact networks in the emergency department: Effects of time, environment, patient characteristics, and staff role

Douglas Lowery-North, *Emory University*

[Vicki Hertzberg](#), *Emory University*

[Lisa Elon](#), *Emory University*

[Christina Mehta](#), *Emory University*

[George Cotsonis](#), *Emory University*

J Baumgardner, *Emory University*

Journal Title: SOCIAL NETWORKS

Volume: Volume 48, Number

Publisher: ELSEVIER | 2017-01-01, Pages 181-191

Type of Work: Article

Publisher DOI: 10.1016/j.socnet.2016.08.005

Permanent URL: <https://pid.emory.edu/ark:/25593/vn2jk>

Final published version: <http://dx.doi.org/10.1016/j.socnet.2016.08.005>

Accessed December 8, 2021 12:12 PM EST



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Contact networks in the emergency department: Effects of time, environment, patient characteristics, and staff role



Vicki Stover Hertzberg^{a,*}, Jason Baumgardner^a, C. Christina Mehta^a, Lisa K. Elon^a, George Cotsonis^a, Douglas W. Lowery-North^b

^a Department of Biostatistics and Bioinformatics, Emory University, Atlanta, GA, United States

^b Department of Emergency Medicine, Emory University, Atlanta, GA, United States

ARTICLE INFO

Article history:

Available online 23 September 2016

Keywords:

Emergency medicine
Infectious disease
Social network

ABSTRACT

Emergency departments play a critical role in the public health system, particularly in times of pandemic. Infectious patients presenting to emergency departments bring a risk of cross-infection to other patients and staff through close proximity interactions or contacts. To understand factors associated with cross-infection risk, we measured close proximity interactions of emergency department staff and patients by radiofrequency identification in a working emergency department. The number of contacts (degree) is not related to patient demographic characteristics. However, the amount of time in close proximity (weighted degree) of patients with ED personnel did differ, with black patients having approximately 15 min more contact with staff than non-white patients. Patients arriving by EMS had fewer contacts with other patients than patients arriving by other means. There are differences in the number of contacts based on staff role and arrival mode. When crowding is low, providers have the most contact time with patients, while administrative staff have the least. However, when crowding is high, this differential is reversed. The effect of arrival mode is modified by the extent of crowding. When crowding is low, patients arriving by EMS had longer contact with administrative staff, compared to patients arriving by other means. However, when crowding is high, patients arriving by EMS had less contact with administrative staff compared to patients arriving by other means. Our findings should help designers of emergency care focus on higher risk situations for transmission of dangerous pathogens in an emergency department. For instance, the effects of arrival and crowding should be considered as targets for engineering or architectural interventions that could artificially increase social distances.

© 2016 Published by Elsevier B.V.

1. Introduction

1.1. Background

Within the last decade the world was swept by the H1N1 pandemic, beginning with cases in Mexico detected in March 2009 (WH Organization, 2009). The first US cases appeared in California in mid-April 2009 (Ginsberg et al., 2009). Vaccine was rushed into production, and was available in October. In January 2010

the CDC declared vaccine had been successfully made available to targeted populations (Centers for Disease Control and Prevention, 2010). The H1N1 epidemic is of particular interest in academic emergency medicine. It is a time when the public health system, in which emergency departments (EDs) play a key role, should have been operating optimally due to heightened awareness of infection control efforts.

Close proximity interactions (contacts) provide a means for cross infection when a susceptible individual inhales airborne microbes shed as large droplets by an infectious individual for diseases such as influenza and severe acute respiratory syndrome (SARS) (Bridges et al., 2003; Tellier, 2006; Wenzel and Edmond, 2003). One setting where contacts between susceptible and infectious individuals occur frequently is the ED, as was dramatically demonstrated in the 2003 SARS epidemic (Varia et al., 2003). Understanding the temporal, environmental, and individual factors associated with contacts may lead to improved infection control

Abbreviations: ED, emergency department; EHR, electronic health record; EMS, emergency medical services; GI, gastrointestinal; PP, patient with patient; PS, patient with staff; RFID, radiofrequency identification; RTLS, real time location sensing; SARS, severe acute respiratory syndrome; SP, staff with patient; SS, staff with staff.

* Corresponding author at: Nell Hodgson Woodruff School of Nursing, Emory University, 1520 Clifton Rd. NE, Atlanta, GA 30322, United States. Fax: +1 404 727 1370. E-mail address: vhertz@emory.edu (V.S. Hertzberg).

<http://dx.doi.org/10.1016/j.socnet.2016.08.005>

0378-8733/© 2016 Published by Elsevier B.V.

efforts in the ED, which may prove to be particularly important in times of pandemic concern.

1.2. Importance

Recent developments in technology have enabled precise measurement of movements of humans and resulting interactions. Researchers are increasingly using real time location sensing (RTLS) systems such as radiofrequency identification (RFID) systems to measure occurrence, duration, and location of contacts typically defined as distance between two individuals of some small fixed distance such as 1 meter (m) by line of sight (Gundlapalli et al., 2009; Hornbeck et al., 2012; Isella et al., 2011a,b; Lucet et al., 2012; Stehle et al., 2011; Vanhems et al., 2013; Salathe et al., 2010).

Sensors have been used to study the interactions of patients and staff in a hospital ICU (Hornbeck et al., 2012), a hospital pediatric ward (Isella et al., 2011a), a hospital geriatric ward (Vanhems et al., 2013), and a hospital ward in which patients were under airborne precautions (Lucet et al., 2012). Sensor data have been examined in comparison to electronic medical system log-in information in an ED setting to determine if the latter has utility in determining contacts (Gundlapalli et al., 2009). Other settings in which such technology has been used include schools (Stehle et al., 2011; Salathe et al., 2010) and scientific conferences (Isella et al., 2011b). Due to the costs associated with such technology, these investigations of contacts have been, with few exceptions, one-time investigations of short duration, e.g., 1 week or less. Such snapshots may not be representative of relationships throughout time, in particular failing to account for seasonal or even week-to-week variation. One study in high school students (Fournet and Barrat, 2014) for 11 days in two years showed that there was little variation in contact matrices by time of day, between days, and between years. However it is not clear how these results generalize to other populations. In general, the few studies involving longer periods of time report summary network measures for the period that do not permit inference to the general population of such networks. There is little information on contacts of patients as they interact with staff and with other patients. It is unknown if there are differences associated with patient age, sex, race, or clinical syndrome. There is also limited information about contacts of staff as they interact with other staff and with patients, particularly with respect to staff role (i.e., provider, nurse, administrative).

1.3. Goals of this investigation

A better understanding of the nature of contacts may lead to improvements in infection control in the ED. We measured contacts among patients and staff of the ED of a large urban hospital in Atlanta, GA (Lowery-North et al., 2013). In this paper we describe a secondary analysis of those data. In particular we examine here the characteristics of the resulting contact networks, relating nodal- and network-level metrics with shift characteristics such as time (i.e., season of year, weekday vs. weekend, day vs. night) and environment (e.g., volume of patients, percent high acuity patients). We also describe relationships between patient contacts and patient characteristics as well as between staff contacts and staff role.

2. Methods

2.1. Design

This is a prospective study. We measured contacts among patients and staff of the ED of a large urban hospital in Atlanta, GA during 81 randomly selected 12-h shifts between 1 July 2009 and 30 June 2010 (Lowery-North et al., 2013).

2.2. Setting

As detailed in Lowery-North et al. (2013), we installed an RFID system to determine contacts ≤ 1 meter (m) between and within patients and staff in a busy hospital ED of modern design, part of an urban academic center. This is a modern ED with centralized workspaces and walled patient treatment rooms. The ED was designed for 50,000 patient encounters annually; over the year of the study there were over 57,000 patient encounters.

2.3. Participants

We placed permanent RFID tags on all staff, and placed temporary RFID tags on all patients during predetermined, randomly chosen shifts. The protocol was reviewed and approved by the Emory University Institutional Review Board. All ED staff were invited to participate. Staff participation was voluntary and anonymous. All non-incarcerated patients who were not mentally incapacitated were eligible. Patient participation was voluntary as well.

2.4. Observation periods

In this paper we restrict our analysis to data from the first 6 months of the study (35 shifts of the 81 observed). Data limitations that led to restricted analysis on a subset of shifts are as follows: examination of participation by patients and staff across the year showed a significant decline. We attribute staff participation decline to a system failure that did not alert us to battery depletion in permanent tags worn by staff. There is no similar physical reason for decline in patient participation, thus we attribute it to waning abilities of the research team to keep up with a task that was too large for them alone. Biases in estimates of measures of interest can result from missing individuals and their concomitant contacts.

We restricted analyses to shifts in the first 6 months of our observation period for two reasons: (1) the decline in staff participation starts at the beginning of the second half of the year and thus these observations should not be biased by the presence missing data; (2) the H1N1 epidemic swept through the state of Georgia during these first six months.

2.5. Variables

2.5.1. Outcome variables

The RFID data were used to create a contact network for each shift depicting interactions (the edges) between patient and staff participants (the nodes), resulting in weighted and unweighted adjacency matrices (Lowery-North et al., 2013; Newman, 2010). From these matrices, the following node-level measures were calculated for each participant: degree (number of contacts), time-weighted degree (time-weighted contacts), relative degree (number of contacts normalized to the interval (0, 1)), closeness centrality (inverse of the average shortest path to all other individuals; range is 0–1), and eigenvector centrality (how well they were associated with other central individuals) (Newman, 2010). In addition, for each shift, the following network-level measures were calculated from these matrices: density, average path length, diameter, time-weighted diameter, radius, maximum spectral gap, number of weak components, and average clustering coefficient (transitivity) (Newman, 2010).

2.5.2. Participant-level independent variables

The only information associated with staff RFID tag number was role (provider, nurse, other). Providers comprised attending and resident physicians, nurse practitioners, and physician assistants.

Nurses included all non-clerical and non-provider ED staff. Staff roles classified as other were primarily clerical, and these were primarily patient registration.

Individual patient data elements were obtained from the electronic health record (EHR) and included age, race, sex, acuity, and mode of arrival (ambulance vs. other), patient arrival and departure times, and ICD 9 codes. ICD 9 codes associated with the presenting complaint were categorized using the ESSENCE criteria (Lewis et al., 2002) into the following categories: gastrointestinal (GI), respiratory, neurological, chest pain, musculoskeletal, other. Because contact patterns vary over the course of a patient's stay, we categorized the RFID observation period in terms of its overlap with patient arrival and departure times. Specifically we separated patients into four groups as follows: those whose entire ED visit was observed during the shift under observation; those whose ED visit began before the shift under observation and completed during the observation period; those whose ED visit began during the shift under observation and completed after the end of the observation period, and those whose ED visit began before the shift under observation and completed after the end of the observation period.

Based on ED EHRs, for both staff members and patients we calculated the maximum patient census during each individual's observation period, using this information as a measure of ED crowding.

2.5.3. Shift-level independent variables

Data regarding environment, that is, overall patient mix, were summarized for each shift with respect to median age, percent female, percent arriving via EMS, percent with high acuity illness, percent presenting with respiratory chief complaint, and percent presenting with musculoskeletal chief complaint. Temporal variables for each shift included day (7 am – 7 pm) vs. night (7 pm – 7 am) shift, weekday vs. weekend, and season of year (July/August/September vs. October/November/December). In addition we used the maximum patient census during the observation period as another measure of ED crowding.

2.5.4. Control variables

Variables that were inextricably related to network quantities of interest served as control variables. For node-level outcomes, control variables included length of shift, counts of staff and patients observed by RFID in that shift, and length of observation period for each individual. For network-level outcomes, control variables included total number of participants and length of shift.

2.6. Statistical methods

Data were first summarized as descriptive statistics, such as medians and percentages. To understand variability of the networks (the 35 randomly selected shifts), network characteristics were summarized and the distribution of these summary statistics reported. We typically report median values as they are robust to extreme values. In doing so we use the language of “on average” in the sense of a typical value (Stevenson, 2010) rather than the arithmetic mean.

To evaluate whether temporal and environmental characteristics were associated with node- and network-level metrics, a series of multivariable models were analyzed. Models of node level outcomes used a mixed models approach to account for the clustering of observations within each shift using SAS's PROC MIXED (SI Inc., 2008a) or PROC GLIMMIX (SI Inc., 2008b) with a random statement for shifts. Most staff members had observations in multiple shifts, another source of clustering, but addition of this random effect was found to be unnecessary to the models. Normal linear regression was used except for the following dependent variables: for number of weak components, Poisson regression was used; for indicators

of radius >2 or not and diameter >5 or not, logistic regression was used. Regression diagnostic techniques confirmed linearity and lack of multicollinearity. Problematic residual distributions led to the use of bootstrapping (Carroll and Ruppert, 1988) to quantify sampling error, with 95% confidence intervals calculated utilizing 1000 bootstrapped estimates (based on bootstrapped samples of shifts). Before bootstrapping we screened for inclusion of 2-way interaction terms as appropriate: patient arrival mode with patient acuity, ED crowding during individual observation period, and chief complaint; patient acuity with percent population with high acuity, ED crowding during individual observation period, chief complaint, and day/night; day/night with percent population with high acuity and ED crowding during individual observation period; and patient age with chief complaint. Criterion for including interaction terms in a final model was $\alpha = 0.001$.

Node-level outcomes degree, weighted degree, and relative degree were modeled for four subsets determined by type of contact edge and participant type (patient or staff): (1) patient with other patient (PP), (2) among patients only, patient with staff person (PS), (3) among staff only, a staff person with a patient (SP), and (4) staff person with another staff person (SS). This permitted some clarity about the quite different contact experiences of these subsets. It allowed patient characteristics to be examined as predictors in patient-centered models, and staff member's role (provider, nurse, other) to be included in staff-centered models, in addition to the temporal and environmental characteristics and control variables included in all models.

For the PP outcomes the models took the following form:

$$E(m_{ij}) = \beta_0 + \sum_{k=1}^{28} \beta_k X_{kij}$$

for $i = 1, \dots, n_j$, $j = 1, \dots, 35$, where m_{ij} is one of the three measures considered (degree, weighted degree, relative degree) for patient i in shift j ; n_j is the number of patients observed in shift j ; X_{1ij} = age of patient i in shift j ; X_{2ij} = 1 if subject i in shift j is female, 0 otherwise; X_{3ij} = 1 if patient i in shift j is non-black, 0 otherwise; X_{4ij} = 1 if patient i in shift j arrived by emergency medical services (EMS) ambulance; X_{5ij} = patient census if patient i in shift j arrived by EMS, 0 otherwise; X_{6ij} = 1 if patient i in shift j had acuity of stable or non-urgent, 0 otherwise; X_{7ij} = 1 if patient i in shift j had acuity of urgent, 0 otherwise; X_{8ij} = 1 if patient i in shift j had gastrointestinal (GI) chief complaint, 0 otherwise; X_{9ij} = 1 if patient i in shift j had respiratory chief complaint, 0 otherwise; X_{10ij} = 1 if patient i in shift j had neurological chief complaint, 0 otherwise; X_{11ij} = 1 if patient i in shift j had chest pain chief complaint, 0 otherwise; X_{12ij} = 1 if patient i in shift j had other chief complaint, 0 otherwise; X_{13ij} = 1 if the visit for patient i started before and ended during shift j , 0 otherwise; X_{14ij} = 1 if the visit for patient i started during and ended after shift j ; X_{15ij} = 1 if the visit for patient i started before and ended shift j ; X_{16ij} = 1 if shift j was a day shift (7 am–7 pm), 0 otherwise; X_{17ij} = 1 if shift j was a weekend shift, 0 otherwise; X_{18ij} = 1 if shift j occurred in July, August, or September of 2009, and 0 if it occurred in October–December of 2009; X_{19ij} = maximum census of the ED during shift j ; X_{20ij} = median age (years) of patients seen during shift j ; X_{21ij} = percent of patients seen during shift j that were female; X_{22ij} = percent of patients seen during shift j that arrived by EMS; X_{23ij} = percent of patients seen during shift j with urgent acuity class; X_{24ij} = percent of patients seen during shift j that had respiratory chief complaint; X_{25ij} = percent of patients seen during shift j that had musculoskeletal chief complaint; X_{26ij} = length of shift j (hours); X_{27ij} = duration of observation; and X_{28ij} = number of patients in the ED during shift j .

For the PS outcomes the models took the following form:

$$E(m_{ij}) = \beta_0 + \sum_{k=1}^{29} \beta_k X_{kij}$$

for $i = 1, \dots, n_j$, $j = 1, \dots, 35$, where m_{ij} is one of the three measures considered (degree, weighted degree, relative degree) for patient i in shift j ; n_j is the number of patients observed in shift j ; X_{1ij}, \dots, X_{28ij} are as defined above, and X_{29ij} = number of staff working in the ED during shift j .

For the SP outcomes the model took the following form:

$$E(m_{ij}) = \beta_0 + \sum_{k=1}^{18} \beta_k Z_{kij}$$

for $i = 1, \dots, n_j$, $j = 1, \dots, 35$, where m_{ij} is one of the three measures considered (degree, weighted degree, relative degree) for staff i in shift j ; n_j is the number of staff observed in shift j ; $Z_{1ij} = 1$ if staff i in shift j is a provider, 0 otherwise; $Z_{2ij} = 1$ if staff i in shift j is a nurse, 0 otherwise; Z_{3ij} = the census for shift j if staff i is a provider; Z_{4ij} = the census for shift j if staff i is a nurse; $Z_{5ij} = 1$ if shift j is a day shift, 0 otherwise; $Z_{6ij} = 1$ if shift j is a weekend shift, 0 otherwise; $Z_{7ij} = 1$ if shift j occurred in July–September of 2009, 0 if it occurred during October–December of 2009; Z_{8ij} = maximum census of the ED during shift j ; Z_{9ij} = median age (years) of patients seen during shift j ; Z_{10ij} = percent of patients seen during shift j that were female; Z_{11ij} = percent of patients seen during shift j that arrived by EMS; Z_{12ij} = percent of patients seen during shift j with urgent acuity class; Z_{13ij} = percent of patients seen during shift j that had respiratory chief complaint; Z_{14ij} = percent of patients seen during shift j that had musculoskeletal chief complaint; Z_{15ij} = length of shift j (hours); Z_{16ij} = duration observed; Z_{17ij} = number of staff working in the ED during shift j ; Z_{18ij} = number of patients in the ED during shift j .

For the SS outcomes the model took the following form:

$$E(m_{ij}) = \beta_0 + \sum_{k=1}^{18} \beta_k Z_{kij}$$

for $i = 1, \dots, n_j$, $j = 1, \dots, 35$, where m_{ij} is one of the three measures considered (degree, weighted degree, relative degree) for staff i in shift j ; n_j is the number of staff observed in shift j ; Z_{1ij}, \dots, Z_{18ij} are as defined above; the two interaction terms were not significant and the number of patients was not relevant.

For eigenvector and closeness centrality measures, the models took the form

$$E(m_{ij}) = \beta_0 + \sum_{k=1}^{20} \beta_k U_{kij}$$

for $i = 1, \dots, n_j$, $j = 1, \dots, 35$, where m_{ij} is one of the two measures considered for participant i in shift j ; n_j is the number of participants observed in shift j ; $U_{1ij} = 1$ if participant i in shift j is a provider, 0 otherwise; $U_{2ij} = 1$ if participant i in shift j is a nurse, 0 otherwise; $U_{3ij} = 1$ if participant i in shift j is administrative staff, 0 otherwise; U_{4ij} = census maximum in shift j ; U_{5ij} = census maximum if participant i in shift j is a provider, 0 otherwise; U_{6ij} = census maximum if participant i in shift j is a nurse; U_{7ij} = census maximum if participant i in shift j is administrative staff, 0 otherwise; $U_{8ij} = 1$ if shift j was a day shift (7 am to 7 pm), 0 otherwise; $U_{9ij} = 1$ if shift j was a weekend shift, 0 otherwise; $U_{10ij} = 1$ if shift j was occurred in July, August, or September of 2009, and 0 if it occurred in October–December of 2009; U_{11ij} = median age (years) of all patients seen during shift j ; U_{12ij} = percent of patients seen during shift j that were female; U_{13ij} = percent of patients seen during

shift j that arrived by EMS; U_{14ij} = percent of patients seen during shift j with urgent acuity class; U_{15ij} = percent of patients seen during shift j that had respiratory chief complaint; U_{16ij} = percent of patients seen during shift j that had musculoskeletal chief complaint; U_{17ij} = length of shift j (hours); U_{18ij} = duration of observation (hours); U_{19ij} = number of staff working in the ED during shift j ; U_{20ij} = number of patients in the ED during shift j .

For the network-level outcomes, separate models for each temporal and environmental predictor were run, with shift length and network size included as control variables. Since there were only 35 network-level observations, we did not perform extensive multivariable analysis beyond these models. Regression diagnostics confirmed linearity, residual assumptions, and lack of multicollinearity.

For these measures the models took the form

$$E(m_j) = \beta_0 + \sum_{k=1}^{10} \beta_k V_{kj}$$

for $j = 1, \dots, 35$, where m_j is one of the measures for shift j ; $V_{1j} = 1$ if shift j was a day shift, 0 otherwise; $V_{2j} = 1$ if shift j was a weekend shift, 0 otherwise; $V_{3j} = 1$ if shift j occurred in July–September of 2009, 0 if it occurred during October, November, December of 2009, 0 if it occurred during October, November, December of 2009; V_{4j} = maximum ED census during shift j ; V_{5j} = median age of all patients in the ED during shift j (years); V_{6j} = percentage of all patients present in ED who were female during shift j ; V_{7j} = percentage of all patients present in ED during shift j who arrived by ambulance; V_{8j} = percentage of all patients present in ED during shift j who were high acuity; V_{9j} = percentage of all patients present in ED during shift j who had respiratory syndrome; V_{10j} = percentage of all patients present in ED during shift j who had musculoskeletal chief complaint.

Gephi (Bastian et al., 2009) was used to calculate nodal- and network-level measures. SAS v9.3 (SI Inc., 2012) was used for all statistical analyses.

3. Results

3.1. Shift characteristics

Table 1 provides descriptive statistics for participant, temporal, and environmental characteristics. There were, on average, 36 staff and 68 patient participants observed per shift. Staff were observed for 575 min per shift while patients were observed for 192 min. Although the goal was to measure 12-h shifts, at least 25% of shifts were observed for 11 or fewer hours. The majority (74%) of shifts were on weekdays, and 57% were day shifts. Shifts split fairly evenly between summer months (July–September) and autumn months (October–December), with 46% of shifts being measured in summer months. The median of shift-level environmental characteristics are as follows: median age of 44 years, 57% female, 29% arrived by EMS, 28% were of high acuity, 12% presented with respiratory syndrome, and 12% presented with musculoskeletal syndrome.

3.2. Participant-level characteristics

3.2.1. Degree

Degree, relative degree, and weighted degree are given in Table 2. The regression analysis parameter estimates for patient–patient (PP), patient–staff (PS), staff–patient (SP), and staff–staff (SS) degree on environmental, temporal, and subject characteristics are shown in Tables 3–6 respectively. For each table, the models take the form of the equations given in section 2 above.

With respect to PP degree, the patients arriving by EMS had fewer contacts with other patients than patients arriving by other modes. Similarly patients in the least urgent acuity classes had more

Table 1
Participant, temporal, and environmental characteristics of the 35 ED contact networks.

	N	%	Median	(1st quartile, 3rd quartile)	Min–max
Participants					
Staff ^a			36	(33, 39)	25–47
Provider			7	(5, 8)	2–10
RN			15	(13, 16)	10–22
Other staff			15	(14, 16)	10–19
Patients participating ^a			68	(61, 80)	35–90
Total participants ^a			104	(92, 113)	66–133
Minutes observed by RFID^a					
Staff			575	(463, 638)	250–694
Patients			192	(156, 207)	114–235
Census maximum during observation^{a,b}					
Staff			51	(44, 57)	35–71
Patients			50	(47, 57)	35–69
Temporal characteristics					
Daytime shift	20	57%			
Weekday shift	26	74%			
Summer	16	46%			
Shift length (h)			12	(11, 12)	5–12
Environmental characteristics^c					
Median age, years			44	(41, 46)	37–51
% Female			57	(54, 59)	48–67
% Arriving by EMS ^d			29	(27, 33)	20–41
% High acuity			28	(25, 30)	19–41
% Respiratory syndrome			12	(10, 16)	3–22
% Musculoskeletal complaint			12	(9, 15)	7–20

^a Shift median was first calculated, and the distributions of these values are reported.
^b During an individual’s observation, the maximum number of patients registered in the ED during any 15 min period.
^c Shift summary statistic was first calculated and the distributions of these values are reported.
^d Emergency medical services.

Table 2
Degree measures (by edge type), closeness centrality, and eigenvector centrality of the 35 ED contact networks.

Node-level characteristic	Summary of shift median values ^a			
	Median	(1st quartile, 3rd quartile)	Shift median min–max	Individual min–max
Degree				
Patient with other patient	6	(5, 10)	3–12	0–40
Patient with a staff member	3	(2, 4)	1–8	0–28
Staff with a patient	4	(1, 10)	0–25	0–64
Staff with other staff	19	(16, 22)	11–28	0–39
Weighted degree (min)				
Patient with other patient	49	(31, 75)	16–94	0–1273
Patient with a staff member	8	(4, 22)	<1–71	0–2859
Staff with a patient	26	(4, 63)	0–198	0–1648
Staff with other staff	1106	(806, 1584)	529–2149	0–8294
Relative degree^b				
Patient with other patient	0.11	(0.08, 0.13)	0.04–0.18	0–0.48
Patient with a staff member	0.09	(0.05, 0.12)	0.03–0.20	0–0.89
Staff with a patient	0.06	(0.03, 0.14)	0–0.28	0–0.82
Staff with other staff	0.58	(0.50, 0.64)	0.29–0.87	0–1.00
Closeness centrality	0.49	(0.46, 0.51)	0.37–0.54	0–1
Eigenvector centrality	0.20	(0.15, 0.26)	0.08–0.42	0–1

^a The median for each shift was first calculated, and the statistics for these values are reported.
^b Relative degree = # contacts with other patients/(# patients-1).

contacts with other patients than patients in more acute classes. Patients with GI complaints had fewer contacts than patients with musculoskeletal complaints. PP degree was positively associated with crowding.

With respect to PP weighted degree, patients arriving by EMS were greater than patients arriving by other modes. There was also a positive association with crowding. There was also a significant interaction between arrival mode and crowding. When crowding was low (say census = 20), patients arriving by EMS had

more person-minutes of contact with other patients, compared to patients arriving by other modes. However, when crowding was high (say census = 70), patients arriving by EMS had fewer person-minutes of contact with other patients, compared to patients arriving by other modes. Moreover, patients in the stable or non-urgent acuity class had approximately 30 person-minutes of contact more than patients in the most acute category.

With respect to PS degree, patients in the least urgent acuity classes had more contacts with other patients than patients in more

Table 3
Among patient–patient contacts: relationship of patient, temporal, and environmental characteristics in the ED with degree, weighted degree, and relative degree.

	Patient contact with other patients					
	Degree		Weighted degree ^a		Relative degree	
	Estimate	(95% CL)	Estimate	(95% CL)	Estimate	(95% CL)
Intercept	1.46	(−1.74, 4.48)	−225	(−452, −19)	0.15	(−0.09, 0.36)
Patient characteristics						
Age (per 10 years)	−0.02	(−0.03, −0.01)	−1	(−3, 1)	0	(0, 0)
Female (vs. Male)	0.02	(−0.01, 0.06)	2	(−5, 10)	0	(0, 0.01)
Non-black (vs. Black)	−0.02	(−0.08, 0.05)	−6	(−17, 3)	0	(−0.01, 0.01)
EMS ^b arrival (vs. other)	−0.13	(−0.19, −0.07)	100	(43, 143)	0.04	(0.01, 0.07)
EMS arrival X census ^c	−	−	−25	(−34, −12)	−0.01	(−0.02, −0.01)
Acuity category (vs. emergent)						
Stable/non-urgent	0.20	(0.09, 0.31)	30	(9, 52)	0.02	(0.01, 0.04)
Urgent	0.06	(0, 0.12)	8	(−1, 18)	0.01	(0, 0.01)
Chief complaint (vs. musculoskeletal)						
GI ^d	−0.11	(−0.21, −0.04)	−3	(−13, 7)	−0.01	(−0.03, 0)
Respiratory	−0.07	(−0.14, 0.01)	−6	(−22, 8)	−0.01	(−0.02, 0)
Neurological	−0.08	(−0.21, 0.05)	−9	(−29, 8)	−0.01	(−0.02, 0.01)
Chest pain	−0.09	(−0.18, 0.01)	−1	(−17, 16)	−0.01	(−0.02, 0)
Other	−0.07	(−0.13, 0)	−4	(−14, 6)	−0.01	(−0.02, 0)
Patient stay ^e (vs. all observed)						
Ends during	−0.19	(−0.31, −0.06)	−18	(−39, 1)	−0.02	(−0.03, −0.01)
Starts during	0.02	(−0.14, 0.16)	13	(−12, 37)	0	(−0.01, 0.02)
Starts before/ends after	−0.65	(−0.88, −0.43)	−63	(−115, −15)	−0.08	(−0.11, −0.05)
Temporal characteristics						
Daytime shift (vs. evening)	−0.05	(−0.32, 0.18)	−16	(−34, 6)	−0.01	(−0.03, 0.01)
Weekend shift (vs. weekday)	0	(−0.32, 0.32)	−18	(−46, 0)	0	(−0.03, 0.02)
Fall (vs. summer)	0.21	(−0.15, 0.52)	14	(−9, 44)	0.02	(−0.01, 0.05)
Environmental characteristics						
Census maximum during observation ^f	0.31	(0.22, 0.39)	27	(18, 35)	0.03	(0.02, 0.03)
Median age, years ^f	−0.04	(−0.40, 0.35)	13	(−24, 50)	0	(−0.03, 0.03)
% Female ^g	−0.10	(−0.41, 0.23)	15	(−7, 36)	0	(−0.03, 0.02)
% Arriving by EMS ^g	−0.14	(−0.39, 0.24)	6	(−14, 36)	−0.01	(−0.03, 0.02)
% High acuity ^g	−0.24	(−0.34, 0.19)	−12	(−33, 8)	−0.01	(−0.02, 0.02)
% Respiratory syndrome ^g	−0.33	(−0.64, 0.05)	−8	(−29, 17)	−0.03	(−0.06, −0.01)
% Musculoskeletal complaint ^g	−0.18	(−0.61, 0.18)	−12	(−46, 14)	−0.02	(−0.05, 0)
Control variables						
Shift length (per hour)	0.01	(−0.08, 0.09)	−2	(−10, 5)	0	(−0.01, 0.01)
Duration observed (per hour)	0.13	(0.11, 0.15)	18	(14, 22)	0.02	(0.01, 0.02)
# Patients (per 10)	0.07	(−0.12, 0.15)	8	(−3, 16)	−0.01	(−0.02, 0)

Bold font indicates estimate is significantly different from 0 at the 0.05 level.

^a Minutes.

^b Emergency medical services.

^c Estimates reported per 10 patient increase of 'maximum number of patients registered in the ED during any 15 min period for individual participants'.

^d Gastrointestinal.

^e Relation of patients ED stay to the RFID observation period.

^f Estimates reported per 10 year increase in median age.

^g Estimates reported per 10% point increase.

acute classes. There was also an interaction between arrival mode and crowding. When crowding was low, patients arriving by EMS were similar to patients arriving by other modes, but as crowding increased, patient contacts with staff increased among patients arriving by other modes while patient contacts with staff did not increase significantly if arriving by EMS.

With respect to PS weighted degree, non-black patients have approximately 15 person-minutes less contact than black patients. Weekends are greater than weekdays by 32 person-minutes. Patients in the stable or non-urgent acuity class had approximately 34 person-minutes of contact more than patients in the most acute category. There was a significant interaction between arrival mode and crowding. When crowding was low, patients arriving by EMS had more person-minutes of contact with staff, compared to patients arriving by other modes. However, when crowding was high, patients arriving by EMS had fewer person-minutes of contact with staff, compared to patients arriving by other modes.

With respect to SP degree, both providers and nurses had more contacts with patients than other staff, with providers having the

most contacts. Contacts of staff with patients increased with crowding. However there was no interaction between staff role and crowding.

With respect to SP weighted degree, there is a significant interaction between staff role and crowding. When crowding is low, providers have the most contact time with patients and staff the least. However, when crowding is high, staff have the most contact time with patients. Staff had 80 fewer person-minutes of contact with patients on weekends than on weekdays.

With respect to SS degree, both providers and nurses have fewer contacts with all types of staff than staff in the other category, with providers having significantly fewer than nurses. There was a positive association with crowding.

With respect to SS weighted degree, providers and RNs are less than other staff. The average person-minutes in contact increased by 478 person-minutes for each 10% increase in population percent with high acuity.

Table 4
Among patient–staff contacts: relationship of patient, temporal, and environmental characteristics in the ED with degree, weighted degree, and relative degree.

	Patient contact with staff members					
	Degree		Weighted degree ^a		Relative degree	
	Estimate	(95% CL)	Estimate	(95% CL)	Estimate	(95% CL)
Intercept	−0.96	(−5.86, 5.24)	−101	(−494, 317)	−0.05	(−0.50, 0.51)
Patient characteristics						
Age (per 10 years)	0	(−0.03, 0.03)	−2	(−5, 1)	0	(0, 0)
Female (vs. male)	−0.01	(−0.07, 0.07)	0	(−12, 11)	0	(−0.01, 0.01)
Non-black (vs. black)	−0.04	(−0.14, 0.05)	−15	(−29, −2)	0	(−0.01, 0.01)
EMS ^b arrival (vs. other)	0.47	(−0.04, 1.00)	102	(37, 164)	0.08	(0, 0.15)
EMS arrival X census ^c	−0.15	(−0.26, −0.03)	−24	(−36, −10)	−0.02	(−0.04, −0.01)
Acuity category (vs. emergent)						
Stable/non-urgent	0.18	(0.06, 0.30)	34	(10, 59)	0.03	(0.01, 0.05)
Urgent	0.04	(−0.04, 0.12)	5	(−11, 19)	0.01	(−0.01, 0.01)
Chief complaint (vs. musculoskeletal)						
GI ^d	0.03	(−0.08, 0.15)	3	(−12, 20)	0	(−0.01, 0.02)
Respiratory	−0.01	(−0.13, 0.12)	8	(−9, 27)	0	(−0.02, 0.02)
Neurological	−0.01	(−0.2, 0.17)	2	(−19, 23)	0	(−0.03, 0.02)
Chest pain	−0.03	(−0.19, 0.11)	16	(−3, 37)	0	(−0.03, 0.02)
Other	−0.05	(−0.15, 0.06)	4	(−8, 16)	−0.01	(−0.03, 0.01)
Patient stay ^e (vs. all observed)						
Ends during	−0.46	(−0.64, −0.31)	−26	(−43, −11)	−0.06	(−0.08, −0.04)
Starts during	0.14	(−0.01, 0.27)	24	(5, 45)	0.03	(0.01, 0.06)
Starts before/ends after	−1.07	(−1.42, −0.76)	−65	(−115, −22)	−0.13	(−0.19, −0.09)
Temporal characteristics						
Daytime shift (vs. evening)	0.08	(−0.25, 0.44)	−5	(−41, 32)	0.01	(−0.03, 0.05)
Weekend shift (vs. weekday)	−0.26	(−0.68, 0.11)	−32	(−75, −1)	−0.04	(−0.09, 0.01)
Fall (vs. summer)	0.27	(−0.22, 0.70)	2	(−42, 44)	0.03	(−0.02, 0.08)
Environmental characteristics						
Census maximum during observation ^b	0.10	(−0.01, 0.22)	10	(−4, 23)	0.02	(0, 0.03)
Median age, years ^f	−0.15	(−0.60, 0.26)	−16	(−65, 25)	−0.02	(−0.07, 0.03)
% Female ^g	0.30	(−0.41, 0.87)	28	(−20, 78)	0.04	(−0.02, 0.11)
% Arriving by EMS ^g	0.07	(−0.30, 0.58)	9	(−24, 54)	0.02	(−0.02, 0.08)
% High acuity ^g	−0.20	(−0.54, 0.12)	−9	(−42, 23)	−0.02	(−0.07, 0.01)
% Respiratory syndrome ^g	0.07	(−0.38, 0.50)	5	(−26, 47)	0	(−0.04, 0.05)
% Musculoskeletal complaint ^g	0	(−0.68, 0.55)	−3	(−55, 43)	−0.01	(−0.08, 0.04)
Control variables						
Shift length (per hour)	0.09	(−0.05, 0.21)	4	(−7, 13)	0.01	(0, 0.03)
Duration observed (per hour)	0.10	(0.08, 0.13)	15	(12, 19)	0.01	(0.01, 0.02)
# Staff (per 10)	−0.02	(−0.54, 0.52)	−11	(−55, 37)	−0.04	(−0.10, 0.03)
# Patients (per 10)	−0.02	(−0.18, 0.14)	−2	(−17, 14)	−0.01	(−0.03, 0.01)

Bold font indicates that the estimate is significantly different from 0 at the 0.05 level.

^a Minutes.

^b Emergency medical services.

^c Estimates reported per 10 patient increase of 'maximum number of patients registered in the ED during any 15 min period for individual participants'.

^d Gastrointestinal.

^e Relation of patients ED stay to the RFID observation period.

^f Estimates reported per 10 year increase in median age.

^g Estimates reported per 10% point increase.

3.2.2. Other centrality measures

Regression parameter estimates for eigenvector centrality are shown in Table 7. There is a significant interaction between role and crowding. Eigenvector centrality increased slightly more with increasing crowding for other staff compared to patients, although there were not differences in slopes for providers vs. patients and nurses vs. patients.

The regression parameter estimates for closeness centrality are also shown in Table 7. There is an interaction between role and crowding. Providers are lower than patients at low crowding, but have higher values than patients with increased crowding.

3.3. Network-level characteristics

Simple descriptive statistics of the network-level characteristics are shown in Table 8. The median density of networks was 0.16, indicating that the individuals had interactions with each other at

the frequency of 16% of the number of possible 2-way interactions. The median radius was 3, with 34% of the networks having a radius less than 3, while the maximum radius was 4. The median diameter was 5, with 43% of networks with a diameter less than 5, while the maximum diameter was 6. The maximum spectral gap varied from 1 to 14, with a median value of 6. The median clustering coefficient was 0.56.

The results of regression analyses of the network-level characteristics on individual temporal and environmental characteristics while controlling for shift length and network size are given in Table 9. Among temporal characteristics time of day and season were associated with some network values, while weekday/weekend was not. Specifically, shifts occurring in daytime were more likely to have radius >2 (estimated beta (se) = 2.46 (1.00)) and had smaller spectral gap (−4.67 (1.07)) than night shifts. Shifts falling in autumn months (October–December) were less dense than shifts falling in summer months (July–September) (est. b

Table 5
Among staff–patient contacts: relationship of staff, temporal, and environmental characteristics in the ED with degree, weighted degree, and relative degree.

	Staff contact with patients					
	Degree		Weighted degree ^a		Relative degree	
	Estimate	(95% CL)	Estimate	(95% CL)	Estimate	(95% CL)
Intercept	–1.57	(–8.14, 7.06)	–348	(–1223, 571)	–0.05	(–0.67, 0.63)
Staff position						
Staff provider (vs. other)	0.76	(0.44, 0.99)	371	(157, 583)	0.24	(0.13, 0.34)
Nurse (vs. other)	0.31	(0.06, 0.52)	282	(125, 431)	0.09	(0.02, 0.17)
Staff (provider vs. other) X census ^b	–	–	–62	(–110, –15)	–0.03	(–0.05, –0.01)
Staff (nurse vs. other) X census	–	–	–62	(–96, –25)	–0.01	(–0.03, 0)
Temporal characteristics						
Daytime shift (vs. evening)	0.22	(–0.17, 0.71)	–23	(–87, 48)	0	(–0.04, 0.04)
Weekend shift (vs. weekday)	–0.34	(–0.93, 0.22)	–80	(–173, –17)	–0.04	(–0.11, 0.01)
Fall (vs. summer)	0.51	(–0.31, 1.31)	24	(–64, 124)	.03	(–0.03, 0.09)
Environmental characteristics						
Census maximum during observation ^b	0.44	(0.23, 0.72)	59	(25, 98)	0.02	(0, 0.04)
Median age, years ^c	0.15	(–0.46, 0.74)	–15	(–113, 76)	0.01	(–0.05, 0.06)
% Female ^d	0.07	(–0.86, 0.79)	49	(–52, 147)	0.03	(–0.05, 0.10)
% Arriving by EMS ^{d,e}	0.09	(–0.52, 1.1)	11	(–52, 120)	0.01	(–0.04, 0.08)
% High acuity ^d	–0.17	(–0.77, 0.30)	–12	(–90, 53)	–0.02	(–0.06, 0.03)
% Respiratory syndrome ^d	–0.01	(–0.77, 0.72)	10	(–53, 98)	0	(–0.06, 0.06)
% Musculoskeletal complaint ^d	–0.13	(–1.21, 0.56)	–11	(–123, 98)	–0.02	(–0.11, 0.04)
Control variables						
Shift length (per hour)	0.08	(–0.07, 0.28)	5	(–17, 31)	0	(–0.01, 0.02)
Duration observed (per hour)	0.14	(0.1, 0.17)	12	(8, 15)	.01	(0.01, 0.02)
# Staff (per 10)	–0.71	(–1.57, –0.13)	–65	(–165, 49)	–0.06	(–0.13, 0.02)
# Patients (per 10)	0.10	(–0.32, 0.26)	13	(–32, 45)	0	(–0.03, 0.02)

Bold font indicates that the estimate is significantly different from 0 at the 0.05 levels.

^a Minutes.

^b Estimates reported per 10 patient increase of 'maximum number of patients registered in the ED during any 15 min period for individual participants'.

^c Estimates reported per 10 year increase in median age.

^d Estimates reported per 10% point increase.

^e Emergency medical services.

Table 6
Among staff–staff contacts: relationship of staff, temporal, and environmental characteristics in the ED with degree, weighted degree, and relative degree.

	Staff contact with other staff					
	Degree		Weighted degree ^a		Relative degree	
	Estimate	(95% CL)	Estimate	(95% CL)	Estimate	(95% CL)
Intercept	1.13	(–1.37, 4.57)	–3653	(–8939, 2207)	.06	(–1.03, 1.25)
Staff position						
Staff provider (vs. other)	–0.58	(–0.70, –0.47)	–1461	(–1712, –1276)	–0.19	(–0.23, –0.16)
Nurse (vs. other)	–0.13	(–0.19, –0.07)	–970	(–1219, –752)	–0.10	(–0.14, –0.06)
Temporal characteristics						
Daytime shift (vs. evening)	0.20	(–0.01, 0.41)	–60	(–404, 330)	0.07	(–0.02, 0.15)
Weekend shift (vs. weekday)	0.05	(–0.18, 0.31)	109	(–394, 495)	–0.01	(–0.12, 0.08)
Fall (vs. summer)	0.15	(–0.06, 0.35)	147	(–169, 538)	0.08	(0.01, 0.16)
Environmental characteristics						
Census maximum during observation ^b	0.12	(0.03, 0.21)	–1	(–114, 146)	0.01	(–0.01, 0.03)
Median age, years ^c	–0.01	(–0.34, 0.26)	283	(–267, 736)	0.01	(–0.12, 0.13)
% Female ^d	–0.01	(–0.38, 0.38)	–21	(–657, 432)	0.04	(–0.09, 0.20)
% Arriving by EMS ^{d,e}	0.06	(–0.16, 0.39)	189	(–209, 607)	0.02	(–0.06, 0.13)
% High acuity ^d	0.04	(–0.16, 0.26)	470	(223, 873)	0.01	(–0.06, 0.09)
% Respiratory syndrome ^d	–0.03	(–0.33, 0.20)	–81	(–541, 348)	0.01	(–0.09, 0.09)
% Musculoskeletal complaint ^d	–0.10	(–0.52, 0.12)	220	(–389, 731)	–0.03	(–0.19, 0.05)
Control variables						
Shift length (per hour)	–0.02	(–0.09, 0.05)	–49	(–145, 67)	–0.01	(–0.04, 0.02)
Duration observed (per hour)	0.07	(0.05, 0.08)	185	(154, 216)	0.02	(0.02, 0.03)
# Staff (per 10)	0.19	(–0.15, 0.43)	478	(–10, 958)	–0.01	(–0.13, 0.09)

Bold font indicates estimates that are significantly different from 0 at the 0.05 level.

^a Minutes.

^b Estimates reported per 10 patient increase of 'maximum number of patients registered in the ED during any 15 min period for individual participants'.

^c Estimates reported per 10 year increase in median age.

^d Estimates reported per 10% point increase.

^e Emergency medical services.

Table 7
Relationship of participant, temporal, and environmental characteristics in the ED with eigenvector and closeness centrality.

	Eigenvector centrality		Closeness centrality	
	EST	95% CL	EST	95% CL
Intercept	0.01	(−0.75, 0.72)	0.39	(−0.22, 1.17)
Participant characteristics				
Participant type (vs. patient)				
Provider	0.16	(−0.01, 0.30)	−0.10	(−0.20, −0.02)
Nurse	0.19	(−0.04, 0.39)	−0.04	(−0.10, 0.02)
Other staff	0.17	(−0.07, 0.36)	−0.04	(−0.09, 0.01)
Census max during observation (per 10)				
Census max X (MD vs. patient)	−0.02	(−0.04, 0.02)	0.02	(0.003, 0.04)
Census max X (RN vs. patient)	0.01	(−0.03, 0.06)	0.01	(0, 0.02)
Census max X (other staff vs. patient)	0.04	(0.002, 0.08)	0.01	(0, 0.02)
Temporal characteristics				
AM shift (vs. PM)	0.02	(−0.03, 0.07)	0.04	(−0.01, 0.08)
Weekend (vs. weekday)	−0.01	(−0.07, 0.05)	−0.02	(−0.07, 0.03)
Summer (vs. Fall)	0.04	(−0.03, 0.12)	0.05	(−0.02, 0.10)
Environmental characteristics				
Median age (per 10 year)	0.02	(−0.07, 0.10)	−0.02	(−0.07, 0.03)
% Female (per 10%)	0	(−0.08, 0.08)	0	(−0.08, 0.07)
% Arriving by EMS (per 10%)	0	(−0.07, 0.09)	0.01	(−0.05, 0.08)
% High acuity (per 10%)	−0.02	(−0.10, 0.03)	−0.02	(−0.06, 0.02)
% With respiratory syndrome (per 10%)	−0.02	(−0.09, 0.06)	−0.01	(−0.07, 0.05)
% With musculoskeletal syndrome (per 10%)	−0.03	(−0.12, 0.05)	−0.01	(−0.10, 0.05)
Control variables				
Shift length	0.01	(−0.01, 0.03)	0.01	(−0.01, 0.03)
Duration observed (per hour)	0.02	(0.01, 0.02)	0.01	(0.01, 0.01)
Staff count (per 10)	−0.04	(−0.11, 0.04)	−0.03	(−0.10, 0.03)
Patient count (per 10)	−0.01	(−0.05, 0.01)	−0.01	(−0.03, 0.01)

Bold font indicates estimates that are significantly different from 0 at the 0.05 level.

Table 8
Network level characteristics of the 35 ED contact networks.

Network characteristic	Median	(1st quartile, 3rd quartile)	Min–Max	
Density	0.16	(0.14, 0.18)	0.10–0.25	
Average path length	2.12	(1.99, 2.27)	1.86–2.77	
Diameter	5	(4, 5)	3–6	(% <5) = 43%
Weighted diameter	0.30	(0.15, 0.69)	0.04–2.50	
Radius	3	(2, 3)	1–4	(% <3) = 34%
Maximum spectral gap	6	(3, 10)	1–14	
# Weak components	3	(2, 4)	1–7	
Clustering coefficient	0.56	(0.54, 0.61)	0.48–0.63	

Table 9
Relationship of temporal and environmental characteristics with network-level measures, controlling for shift length and network size^a; 35 ED contact networks.

Characteristic	Parameter estimates (se) ^b							
	Density	Average path length	Log (wt. diameter)	Maximum spectral gap	Clustering coefficient	# Weak components	Diameter (>5 vs. ≤5)	Radius (>2 vs. ≤2)
Temporal characteristics								
Daytime shift (vs. evening)	0 (.01)	0.05 (0.08)	0.34 (0.40)	−4.67 (1.07)	−0.03 (0.01)	−0.22 (0.56)	0.77 (0.81)	2.46 (1.00)
Weekendshift (vs. weekday)	0.01 (.01)	0.02 (0.08)	0.16 (0.41)	0.49 (1.40)	0.01 (0.02)	−0.49 (0.57)	0.98 (0.85)	0.20 (0.90)
Fall (vs. summer)	−0.03 (0.01)	0.18 (0.07)	0.36 (0.40)	3.05 (1.25)	−0.01 (0.01)	1.23 (0.52)	1.15 (0.83)	0.08 (0.88)
Environmental characteristics								
Census maximum during shift ^c	0.01 (0.01)	−0.02 (0.04)	0.07 (0.24)	0.53 (0.79)	0.01 (0.01)	−0.27 (0.33)	0.82 (0.53)	1.79 (0.79)
Median Age, years ^d	−0.02 (0.02)	0.11 (0.11)	1.06 (0.53)	0.96 (1.91)	−0.04 (0.02)	1.04 (0.77)	0.72 (1.15)	0.62 (1.26)
% Female ^e	0.03 (0.02)	−0.15 (0.09)	−1.09 (0.45)	−1.61 (1.63)	0.03 (0.02)	−1.07 (0.65)	−0.19 (0.96)	0.41 (1.08)
% Arriving by EMS ^e	0 (0.01)	0.04 (0.08)	0.19 (0.42)	0.94 (1.43)	0 (0.02)	0.37 (0.59)	−1.09 (0.88)	0.52 (0.92)
% High acuity ^e	−0.01 (0.01)	0.08 (0.07)	0.85 (0.33)	1.45 (1.21)	0.01 (0.01)	0.12 (.51)	1.60 (0.87)	1.61 (0.97)
% Respiratory syndrome ^e	0 (0.01)	−0.08 (0.07)	−0.66 (0.38)	1.88 (1.30)	0.01 (0.01)	0.28 (0.55)	0.13 (0.78)	−1.33 (0.91)
% Musculoskeletal complaint ^e	−0.03 (0.02)	0.08 (0.10)	0.26 (0.52)	0.51 (1.77)	−0.05 (0.02)	1.58 (0.70)	−0.80 (1.05)	0.50 (1.14)

Bold font indicates estimates that are significantly different from 0 at the 0.05 level.

^a Density, average path length, log(weighted diameter), maximum spectral gap, and clustering coefficient using multiple linear regression; # weak components using poisson regression, radius and diameter (median splits) using logistic regression; all models contain 3 variables (the variable of interest and control variables regarding shift length and total number of participants (staff plus patients) in each shift).

^b Estimates significant at $p < 0.05$ are in bold.

^c Census reported per 10 patient increase of *shift mean* of the 'maximum number of patients registered in the ED during any 15 min period for individual participants'.

^d Estimates reported per 10 year increase in median age.

^e Estimates reported per 10% point increase.

(se): -0.03 (0.01)) while having longer average path lengths (0.18 (0.07)), larger spectral gap (3.05 (1.25)), and larger number of components (1.23 (0.52)).

Among environmental characteristics, census maximum, percent female patients, percent high acuity patients, and percent musculoskeletal complaint patients were associated with some network measures, while median age, percent arriving by EMS, and percent respiratory complaint patients were not. Specifically, the odds (standard error) of having radius >2 increased with shift census maximum (1.79 (0.79)). The log of the weighted diameter decreased as percentage of female patients present increased (-1.09 (0.45)); however this same outcome measure increased as percentage of high acuity patients present increased (0.85 (0.33)). Number of components increased as percentage of patients with musculoskeletal complaints increased (1.58 (0.07)), but the clustering coefficient decreased with this same covariate (-0.05 (0.02)).

4. Discussion

4.1. Key results

Major findings for patient contacts are as follows: (1) the number of patient contacts (degree) are not related to patient demographic characteristics, but the time-weighted number of contacts (weighted degree) for black patients were higher than for non-black patients; (2) there are differences in number of contacts based on arrival mode and staff role; and (3) the effect of arrival mode is modified by the extent of crowding. With respect to arrival mode, EMS patients have a more consistent geographic pathway of care while patients arriving by other modes are more ambulatory and therefore more easily moved between various waiting, evaluation and treatment areas (for example waiting room, triage, registration, imaging, treatment rooms). Patients arriving by EMS enter through the ambulance bay entry and are moved directly into a treatment room where traditional arrival activities occur (e.g., registration, triage, specimen collection, EKG, etc.). In contrast, for “ambulatory” patients, traditional arrival activities occur in a variety of locations to which the patient moves as opposed to a single treatment room. Thus the increasing number of contacts that is associated with increased crowding does not affect EMS patients – as it does patients arriving by other modes – since they are effectively sequestered in a single room, leading to the observed statistical interaction.

As for SP contacts, degree, weighted degree, and relative degree are all increased with crowding; thus in crowded conditions, staff have more contacts of longer duration with more patients. Because the provider to patient ratio is smaller than the non-provider staff to patient ratio, providers may become task saturated at an earlier point as conditions become more crowded and therefore become the rate limiting step in care delivery. For SS relative degree, providers $<$ nurses $<$ other, which suggests that the non-clinical staff work within smaller spaces filled with other staff colleagues. We will be exploring this possibility further in a planned future analysis of the geography of contacts.

4.2. Limitations

With only 35 networks, the network level models could not control for more covariates. In addition, we could not investigate the whole range of seasonality with observations for only 6 months. This ED does not treat pediatric patients, and thus contact patterns may differ in those settings. We did not measure interactions among and between visitors, i.e., family accompanying patients. The entire span of a patient’s visit is not captured by the RFID observation period although we tried to control for this in our models.

Contact patterns change over the course of a patient visit. Longer continuous observation periods would mean that a higher percentage of patients would have a completely captured ED stay.

4.3. Generalizability

Gundlapalli et al. (2009) used electronic health records coupled with data from a staff RTLS using infrared detection to reconstruct the social network of staff–patient interactions in a hospital ED over the course of one day. During this day there were 21 staff and 40 patients. The average PS degree was 3, while the average SP degree was 6. The average path length was 2.577 and the clustering coefficient was 0.541. The average (SD) eigenvector centrality was 14.7 (22.6) for staff and 6.9 (9.3) for patients. We cannot compare our findings to theirs, however, in that staff–staff and patient–patient interactions are not included. Polgreen et al. (2010) used student observers to record interactions of 148 hospital staff with other staff, patients, and others over the course of 606 h. They recorded 4413 interactions with other staff, and thus the staff–staff network has average degree = 59.6 and density = 0.41. They also indicate that these staff had 1762 interactions with patients. However, without the number of patients and the number of patient–patient interactions we cannot describe the entire network and thus cannot compare our network to theirs. Isella et al. (2011a) used an RFID system to record interactions of patients, staff, and visitors in a ward of a pediatric children’s hospital. There were 119 participants having average degree of 20 and network density of 0.17. Since we did not record interactions of visitors our network is not comparable.

4.4. Implications

Traditional models assume random mixing of members of healthcare networks. Prior work (Gundlapalli et al., 2009; Lowery-North et al., 2013) has demonstrated that in the emergency department network members are circulating in a non-random fashion. The data presented here further emphasize the non-random mixing characteristics of the ED population. These results should help focus designers of emergency care on higher risk situations for the transmission of dangerous pathogens in an emergency department. In particular given the differential in contacts with EMS vs. non-EMS patients, it would appear that infection control efforts should consider non-EMS patients, particularly during times of ED crowding, as high priority targets for engineering or architectural interventions to artificially increase social distance with each other and with staff. Similar measures should also be directed at staff to increase their social distance from one another.

In summary, we investigated the association of contacts with individual patient characteristics, staff role, and shift level characteristics such as patient mix. We have observed repeated shifts while other studies have studied short fixed intervals at one fixed time. Our findings suggest that optimization of the geographic pathway of care is necessary, particularly for patients not arriving by EMS, in order to reduce the number of contacts with other patients, thus reducing cross infection risk. Such optimization might limit intradepartmental motion of patients and thus reduce the number of patient contact events. Therefore we need to examine movement patterns of EMS arrivals vs. other arrivals to verify our conjecture. We also need to better understand not only the geographic pathways of care but also the locations where PP interactions are occurring under crowded conditions (e.g., waiting room, transit areas like hallways) in order to find ways to reduce the number and duration of contacts taking place there. Social network analysis enabled characterization of the ED in a novel manner. Our findings would not have been possible without the ability to look at ED patient and staff interactions through this lens.

Competing interests

The authors declare that they have no competing interests.

Acknowledgement

This research was supported by NIH contract HHSN266200700006C.

References

- Bastian, M., Heymann, S., Jacomy, M., 2009. Gephi: an open source software for exploring and manipulating networks. In: *International AAAI Conference on Weblogs and Social Media*, San Jose, CA.
- Bridges, C.B., Kuehnert, M.J., Hall, C.B., 2003. Transmission of influenza: implications for control in health care settings. *Clin. Infect. Dis.* 37, 1094–1101.
- Carroll, R., Ruppert, D., 1988. *Transformation and Weighting in Regression*. Chapman & Hall, London.
- Centers for Disease Control and Prevention, 2010. Interim results: influenza A (H1N1) 2009 monovalent vaccination coverage—United States, October–December 2009. *MMWR Morbidity and mortality weekly report* 59: 44.
- Fournet, J., Barrat, A., 2014. Contact patterns among high school students. *PLoS One* 9.
- Ginsberg, M., Hopkins, J., Maroufi, A., Dunne, G., Sunega, D., et al., 2009. Swine influenza A (H1N1) infection in two children—Southern California, March–April 2009. *Morbidity and Mortality Weekly Report* 58: 400–402.
- Gundlapalli, A., Ma, X., Benuzillo, J., Pettey, W., Greenberg, R., et al., 2009. Social network analyses of patient–healthcare worker interactions: implications for disease transmission. *AMIA Annual Symposium Proceedings*, 2009, 213–217.
- Hornbeck, T., Naylor, D., Segre, A.M., Thomas, G., Herman, T., et al., 2012. Using sensor networks to study the effect of peripartetic healthcare workers on the spread of hospital-associated infections. *J. Infect. Dis.* 206, 1549–1557.
- Isella, L., Romano, M., Barrat, A., Cattuto, C., Colizza, V., et al., 2011a. Close encounters in a pediatric ward: measuring face-to-face proximity and mixing patterns with wearable sensors. *PLoS One* 6, e17144.
- Isella, L., Stehle, J., Barrat, A., Cattuto, C., Pinton, J.F., et al., 2011b. What's in a crowd? Analysis of face-to-face behavioral networks. *J. Theor. Biol.* 271, 166–180.
- Lewis, M.D., Pavlin, J.A., Mansfield, J.L., O'Brien, S., Boomsma, L.G., et al., 2002. Disease outbreak detection system using syndromic data in the greater Washington DC area. *Am. J. Prev. Med.* 23, 180–186.
- Lowery-North, D.W., Hertzberg, V.S., Elon, L., Cotsonis, G., Hilton, S.A., et al., 2013. Measuring social contacts in the emergency department. *PLoS One* 8, e70854.
- Lucet, J.C., Laouenan, C., Chelius, G., Veziris, N., Lepelletier, D., et al., 2012. Electronic sensors for assessing interactions between healthcare workers and patients under airborne precautions. *PLoS One* 7.
- Newman, M.E.J., 2010. *Networks: An Introduction*. Oxford University Press, New York.
- Polgreen, P.M., Tassier, T.L., Pemmaraju, S.V., Segre, A.M., 2010. Prioritizing health-care worker vaccinations on the basis of social network analysis. *Infect. Control Hosp. Epidemiol.* 31, 893–900.
- SI Inc., 2008a. The MIXED Procedure. *SAS/STAT 92 User's Guide*. Cary, NC, pp. 3886–4086.
- SI Inc., 2008b. The GLIMMIX Procedure. *SAS/STAT 92 User's Guide*. Cary, NC, pp. 2078–2428.
- SI Inc., 2012. *What's New in SAS 9.3*. SAS Institute Inc., Cary, NC.
- Salathe, M., Kazandjieva, M., Lee, J.W., Levis, P., Feldman, M.W., et al., 2010. A high-resolution human contact network for infectious disease transmission. *Proc. Natl. Acad. Sci. U. S. A.* 107, 22020–22025.
- Stehle, J., Voirin, N., Barrat, A., Cattuto, C., Isella, L., et al., 2011. High-resolution measurements of face-to-face contact patterns in a primary school. *PLoS One* 6.
- Stevenson, A., 2010. *Oxford Dictionary of English*. Oxford University Press.
- Tellier, R., 2006. Review of aerosol transmission of influenza A virus. *Emerg. Infect. Dis.* 12, 1657–1662.
- Vanhems, P., Barrat, A., Cattuto, C., Pinton, J.-F., Khanafer, N., et al., 2013. Estimating potential infection transmission routes in hospital wards using wearable proximity sensors. *PLoS One* 8, e73970.
- Varia, M., Wilson, S., Sarwal, S., McGeer, A., Gournis, E., et al., 2003. Investigation of a nosocomial outbreak of severe acute respiratory syndrome (SARS) in Toronto, Canada. *Can. Med. Assoc. J.* 169, 285–292.
- WH Organization, 2009. Influenza-like illness in the United States and Mexico.
- Wenzel, R.P., Edmond, M.B., 2003. Managing SARS amidst uncertainty. *N. Engl. J. Med.* 348, 1947–1948.