



EMORY
LIBRARIES &
INFORMATION
TECHNOLOGY

OpenEmory

Processes of Change in Trauma-Focused Cognitive Behavioral Therapy for Youths: An Approach Informed by Emotional Processing Theory

Elizabeth Alpert, *University of Delaware*

Adele M Hayes, *University of Delaware*

[Charlotte Yasinski](#), *Emory University*

Charles Webb, *State of Delaware Division of Prevention and Behavioral Health Services*

Esther Deblinger, *Rowan University*

Journal Title: CLINICAL PSYCHOLOGICAL SCIENCE

Volume: Volume 9, Number 2

Publisher: SAGE PUBLICATIONS INC | 2021-02-16, Pages 270-283

Type of Work: Article | Post-print: After Peer Review

Publisher DOI: 10.1177/2167702620957315

Permanent URL: <https://pid.emory.edu/ark:/25593/vt4s3>

Final published version: <http://dx.doi.org/10.1177/2167702620957315>

Accessed September 25, 2023 12:27 AM EDT



Published in final edited form as:

Clin Psychol Sci. 2021 March 1; 9(2): 270–283. doi:10.1177/2167702620957315.

Processes of Change in Trauma-Focused Cognitive Behavioral Therapy for Youth: An Emotional Processing Theory Informed Approach

Elizabeth Alpert¹, Adele M. Hayes¹, Carly Yasinski², Charles Webb³, Esther Deblinger⁴

¹University of Delaware, Department of Psychological and Brain Sciences

²Emory University, Department of Psychiatry and Behavioral Sciences

³State of Delaware Division of Prevention and Behavioral Health Services

⁴Rowan University, School of Osteopathic Medicine

Abstract

This study examines processes of change in trauma-focused cognitive behavioral therapy (TF-CBT) delivered to a community sample of 81 youth. Emotional processing theory (EPT) is used as an organizational framework. EPT highlights activating and changing pathological trauma-related responses and increasing adaptive responses across cognitive, emotional, behavioral, and physiological domains. We coded sessions during the trauma processing phase of TF-CBT to examine the extent to which pathological and adaptive trauma-related responses were activated across domains. Higher scores indicate that more domains (0–4) were activated at a threshold of moderate to high intensity. Curvilinear change (inverted U, increase then decrease) in multimodal negative response scores across sessions predicted improvement in internalizing and PTSD symptoms at posttreatment. Linear increases in multimodal positive responses predicted improvement in externalizing symptoms. Findings suggest value in activating and changing both pathological and adaptive trauma responses across multiple domains and examining nonlinear patterns of change.

Keywords

childhood trauma; posttraumatic stress disorder; emotional processing; trauma-focused cognitive behavioral therapy; psychotherapy process

Childhood trauma is associated with numerous consequences that can persist throughout development, including poor academic performance, substance use, violent behavior, depression, and suicide (Paolucci, Genuis, & Violato, 2001; Roth, Newman, Pelcovitz, van

Primary corresponding author: Elizabeth Alpert, Department of Psychological and Brain Sciences, University of Delaware, 108 Wolf Hall, Newark, DE, 19716, USA., ealpert@udel.edu; Secondary corresponding author: Adele M. Hayes, Department of Psychological and Brain Sciences, University of Delaware, 108 Wolf Hall, Newark, DE, 19716, USA., ahayes@udel.edu.

Author Roles: EA developed the rationale and method for applying emotional processing theory to this investigation of change processes in TF-CBT and conducted the data analyses and interpretation under the supervision of AMH. EA and AMH drafted the manuscript. AMH designed and conducted the clinical trial on which this study was based and developed the CHANGE coding system used in the current study. CY trained coders and supervised coder meetings. CW and ED trained and supervised study therapists throughout the clinical trial. All authors provided critical revisions and approved the final version for submission.

der Kolk, & Mandel, 1997; Tyler, 2002). Trauma-focused cognitive behavioral therapy (TF-CBT; Cohen, Mannarino, & Deblinger, 2006; 2017) is an evidence-based treatment for youth who have experienced a wide range of childhood traumas, and there is strong empirical support for its effectiveness in reducing PTSD and other trauma-related symptoms (Gutermann et al., 2016; Lenz & Hollenbaugh, 2015). However, as is the case with many evidence-based treatments for children and adolescents (Weisz & Kazdin, 2010), little is known about processes of therapeutic change in TF-CBT. A better understanding of which in-session variables predict treatment outcomes could help clinicians to focus on those key targets to improve treatment efficacy.

Emotional Processing Theory as a Framework for Studying Processes of Change in TF-CBT

A number of trauma treatments for children and adults focus on the processing of traumatic experiences. These treatments include TF-CBT (Cohen et al., 2006; 2017; 2018), prolonged exposure (PE; Foa, Hembree, & Rothbaum, 2007), cognitive processing therapy (CPT; Resick, Monson, & Chard, 2017; Resick & Schnicke, 1992), cognitive therapy (e.g., Perrin et al., 2017), emotion-focused therapy (e.g., Paivio & Nieuwenhuis, 2001; Paivio et al., 2001), written exposure therapy (Sloan & Marx, 2019), and narrative exposure therapy (Schauer, Neuner, & Elbert, 2011). Emotional processing theory (EPT; Foa, Huppert, & Cahill, 2006; Foa & Kozak, 1986) can therefore be a useful common framework to study how therapeutic change occurs in TF-CBT and other trauma treatments for youth and adults.

EPT posits that psychopathology related to anxiety and traumatic stress is represented by a pathological network of stimuli, response elements (cognitive, emotional, behavioral, physiological), and their meaning. Emotional processing occurs when the pathological network is activated and incompatible information is introduced to promote corrective learning of more adaptive associations and responses. Although EPT has been studied most often in the context of exposure-based treatments, emotional processing was originally proposed as a transdiagnostic mechanism of change across different types of treatments (Foa & Kozak, 1986) for adults and youth (Peterman, Carper, & Kendall, 2019). Fear activation and habituation were initially emphasized as indicators of emotional processing, but updates to EPT (Foa et al., 2006) highlight the importance of changing responses to fear stimuli across multiple domains of functioning, including cognitive, emotional, behavioral, and physiological domains. The updated theory (Foa et al., 2006; Foa & McNally, 1996) also incorporates findings on inhibitory learning (e.g., Bouton, 2000; see Craske et al., 2008 for a review), which suggest that new responses learned in treatment do not replace the pathological learning, but rather can inhibit or buffer the old responses. This suggests that it is also important to measure new, more adaptive responses across multiple domains.

TF-CBT is based in part on emotional processing theory (Cohen et al., 2018), especially the trauma narration and processing (second) phase of treatment. Clinicians help youth develop a trauma narrative, which activates the network of cognitive, emotional, behavioral, and physiological responses to trauma memories. Clinicians then guide youth to process their traumatic experiences, and they also provide corrective information to help them learn more

adaptive responses (Cohen et al., 2018). EPT is therefore relevant to the study of change in TF-CBT and can help to link findings from this study to the broader literature on change in other child and adult treatments for PTSD.

Emotional Processing and Multimodal Change

According to EPT and related treatments, processing of traumatic memories is most potent if multiple types of trauma responses are targeted (Cohen et al., 2006; 2017; Foa et al., 2007; Foa et al., 2006; Foa & McLean, 2016). This multimodal focus includes trauma-related cognitions, emotions, behaviors, and physiological responses. However, most studies on the process of change in PTSD treatments have examined only one domain of functioning rather than multimodal activation and change.

For instance, decreases in negative cognitions predict symptom reduction in TF-CBT (Pfeiffer, Sachser, de Haan, Tutus, & Goldbeck, 2017) and other trauma-focused therapies for children and adolescents (e.g., König et al., 2019; McLean, Yeh, Rosenfield, & Foa, 2015; Smith et al., 2007) and for adults (e.g., McLean, Su, & Foa, 2015; McLean et al., 2019; Scher, Suvak, & Resick, 2017; Zalta et al., 2014). Reductions in self-reported fear reactivity across sessions predict better treatment outcomes (Asnaani, McLean, & Foa, 2016; Cooper, Clifton, & Feeny, 2017), and other negative emotions such as anger, shame, guilt, and disgust also play an important role in PTSD (McLean & Foa, 2017) and its treatment (Foa, Riggs, Massie, & Yarczower, 1995; Kaczurkin, Asnaani, Zhong, & Foa, 2016; Rizvi, Vogt, & Resick, 2009). In the behavioral domain, avoidance (e.g., avoidance of trauma cues, homework noncompliance) predicts worse posttreatment outcomes (Cooper, Kline, et al., 2017; Gloth, 2017). In addition, there is evidence that changes in a number of physiological indicators of distress predict better treatment outcomes in exposure therapy for PTSD, such as reductions in physiological fear response (e.g., heart rate) across sessions (Asnaani et al., 2016; Cooper, Clifton, et al., 2017), decreases in trauma-potentiated startle from pretreatment to posttreatment (Maples-Keller et al., 2019), and an increase then decrease in trauma-potentiated startle (Robison-Andrew et al., 2014). Cisler and colleagues (2015) also found that at pretreatment, adolescent girls who improved more in TF-CBT showed greater differential amygdala reactivity to fear faces than to neutral faces (i.e., better threat vs. safety discrimination) than nonresponders. Although these studies focus on one specific domain of functioning, taken together, they illustrate the importance of a more multimodal focus on negative trauma responses across multiple domains.

As with pathological responses to trauma content, some research has examined more adaptive responses learned in treatment. Again, these studies typically examine one domain in isolation rather than a multimodal set of responses. For example, accommodation (a cognitive construct that captures healthy, balanced beliefs about oneself, others, and the future) has been shown to predict better outcomes in TF-CBT (Ready et al., 2015) and in cognitive processing therapy for youth (König et al., 2019) and for adults (Dondanville et al., 2016; Iverson et al., 2015). There is some evidence that positive emotions reduce the reacquisition of fear, which could have implications for trauma treatments (Dour, Brown, & Craske, 2016; Zbozinek & Craske, 2017). Increases in approach behaviors toward previously avoided trauma-related situations and reminders are also associated with PTSD symptom

improvement (Porter, Romero, & Barone, 2018). Mindfulness-based interventions, which are associated with positive physiological responses (e.g., lower heart rate, increased parasympathetic activity; Ditto, Eclache, & Goldman, 2006; Mazzei, 2017), have been associated with reduction in PTSD symptoms (Colgan, Christopher, Michael, & Wahbeh, 2016; Held, Owens, Monroe, & Chard, 2017). Together, these findings suggest value in measuring the development of more adaptive responses across domains of functioning, in addition to the reduction of pathological responses.

PTSD is a complex system of pathology (Sripada, Rauch, & Liberzon, 2016), and studies on processes of change in PTSD treatments should capture the breadth of pathological and adaptive trauma responses that a client can experience. We used an observational coding system of therapeutic change processes, the CHANGE (Hayes, Feldman, & Goldfried, 2007), to code therapy sessions from TF-CBT. The CHANGE uses verbal and nonverbal information and can assess trauma reactions expressed by clients across the cognitive, emotional, behavioral, and physiological domains of interest in this study. The measure also assesses both pathological and adaptive responses.

Emotional Processing and Nonlinear Change

Another implication of EPT is that research on processes of change in treatment might be advanced by examining not only linear, but also nonlinear patterns of change. According to EPT, emotional processing involves the multimodal activation of the pathological network of trauma-related responses, followed by the weakening of these responses across domains (Foa et al., 2006; Foa & Kozak, 1986). Thus, this pattern of change might be captured by a quadratic (concave down, inverted U) pattern of change, beyond a linear trajectory of change. Indeed, change in psychotherapy does not always follow a linear trajectory (Hayes, Laurenceau, Feldman, Strauss, & Cardaciotto, 2007). For instance, data suggest that symptom trajectories in some PTSD treatments are better represented by quadratic than by linear functions (Nishith, Resick, & Griffin, 2002; Schumm, Walter, & Chard, 2013). Robison-Andrew and colleagues (2014) also found that quadratic change in startle response measured at pre-, mid-, and posttreatment predicted better treatment response in prolonged exposure and present-centered therapy for PTSD. Quadratic change in negative trauma responses across domains may similarly predict treatment outcome in TF-CBT, although this has not yet been examined.

The Current Study

Using EPT as an organizational framework (Foa et al., 2006), we examined change in both pathological and adaptive responses to trauma-related content during the trauma narration and processing phase of TF-CBT for trauma-exposed youth (Cohen et al., 2006; 2017). This phase of treatment was selected because it focuses specifically on the activation and processing of the traumatic memories and their impacts. The CHANGE coding system (Hayes, Feldman, et al., 2007) was used to assess the extent to which pathological and adaptive responses to trauma-related content were multimodal (i.e., activated at a moderate to high level across cognitive, emotional, behavioral, and physiological domains). Higher multimodal response scores indicated that more domains (range 0–4) were activated at

threshold. Trauma responses were coded for each session during this treatment phase to allow for an evaluation of linear and curvilinear trajectories as predictors of internalizing, externalizing, and PTSD symptoms at posttreatment.

In line with emotional processing theory (Foa et al., 2006; Foa & Kozak, 1986) and reviews of its evidence (Asnaani et al., 2016; Foa et al., 2006), we expected that in addition to a linear decrease in multimodal negative trauma-related responses, a curvilinear pattern would also predict more improvement in posttreatment measures of psychopathology (internalizing, externalizing, and PTSD symptoms). In other words, multimodal negative response scores that increased and then decreased (weakened) over the processing phase of TF-CBT were expected to predict better outcomes. Given the emphasis in EPT on strengthening new, adaptive associations (Foa et al., 2006) and the importance of multimodal positive responses in new learning (e.g. Dour et al., 2016; Porter et al., 2018), we expected greater linear increases in multimodal positive responses to predict better posttreatment outcomes. There is no theoretical reason to expect that positive responses would change in a curvilinear way, so only linear change was examined.

Method

Participants

Participants were recruited as part of a larger effectiveness trial of TF-CBT for trauma-exposed youth (see Webb, Hayes, Grasso, Laurenceau, & Deblinger, 2014 for trial outcomes, informed consent procedures, and information on participant recruitment and retention; [ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT01649141) Identifier: NCT01649141). The procedures for this trial were approved by the institutional review boards of all participating agencies. Youth were eligible for the trial if they had a score of 17 or more on the UCLA PTSD Reaction Index for DSM-IV-Abbreviated (UPID-A; Steinberg, Brymer, Decker, & Pynoos, 2004) or endorsed 3 of 9 PTSD symptoms based on an independently verified trauma (e.g., through child welfare). Eligible youth also spoke English, qualified for publicly-funded treatment, and had a non-offending caregiver willing to participate in treatment. Youth were excluded if they had an intellectual disability, untreated psychosis or current substance abuse, required frequent hospitalizations or a higher level of care, or had a sibling already in the study.

Of the 109 participants who met inclusion criteria for the effectiveness trial, 81 began the trauma processing (second) phase of treatment and were included in the present sample. These youth did not differ demographically from the 28 participants who discontinued treatment before this phase of treatment. The current sample included 54 (66.7%) females and 27 (33.3%) males, ages 7–17 years old ($M = 12.6$, $SD = 2.8$). They were 51.9% White, 40.7% Black or African American, 3.7% Hispanic or Latino, and 3.7% Biracial. Youth had experienced on average 3.5 types of trauma ($SD = 1.7$). Thirty-seven percent were in foster care when they enrolled. Youth completed an average of 5.28 sessions of the approximately 6 sessions in the trauma processing phase of TF-CBT ($SD = 2.17$). Using hierarchical piecewise regression analyses with this same sample, Ready and colleagues (2015) reported that there were significant reductions in internalizing ($\gamma = -.98$, $SE = .15$, $p < .001$), externalizing ($\gamma = -.76$, $SE = .21$, $p < .001$), and PTSD symptoms ($\gamma = -2.55$, $SE = .29$, $p < .001$) from pretreatment to posttreatment.

Procedure

Therapy.—Trauma-focused cognitive behavioral therapy (TF-CBT; Cohen et al., 2006; 2017) is an evidence-based treatment for children and adolescents who have experienced a range of childhood traumas, as well as their non-offending caregivers. TF-CBT incorporates psychoeducation, skills-building, gradual exposure to traumatic memories, and processing of trauma-related content. Youth and caregivers meet with a therapist in separate 30- to 45-minute sessions for approximately 12–15 sessions and also in several conjoint sessions, if appropriate.

TF-CBT is divided into three phases. Phase 1, the *stabilization and skills building* phase, emphasizes psychoeducation and coping skills, such as relaxation and emotion regulation skills. In phase 2, the *trauma narration and processing* phase, youth are exposed to the trauma memory and trauma-related content by developing a detailed narrative of their experiences. As youth create and discuss their narratives, therapists help them to emotionally process their experiences and challenge maladaptive beliefs about the trauma, its meaning, and its consequences. The third phase, the *integration and consolidation* phase, includes in vivo mastery activities when needed, trauma-focused conjoint sessions in which youth share the narrative with caregivers if clinically appropriate, and the development of personal safety skills. The present study focuses on sessions from the second phase of treatment, when trauma processing is the focus of sessions.

Therapy was delivered by a team of 25 clinicians who either held a professional degree or were doctoral students in clinical psychology programs. Unlicensed clinicians were supervised by a licensed practitioner. Adherence ratings suggested that therapists delivered TF-CBT with good fidelity (see Ready et al., 2015 for further details).

Measures

Session coding.—The CHANGE coding system (Hayes, Feldman, et al., 2007) was used to code each treatment session in the trauma processing phase of TF-CBT for each participant. Sessions were coded for the extent of multimodal activation of pathological and adaptive trauma-related responses across cognitive, emotional, behavioral, and physiological domains of functioning. The CHANGE is designed to capture a range of variables thought to be central to therapeutic change. This coding system has been used to code written narratives in cognitive behavioral treatments for depression (Hayes, Beevers, Feldman, Laurenceau, & Perlman, 2005; Hayes, Feldman, et al., 2007) and adult PTSD (Alpert, Hayes, Barnes, & Sloan, 2020), as well as audiotaped sessions of prolonged exposure for adult PTSD (Jun, Keller, Zoellner, Feeny, & Hayes, 2016), cognitive therapy for personality disorders (Hayes & Yasinski, 2015), and cognitive therapy for treatment-resistant depression (Abel, Hayes, Henley, & Kuyken, 2016).

In the current study, a team of 19 graduate and undergraduate students coded audio recordings of sessions ($M = 5.28$ sessions per participant, $SD = 2.17$). Coders were trained in the CHANGE coding system and practiced coding with experienced coders until they reached sufficient agreement on target variables (intraclass correlations (ICCs) $> .80$). Two coders rated each session, and each coder was paired with multiple other coders. Weekly

consensus meetings were held to prevent rater drift and to reach group consensus on discrepancies of two or more points on the 4-point scale of the CHANGE. Discrepancies were replaced with consensus ratings, and the ratings of the two coders were then averaged.

Each session was coded for trauma-related responses that occurred in that session or over the past week. To be coded, content had to relate to the trauma or its impact. Negative and positive responses, coded separately, included responses in four domains: *cognitions* about the self, others, or the future (e.g., “I can’t trust anyone” or “I will be able to heal”), *emotions* (e.g., sad or pleased), *behaviors* (e.g., getting in a fight or using coping skills when angry), and *physiological* experiences (e.g., difficulty breathing or feeling relaxed). For the cognitive variable, scores were calculated by taking the highest of three codes representing beliefs about the self, others, and the future. The CHANGE variables are coded for intensity on a four-point scale from 0 (*absent or very low*) to 3 (*high*).

Detailed descriptions of each coding category and examples of session content that would be coded as “high” are presented in Table 1. For each variable, a final intra-class correlation (ICC) was calculated by computing the ICC for each coder pair and then averaging across coders, as recommended by Hallgren (2012). ICCs ranged from .66–.91, which is in the good to excellent range of agreement (Cicchetti, 1994; see Table 1).

Multimodal response scores.—Because many trauma treatments emphasize the importance of activating and changing responses across multiple domains (e.g., Cohen et al., 2006; 2017; Foa et al., 2007), we created a composite score to quantify the extent to which negative and positive responses were activated across four domains: cognitive, emotional, behavioral, and physiological. The intensity scores (0–3) for each of the CHANGE variables in a given session were used to calculate the number of the four domains that were activated at a threshold of moderate to high intensity (score 2 on the 0–3 CHANGE scale). Multimodal response scores can range from 0 to 4 domains activated at threshold; these scores take into account both the intensity and breadth of trauma-related responses.

For example, a child describing abuse by his uncle might express the belief, “I can never trust anyone ever again” (negative cognitions intensity = 3), report feeling very anxious all the time (negative emotions = 3) and a bit shaky one time (negative physiological = 1), and refuse to engage in any discussion of his trauma narrative (negative behavior = 3). In this example, negative trauma-related responses were activated at threshold in three of the four domains (cognitive, emotional, and behavioral), so the multimodal negative response score is 3. Another adolescent discussing her father’s murder may express the beliefs, “I know I’ll be able to move forward from this” and “I feel strong” (positive cognitions intensity = 3), report she is feeling pleased and proud of herself for how she was able to cope with the loss of her father (positive emotions = 2), and report that she tried to have a constructive conversation with a peer about her father’s death (positive behavior = 1). Positive physiological responses were not noted (positive physiological = 0). Positive trauma-related responses in this example were activated at threshold in two domains (cognitive and emotional), so the multimodal positive response score is 2.

Measures of treatment outcomes.

Internalizing and externalizing symptoms.: At pretreatment and at 6 months post-randomization (approximately posttreatment), caregivers completed the Child Behavior Checklist (CBCL; Achenbach & Rescorla, 2001). The CBCL is a well-established measure of mental health problems in children that has good reliability and validity (Achenbach & Rescorla, 2001). Caregivers rate 113 items assessing a range of child emotional and behavioral problems, each rated on a three-point Likert scale from 0 (*not true*) to 2 (*very true or often true*).

The internalizing and externalizing scales of the CBCL were the primary outcome measures used in this study, consistent with previous publications from this trial (Hayes et al., 2017; Ready et al., 2015; Yasinski et al., 2016). The internalizing scale of the CBCL measures anxiety, depression, withdrawal, and somatic complaints. The externalizing scale measures acting out, rule-breaking, and aggressive behaviors. Measures of broadband symptomatology have been recommended over more disorder-specific measures to capture therapeutic change in youth treatment (Becker, Chorpita, & Daleiden, 2011). Further, PTSD includes many internalizing symptoms (e.g., persistent negative emotions, reduced interest, sleep disturbance), but it is also associated with higher levels of externalizing symptoms in youth (Barboza, Dominguez, & Pinder, 2017; Saigh, Yasik, Oberfield, Halamandaris, & McHugh, 2002). Data in adult populations also suggest that more transdiagnostic internalizing and externalizing symptom measures might better capture the range of trauma-related pathology than PTSD-specific measures (Wolf, Miller, Harrington, & Reardon, 2012). Thus, we examined the internalizing and externalizing scales of the CBCL as treatment outcomes. We used raw scores, as T-scores can truncate the range of data and be less sensitive to gradual symptom changes during treatment (Achenbach, 1991). In the current sample, reliability across time points was excellent for both the internalizing (Cronbach's $\alpha = .89$ to $.90$) and externalizing scales ($\alpha = .92$ to $.95$).

PTSD symptoms.: A measure of PTSD symptoms, the self-report version of the UCLA PTSD Reaction Index for DSM-IV (UPID; Steinberg et al., 2004), was completed by youth at baseline and 6 months post-randomization. The UPID can be administered as a questionnaire or structured interview to youth ages 7–18, and it assesses various trauma types and the severity of DSM-IV PTSD symptoms. The UPID shows good test-retest reliability, convergent validity, sensitivity, and specificity in diagnosing PTSD (Steinberg et al., 2004). In the present sample, the UPID had good reliability (Cronbach's $\alpha = .87$ to $.90$).

Data Analytic Approach

Data were analyzed in Mplus 7 (Muthén & Muthén, 1998–2012). Multilevel modeling was used to account for the hierarchical structure of the data (i.e., sessions nested within participants). Within-person slopes of scores for multimodal negative and positive trauma-related responses were entered as predictors of posttreatment outcomes (internalizing, externalizing, and PTSD symptoms). Random slopes were estimated to allow for slopes to vary from person to person, and random slopes can be treated as predictors of between-person outcomes. Two sets of models were repeated across internalizing, externalizing, and

PTSD outcomes for a total of six models. Child age and gender were uncorrelated with all predictor and outcome measures, so they were not included in the final analyses.

In the first three models, within-person slopes of multimodal negative and positive response scores were estimated to represent each individual's linear change in these variables over time. At the within-person level of each model, random slopes of multimodal negative and positive responses were estimated using each individual's repeated measures by regressing each of these variables on session number, with the first session coded as 0. At the between-person level, posttreatment internalizing (Model 1), externalizing (Model 2), or PTSD (Model 3) was regressed on the within-person slopes of both negative and positive responses, controlling for the baseline levels of the outcome variable (e.g., the model predicting posttreatment internalizing included baseline internalizing as a covariate). We controlled for baseline symptoms to account for initial severity and capture changes in symptoms.

In the final three models, the within-person quadratic slope of multimodal negative responses was entered as a predictor of posttreatment internalizing (Model 4), externalizing (Model 5), and PTSD symptoms (Model 6). In order to reduce collinearity between linear and quadratic time terms, session number was grand-mean centered in these analyses. At the within-person level, random slopes were estimated using each individual's repeated measures by regressing negative responses on linear (i.e., centered session number) and quadratic (centered session number squared) session terms. Change in positive responses was not included as a covariate because models including this variable failed to converge. At the between-person level of each model, the treatment outcome (internalizing, externalizing, or PTSD) was regressed on the within-person quadratic slope of change in multimodal negative responses, controlling for the baseline level of the outcome variable.

Results

Descriptive Analyses and Intercorrelations

Descriptive analyses for within-person (session-by-session) variables of interest and their intercorrelations revealed that multimodal negative trauma-related responses ($M = 1.34$, $SD = 1.01$, range 0–4) and multimodal positive responses ($M = .88$, $SD = .91$, range 0–3) were not significantly correlated ($r = .10$, n.s.). On average across sessions in the trauma processing phase of TF-CBT, participants had higher multimodal negative responses than positive responses ($t(412) = 7.42$, $p < .001$). Descriptive statistics for between-person variables (internalizing, externalizing, and PTSD) at baseline and posttreatment and their intercorrelations are presented in Table 2. Internalizing and externalizing symptom severity measures were positively correlated at all time points. Baseline PTSD severity was only correlated significantly with baseline internalizing.

We also examined the fixed and random effects of within-person slopes over time for the multimodal negative and positive response variables that were the predictor variables in the main analyses. None of the slopes of interest were significant on average across the sample (fixed effect = .02 for negative responses linear slope, $-.02$ for negative responses quadratic slope, .04 for positive responses linear slope, all n.s.). The linear slope of negative responses

over time varied significantly across participants (random effect = .01, $p < .05$), but the other slopes of interest did not (random effect = .002 for negative responses quadratic, .01 for positive responses linear, both n.s.). Based on estimates of the averages and variances of these slopes, 95% of the sample ranged from negative to positive values for all slopes of interest (95% sample range = $-.19$ to $.24$ for negative responses linear, $-.11$ to $.06$ for negative responses quadratic, $-.13$ to $.20$ for positive responses linear).

Predictors of Treatment Outcomes

The first three models regressed internalizing, externalizing, and PTSD outcomes respectively on within-person linear slopes of multimodal negative and positive responses, controlling for baseline symptoms. Results of these models (see Table 3) suggest that linear change in negative trauma-related responses over the trauma processing phase of TF-CBT did not predict any of the three posttreatment outcomes of interest. However, as predicted, more linear improvement in positive responses predicted lower posttreatment externalizing (but not internalizing or PTSD) symptoms. The final three models regressed internalizing, externalizing, and PTSD outcomes respectively on the within-person quadratic slope of multimodal negative responses. Results (Table 3) indicate that while the linear slope of negative responses did not predict posttreatment outcomes, the quadratic shape of change did predict posttreatment internalizing and PTSD outcomes, as predicted, but not externalizing symptoms. The effect was such that a more concave down curve (inverted U) predicted lower posttreatment internalizing and PTSD symptoms.

Discussion

The current study extends prior research on processes of change in TF-CBT and other PTSD treatments by moving from a focus on isolated variables (e.g., negative cognitions, fear response) to an examination of multimodal pathological and adaptive trauma responses as predictors of treatment outcome. Using Foa and colleagues' (2006) updated emotional processing theory as an organizational framework, we examined linear and nonlinear changes in the extent to which pathological and adaptive responses to trauma-related content were activated across cognitive, emotional, behavioral, physiological domains. We examined the following predictors of outcome: 1) linear decreases in multimodal pathological responses, 2) curvilinear change (concave down, inverted U) in multimodal pathological responses, and 3) linear increases in multimodal adaptive responses. Linear change in multimodal negative responses did not predict treatment outcomes, but as hypothesized, a *curvilinear* pattern of negative responses did predict lower posttreatment internalizing and PTSD symptoms, though not externalizing symptoms. Also as predicted, linear increases in multimodal positive responses predicted less externalizing behavior at posttreatment, but unexpectedly did not predict improvement in internalizing or PTSD symptoms.

Most studies examining processes of change in TF-CBT and other PTSD treatments have focused on linear changes in single domains of responses to trauma content, and few predictors of TF-CBT outcome have been identified. Consistent with the most recent iteration of EPT (Foa et al., 2006), our findings suggest that linear and nonlinear changes in multimodal pathological and adaptive responses are important in TF-CBT. Although some

studies have described a curvilinear pattern of symptom change across treatment (Nishith et al., 2002; Schumm et al., 2013), they did not examine this pattern as a predictor of treatment outcomes. Our study is the second to our knowledge to examine curvilinear change in trauma responses across treatment sessions as a *predictor* of symptom outcomes. Robison-Andrew and colleagues (2014) found that a quadratic pattern of change in startle response measured at pre-, mid-, and posttreatment predicted better treatment response in prolonged exposure and present-centered therapy for PTSD. Similarly, we found that curvilinear change in multimodal pathological responses (cognitive, emotional, behavioral, and physiological) predicted improvement in both internalizing symptoms and PTSD symptoms. This concave down pattern might represent the activation of multiple domains of responses in the initial trauma processing sessions of TF-CBT followed by weakening of these responses (fewer domains activated at a moderate to high level) over subsequent sessions. This finding suggests that the trauma processing phase of TF-CBT can involve a transient period of distress as the traumatic experiences are recalled and processed, but an increase and decrease in distress is associated with symptom improvement. Although our findings need to be replicated, they suggest the promise of multimodal measures of pathological and adaptive learning and of nonlinear analyses of change (Foa et al., 2006; Hayes, Laurenceau, et al., 2007).

A notable pattern of findings was that change in both negative and positive multimodal responses to trauma content predicted better posttreatment outcomes on one or two symptom measures, but not all three. Negative and positive responses were not significantly correlated in the present sample, suggesting that positive responses represent more than the inverse of negative responses and that both can provide important information. This pattern of findings suggests that changes in pathological responses, which predicted improvement in internalizing and PTSD symptoms, may operate on different sets of symptoms than the development of more adaptive responses, which predicted improvement in externalizing symptoms. There is little research on predictors of externalizing symptoms related to trauma or on the role of developing more adaptive responses. Deblinger, Mannarino, Cohen, Runyon, and Steer (2011) found that more time teaching caregivers parenting skills in TF-CBT was associated with more improvement in child externalizing symptoms. It is possible that increases in more positive responses might have helped youth to better regulate themselves and respond to trauma-related situations with less anger, aggression, and impulsivity. Anger in particular predicts worse outcomes and higher dropout in trauma-focused treatment in adults (Foa et al, 1995; Rizvi et al., 2009), and it might also be an important target in treatments for traumatized youth.

Strengths, Limitations, and Future Directions

One strength of the present study is the repeated, session-by-session measurement of responses to trauma-related content over the phase of TF-CBT that targets trauma exposure and processing. The longitudinal study design allowed for the estimation of within-person slopes of change in pathological and adaptive trauma-related responses, which could then be examined as predictors of subsequent posttreatment outcomes. Because internalizing, externalizing, and PTSD symptoms were only measured at pretreatment and posttreatment rather than at each session, we cannot rule out the possibility that symptom change in the

first phase of treatment influenced change in responses to trauma content in the second phase of treatment. In addition, therapists might have responded differently to clients based on previous symptom change (Stiles, 2009). The current study provides preliminary evidence that changes in multimodal trauma responses predict posttreatment outcomes, but future studies should include session-by-session measurements of symptoms to better establish the temporal precedence of changes in negative and positive trauma responses (Kazdin, 2007). Future studies might also measure therapist interventions during sessions to reveal potential predictors of therapeutic change.

Another strength of the present study is that the CHANGE coding system (Hayes, Feldman, et al., 2007) allowed for the measurement of multiple types of responses to trauma content in one study and as rated by independent raters. The CHANGE can be useful as a single measure that captures multiple domains of pathological and adaptive trauma responses expressed during treatment sessions, in line with EPT's emphasis on multimodal activation and change (Foa et al., 2006). This measure also reduces the possible confound of method variance. However, the CHANGE is reliant on observable verbal and nonverbal content. Future studies could include different types of measures for each of the domains assessed (cognitive, emotional, behavioral, and physiological), including self-reports, lab tasks, behavioral assessments, and psychophysiological measures.

While the constructs that we examined involved multimodal sets of responses, we calculated composite scores rather than conducting network analyses. This was because the density of within-person data and the number of participants were not sufficient for network analyses (Epskamp, Borsboom, & Fried, 2018). With assessments that are daily or more frequent (rather than weekly), network analyses could be used to examine the interconnectivity among the types of responses and to identify those that are most central in predicting outcome (Borsboom, 2017; Epskamp et al., 2018). The present approach still has clinical utility in that findings suggest that clinicians might attend to changes in the *breadth* of their clients' responses to trauma-related content rather than focusing on one type of response or another (e.g., cognitive or emotional). Protocols for exposure therapies for PTSD, including TF-CBT, emphasize multimodal activation during exposure sessions, including sensory details, thoughts, and feelings (Cohen et al., 2006; 2017; Foa et al., 2007; Resick et al., 2017; Resick & Schnicke 1992), but less attention is paid to change in breadth of negative responses over the course of treatment or to the development of multimodal adaptive responses.

Another notable strength of this study is the use of emotional processing theory to guide the investigation of processes of change in TF-CBT, and the examination of change processes in a racially and ethnically diverse community sample of trauma-exposed youth. EPT has typically been studied in adult populations, and our findings contribute to literature showing that EPT is also relevant in treatments for youth (e.g., Cohen et al., 2006; 2017; Foa, McLean, Capaldi, & Rosenfield, 2013; Franklin et al., 2011; Peterman et al., 2019) and in treatments delivered in community settings (e.g., Foa et al., 2005; Webb et al., 2014).

It should also be noted that the present sample only included youth who began the trauma processing (second) phase of TF-CBT. While there were no differences on demographic

measures between these youth and those who discontinued therapy before that point, there could be other differences between these groups. In addition, we focused on the trauma processing phase of TF-CBT, where the focus is on the activation and processing of traumatic memories and related content. Important processes of change are also likely to occur in the other phases of TF-CBT, and these should be examined in future studies.

Another consideration is that some of the regression coefficients that were not significant predictors of outcome had large standard errors and confidence intervals. These included the linear slope of negative responses predicting posttreatment externalizing symptoms, linear slopes of negative and positive responses predicting PTSD symptoms, and the quadratic slope of negative responses predicting externalizing symptoms. Such a pattern could be related to low reliability of the predictor variables, which would increase risk of Type I error; however, the intraclass correlations were all in the good to excellent range (.66 to .91; Cicchetti, 1994), and all of the analyses in question produced nonsignificant findings. Further, other models including the same predictors had much smaller standard errors and confidence intervals, suggesting that the problem was not with the predictor variables. It is possible that we were somewhat underpowered to detect significant effects, due to the nature of the models (i.e., using within-person slopes as predictors) relative to the sample size. These findings will require replication in a larger sample to assure that null findings were not due to insufficient power.

Conclusion

Overall, the present findings add to the literature on processes of change in TF-CBT. Consistent with the broader framework of emotional processing theory (Foa et al., 2006), we found that an increase and then decrease (a curvilinear pattern) of multimodal pathological responses over the course of the trauma processing phase of TF-CBT predicted improvement in internalizing and PTSD symptoms. In addition, increases in multimodal adaptive responses predicted improvement in externalizing symptoms. These findings have implications for both research and clinical practice. Researchers might broaden the measurement of emotional processing and processes of change in PTSD treatments to include pathological and adaptive trauma-related responses across multiple domains, including cognitions, emotions, behaviors, and physiological responses. Our findings also illustrate the importance of examining not only linear, but also nonlinear patterns of change. TF-CBT clinicians might target activation and change in trauma responses across multiple domains, facilitate an increase then a decrease in negative responses during trauma processing, and also help clients to develop new, more positive responses to traumatic experiences.

Acknowledgements

This research was supported by grants from the National Institute of Mental Health (NIMH: R01-MH086558) awarded to Adele M. Hayes and from the Substance Abuse and Mental Health Services Administration (SAMHSA: SM57145) awarded to the Delaware Division of Prevention and Behavioral Health Services. We would like to thank our coding team for their help, Jean-Philippe Laurenceau and Emily Soriano for feedback on statistical analyses, and Edna Foa for her helpful comments on a previous draft of this manuscript.

References

- Abel A, Hayes AM, Henley W, & Kuyken W (2016). Sudden gains in cognitive-behavior therapy for treatment-resistant depression: Processes of change. *Journal of Consulting and Clinical Psychology*, 84(8), 726–737. 10.1037/ccp0000101 [PubMed: 27100125]
- Achenbach TM (1991). *Manual for the Child Behavior Checklist 4–18 and 1991 profile*. Burlington: Department of Psychiatry, University of Vermont.
- Achenbach TM, & Rescorla LA (2001). *Manual for the ASEBA school-age forms & profiles*. Burlington, VT: University of Vermont, Research Center for Children, Youth, & Families.
- Alpert E, Hayes AM, Barnes JB, & Sloan DM (2020). Predictors of dropout in cognitive processing therapy for PTSD: An examination of trauma narrative content. *Behavior Therapy*, 51(5), 774–788. 10.1016/j.beth.2019.11.003 [PubMed: 32800305]
- Asnaani A, McLean CP, & Foa EB (2016). Updating Watson & Marks (1971): How has our understanding of the mechanisms of extinction learning evolved and where is our field going next? *Behavior Therapy*, 47(5), 654–668. 10.1016/j.beth.2016.02.003 [PubMed: 27816078]
- Barboza GE, Dominguez S, & Pinder J (2017). Trajectories of post-traumatic stress and externalizing psychopathology among maltreated foster care youth: A parallel process latent growth curve model. *Child Abuse & Neglect*, 72, 370–382. 10.1016/j.chiabu.2017.09.007 [PubMed: 28917187]
- Becker KD, Chorpita BF, & Daleiden EL (2011). Improvement in symptoms versus functioning: How do our best treatments measure up? *Administration and Policy in Mental Health and Mental Health Services Research*, 38(6), 440–458. 10.1007/s10488-010-0332-x [PubMed: 21207129]
- Borsboom D (2017). A network theory of mental disorders. *World Psychiatry*, 16(1), 5–13. 10.1002/wps.20375 [PubMed: 28127906]
- Bouton ME (2000). A Learning theory perspective on lapse, relapse, and the maintenance of behavior change. *Health Psychology*, 19 (1 Suppl), 57–63. 10.1037/0278-6133.19.Suppl1.57 [PubMed: 10709948]
- Cicchetti DV (1994). Guidelines, criteria, and rules of thumb for evaluating normed and standardized assessment instruments in psychology. *Psychological Assessment*, 6(4), 284–290. 10.1037/1040-3590.6.4.284
- Cisler JM, Sigel BA, Kramer TL, Smitherman S, Vanderzee K, Pemberton J, & Kilts CD (2015). Amygdala response predicts trajectory of symptom reduction during trauma-focused cognitive-behavioral therapy among adolescent girls with PTSD. *Journal of Psychiatric Research*, 71, 33–40. 10.1016/j.jpsychires.2015.09.011 [PubMed: 26522869]
- Cohen JA, Mannarino AP, & Deblinger E (2006). *Treating trauma and traumatic grief in children and adolescents*. New York, NY: Guilford Press.
- Cohen JA, Mannarino AP, & Deblinger E (2017). *Treating trauma and traumatic grief in children and adolescents (2nd Ed.)*. New York, NY: Guilford Press.
- Cohen JA, Mannarino AP, & Deblinger E (2018). In Weisz JR & Kazdin AE (Eds.), *Trauma-focused cognitive-behavioral therapy for traumatized children (3rd ed. ed.)*. New York, NY: Guilford Press.
- Colgan DD, Christopher M, Michael P, & Wahbeh H (2016). The body scan and mindful breathing among veterans with PTSD: Type of intervention moderates the relationship between changes in mindfulness and post-treatment depression. *Mindfulness*, 7(2), 372–383. 10.1007/s12671-015-0453-0 [PubMed: 32863982]
- Cooper AA, Clifton EG, & Feeny NC (2017). An empirical review of potential mediators and mechanisms of prolonged exposure therapy. *Clinical Psychology Review*, 56, 106–121. 10.1016/j.cpr.2017.07.003 [PubMed: 28734184]
- Cooper AA, Kline AC, Graham B, Bedard-Gilligan M, Mello PG, Feeny NC, & Zoellner LA (2017). Homework “dose,” type, and helpfulness as predictors of clinical outcomes in prolonged exposure for PTSD. *Behavior Therapy*, 48(2), 182–194. 10.1016/j.beth.2016.02.013 [PubMed: 28270329]
- Craske MG, Kircanski K, Zelikowsky M, Mystkowski J, Chowdhury N, & Baker A (2008). Optimizing inhibitory learning during exposure therapy. *Behaviour Research and Therapy*, 46(1), 5–27. 10.1016/j.brat.2007.10.003 [PubMed: 18005936]

- Deblinger E, Mannarino AP, Cohen JA, Runyon MK, & Steer RA (2011). Trauma-focused cognitive behavioral therapy for children: Impact of the trauma narrative and treatment length. *Depression and Anxiety*, 28(1), 67–75. 10.1002/da.20744 [PubMed: 20830695]
- Ditto B, Eclache M, & Goldman N (2006). Short-term autonomic and cardiovascular effects of mindfulness body scan meditation. *Annals of Behavioral Medicine*, 32(3), 227–234. 10.1207/s15324796abm3203_9 [PubMed: 17107296]
- Dondanville KA, Blankenship AE, Molino A, Resick PA, Wachen JS, Mintz J, ... Peterson AL (2016). Qualitative examination of cognitive change during PTSD treatment for active duty service members. *Behaviour Research and Therapy*, 79, 1–6. 10.1016/j.brat.2016.01.003 [PubMed: 26874683]
- Dour HJ, Brown LA, & Craske MG (2016). Positive valence reduces susceptibility to return of fear and enhances approach behavior. *Journal of Behavior Therapy and Experimental Psychiatry*, 50, 277–282. 10.1016/j.jbtep.2015.09.010 [PubMed: 26497447]
- Epskamp S, Borsboom D, & Fried EI (2018). Estimating psychological networks and their accuracy: A tutorial paper. *Behavior Research Methods*, 50(1), 195–212. 10.3758/s13428-017-0862-1 [PubMed: 28342071]
- Foa EB, Hembree EA, Cahill SP, Rauch SAM, Riggs DS, Feeny NC, & Yadin E (2005). Randomized trial of prolonged exposure for posttraumatic stress disorder with and without cognitive restructuring: Outcome at academic and community clinics. *Journal of Consulting and Clinical Psychology*, 73(5), 953–964. 10.1037/0022-006X.73.5.953 [PubMed: 16287395]
- Foa EB, Hembree EA, & Rothbaum BO (2007). *Prolonged exposure therapy for PTSD: Emotional processing of traumatic experiences: Therapist guide*. New York, NY: Oxford University Press.
- Foa EB, Huppert JD, & Cahill SP (2006). Emotional processing theory: An update. In Rothbaum BO (Ed.), *Pathological anxiety: Emotional processing in etiology and treatment* (pp. 3–24). New York, NY: Guilford Press.
- Foa EB, & Kozak MJ (1986). Emotional processing of fear: Exposure to corrective information. *Psychological Bulletin*, 99(1), 20–35. 10.1037/0033-2909.99.1.20 [PubMed: 2871574]
- Foa EB, & McLean CP (2016). The efficacy of exposure therapy for anxiety-related disorders and its underlying mechanisms: The case of OCD and PTSD. *Annual Review of Clinical Psychology*, 12, 1–28. 10.1146/annurev-clinpsy-021815-093533
- Foa EB, McLean CP, Capaldi S, & Rosenfield D (2013). Prolonged exposure vs. supportive counseling for sexual abuse-related PTSD in adolescent girls: A randomized clinical trial. *JAMA: Journal of the American Medical Association*, 310(24), 2650–2657. 10.1001/jama.2013.282829 [PubMed: 24368465]
- Foa EB, & McNally RJ (1996). Mechanisms of change in exposure therapy. In Rapee RM (Ed.), *Current controversies in the anxiety disorders* (pp. 214–227). New York, NY: Guilford Press.
- Foa EB, Riggs DS, Massie ED, & Yarczower M (1995). The impact of fear activation and anger on the efficacy of exposure treatment for posttraumatic stress disorder. *Behavior Therapy*, 26(3), 487–499. 10.1016/S0005-7894(05)80096-6
- Franklin ME, Sapyta J, Freeman JB, Khanna M, Compton S, Almirall D, ... March JS (2011). Cognitive behavior therapy augmentation of pharmacotherapy in pediatric obsessive-compulsive disorder: The pediatric OCD treatment study II (POTS II) randomized controlled trial. *JAMA: Journal of the American Medical Association*, 306(11), 1224–1232. 10.1001/jama.2011.1344 [PubMed: 21934055]
- Gloth CA (2017). *The role of client avoidance on PTSD recovery throughout the course of trauma therapy* (Unpublished doctoral dissertation). University of Missouri - Saint Louis, Saint Louis, MO.
- Gutermann J, Schreiber F, Matulis S, Schwartzkopff L, Deppe J, & Steil R (2016). Psychological treatments for symptoms of posttraumatic stress disorder in children, adolescents, and young adults: A meta-analysis. *Clinical Child and Family Psychology Review*, 19(2), 77–93. 10.1007/s10567-016-0202-5 [PubMed: 27059619]
- Hallgren KA (2012). Computing inter-rater reliability for observational data: An overview and tutorial. *Tutorials for Quantitative Methods in Psychology*, 8(1), 23–34. 10.20982/tqmp.08.1.p023

- Hayes AM, Beevers CG, Feldman GC, Laurenceau J, & Perlman C (2005). Avoidance and processing as predictors of symptom change and positive growth in an integrative therapy for depression. *International Journal of Behavioral Medicine*, 12(2), 111–122. 10.1207/s15327558ijbml202_9 [PubMed: 15901220]
- Hayes AM, Feldman GC, & Goldfried MR (2007). The change and growth experiences scale: A measure of insight and emotional processing. *Insight in psychotherapy*. (pp. 231–253). American Psychological Association, Washington, DC.
- Hayes AM, Laurenceau J, Feldman G, Strauss JL, & Cardaciotto L (2007). Change is not always linear: The study of nonlinear and discontinuous patterns of change in psychotherapy. *Clinical Psychology Review*, 27(6), 715–723. 10.1016/j.cpr.2007.01.008 [PubMed: 17316941]
- Hayes AM, & Yasinski C (2015). Pattern destabilization and emotional processing in cognitive therapy for personality disorders. *Frontiers in psychology*, 6. 10.3389/fpsyg.2015.00107
- Hayes AM, Yasinski C, Grasso D, Ready CB, Alpert E, McCauley T, ... Deblinger E (2017). Constructive and unproductive processing of traumatic experiences in trauma-focused cognitive-behavioral therapy for youth. *Behavior Therapy*, 48(2), 166–181. 10.1016/j.beth.2016.06.004 [PubMed: 28270328]
- Held P, Owens GP, Monroe JR, & Chard KM (2017). Increased mindfulness skills as predictors of reduced trauma-related guilt in treatment-seeking veterans. *Journal of Traumatic Stress*, 30(4), 425–431. 10.1002/jts.22209 [PubMed: 28741736]
- Iverson KM, King MW, Cunningham KC, & Resick PA (2015). Rape survivors' trauma-related beliefs before and after cognitive processing therapy: Associations with PTSD and depression symptoms. *Behaviour Research and Therapy*, 66, 49–55. 10.1016/j.brat.2015.01.002 [PubMed: 25698164]
- Jun J, Keller S Zoellner LA, Feeny N, & Hayes AM (2016, 11). Examination of processes of change and sudden gains in posttraumatic stress disorder (PTSD) treatment. In Jun J (Chair), *Psychotherapy process-oriented assessment to enhance trauma-focused treatment: In-depth clinical exploration of key change processes*. Symposium conducted at the annual meeting of the Association for Behavioral and Cognitive Therapies, New York, NY.
- Kaczurkin AN, Asnaani A, Zhong J, & Foa EB (2016). The moderating effect of state anger on treatment outcome in female adolescents with PTSD. *Journal of Traumatic Stress*, 29(4), 325–331. 10.1002/jts.22116 [PubMed: 27459380]
- Kazdin AE (2007). Mediators and mechanisms of change in psychotherapy research. *Annual Review of Clinical Psychology*, 3, 1–27. 10.1146/annurev.clinpsy.3.022806.091432
- König J, Kopp B, Ziegelmeier A, Rimane E, Steil R, Renneberg B, & Rosner R (2019). Young people's trauma-related cognitions before and after cognitive processing therapy for post-traumatic stress disorder. *Psychology and Psychotherapy: Theory, Research and Practice*. Advance online publication. 10.1111/papt.12263
- Lenz AS, & Hollenbaugh KM (2015). Meta-analysis of trauma-focused cognitive behavioral therapy for treating PTSD and co-occurring depression among children and adolescents. *Counseling Outcome Research and Evaluation*, 6(1), 18–32. 10.1177/2150137815573790
- Maples-Keller JL, Rauch SAM, Jovanovic T, Yasinski CW, Goodnight JM, ... Norrholm SD (2019). Changes in trauma-potentiated startle, skin conductance, and heart rate within prolonged exposure therapy for PTSD in high and low treatment responders. *Journal of Anxiety Disorders*, 68, 102–147. 10.1016/j.janxdis.2019.102147
- Mazzei RL (2017). Mindfulness and meditation for reducing stress. In Wadhwa S (Ed.), *Stress in the modern world: Understanding science and society* (Vol. 1). Santa Barbara, CA: Greenwood Press/ABC-CLIO.
- McLean CP, & Foa EB (2017). Emotions and emotion regulation in posttraumatic stress disorder. *Current Opinion in Psychology*, 14, 72–77. 10.1016/j.copsyc.2016.10.006 [PubMed: 28813323]
- McLean CP, Su Y, & Foa EB (2015). Mechanisms of symptom reduction in a combined treatment for comorbid posttraumatic stress disorder and alcohol dependence. *Journal of Consulting and Clinical Psychology*, 83(3), 655–661. 10.1037/ccp0000024 [PubMed: 26009787]
- McLean CP, Yeh R, Rosenfield D, & Foa EB (2015). Changes in negative cognitions mediate PTSD symptom reductions during client-centered therapy and prolonged exposure for

- adolescents. *Behaviour Research and Therapy*, 68, 64–69. 10.1016/j.brat.2015.03.008 [PubMed: 25812826]
- McLean CP, Zang Y, Gallagher T, Suzuki N, Yarvis JS, Litz BT, Mintz J, Young-McCaughan S, Peterson AL, Foa EB, & The STRONG STAR Consortium. (2019). Trauma-related cognitions and cognitive emotion regulation as mediators of PTSD change among treatment-seeking active duty military personnel with PTSD. *Behavior Therapy Advance online publication*. 10.1016/j.beth.2019.03.006
- Muthén LK and Muthén BO (1998–2012). *Mplus user's guide* (7th ed.). Los Angeles, CA: Muthén & Muthén.
- Nishith P, Resick PA, & Griffin MG (2002). Pattern of change in prolonged exposure and cognitive-processing therapy for female rape victims with posttraumatic stress disorder. *Journal of Consulting and Clinical Psychology*, 70(4), 880–886. 10.1037/0022-006X.70.4.880 [PubMed: 12182271]
- Paivio SC, Hall IE, Holowaty KAM, Jellis JB, & Tran N (2001). Imaginal confrontation for resolving child abuse issues. *Psychotherapy Research*, 11(4), 433–453. 10.1093/ptr/11.4.433
- Paivio SC, & Nieuwenhuis JA (2001). Efficacy of emotion focused therapy for adult survivors of child abuse: A preliminary study. *Journal of Traumatic Stress*, 14(1), 115–133. 10.1023/A:1007891716593
- Paolucci EO, Genuis ML, & Violato C (2001). A meta-analysis of the published research on the effects of child sexual abuse. *The Journal of Psychology: Interdisciplinary and Applied*, 135(1), 17–36. 10.1080/00223980109603677 [PubMed: 11235837]
- Perrin S, Leigh E, Smith P, Yule W, Ehlers A, & Clark DM (2017). Cognitive therapy for PTSD in children and adolescents. In Landolt MA, Cloitre M, & Schnyder U (Eds.), *Evidence-based treatments for trauma related disorders in children and adolescents* (pp. 187–207). Cham: Springer International Publishing.
- Peterman JS, Carper MM, & Kendall PC (2019). Testing the habituation-based model of exposures for child and adolescent anxiety. *Journal of Clinical Child & Adolescent Psychology*, 48, S34–S44. 10.1080/15374416.2016.1163707 [PubMed: 27355694]
- Pfeiffer E, Sachser C, de Haan A, Tutus D, & Goldbeck L (2017). Dysfunctional posttraumatic cognitions as a mediator of symptom reduction in trauma-focused cognitive behavioral therapy with children and adolescents: Results of a randomized controlled trial. *Behaviour Research and Therapy*, 97, 178–182. 10.1016/j.brat.2017.08.001 [PubMed: 28810136]
- Porter E, Romero EG, & Barone MD (2018). Description and preliminary outcomes of an in vivo exposure group treatment for posttraumatic stress disorder. *Journal of Traumatic Stress*, 31(3), 410–418. 10.1002/jts.22285 [PubMed: 29924421]
- Ready CB, Hayes AM, Yasinski CW, Webb C, Gallop R, Deblinger E, & Laurenceau J (2015). Overgeneralized beliefs, accommodation, and treatment outcome in youth receiving trauma-focused cognitive behavioral therapy for childhood trauma. *Behavior Therapy*, 46(5), 671–688. 10.1016/j.beth.2015.03.004 [PubMed: 26459847]
- Resick PA, Monson CM, & Chard KM (2017). *Cognitive processing therapy for PTSD: A comprehensive manual*. New York, NY: Guilford Press.
- Resick PA, & Schnicke MK (1992). Cognitive processing therapy for sexual assault victims. *Journal of Consulting and Clinical Psychology*, 60(5), 748–756. [PubMed: 1401390]
- Rizvi SL, Vogt DS, & Resick PA (2009). Cognitive and affective predictors of treatment outcome in cognitive processing therapy and prolonged exposure for posttraumatic stress disorder. *Behaviour Research and Therapy*, 47(9), 737–743. 10.1016/j.brat.2009.06.003 [PubMed: 19595295]
- Robison-Andrew E, Duval ER, Nelson CB, Echeverri-Cohen A, Giardino N, Defever A, ... Rauch SAM (2014). Changes in trauma-potentiated startle with treatment of posttraumatic stress disorder in combat veterans. *Journal of Anxiety Disorders*, 28(4), 358–362. 10.1016/j.janxdis.2014.04.002 [PubMed: 24786361]
- Roth S, Newman E, Pelcovitz D, van der Kolk B, & Mandel FS (1997). Complex PTSD in victims exposed to sexual and physical abuse: Results from the DSM-IV field trial for posttraumatic stress disorder. *Journal of Traumatic Stress*, 10(4), 539–555. 10.1023/A:1024837617768 [PubMed: 9391940]

- Saigh PA, Yasik AE, Oberfield RA, Halamandaris PV, & McHugh M (2002). An analysis of the internalizing and externalizing behaviors of traumatized urban youth with and without PTSD. *Journal of Abnormal Psychology*, 111(3), 462–470. 10.1037/0021-843X.111.3.462 [PubMed: 12150422]
- Schauer M, Neuner F, & Elbert T (2011). *Narrative exposure therapy: A short-term treatment for traumatic stress disorders* (2nd ed.). Cambridge, MA: Hogrefe Publishing.
- Scher CD, Suvak MK, & Resick PA (2017). Trauma cognitions are related to symptoms up to 10 years after cognitive behavioral treatment for posttraumatic stress disorder. *Psychological Trauma: Theory, Research, Practice, and Policy*, 9(6), 750–757. 10.1037/tra0000258
- Schumm JA, Walter KH, & Chard KM (2013). Latent class differences explain variability in PTSD symptom changes during cognitive processing therapy for veterans. *Psychological Trauma: Theory, Research, Practice, and Policy*, 5(6), 536–544. 10.1037/a0030359
- Sloan DM & Marx BP (2019). *Written Exposure Therapy for PTSD: A Brief Treatment Approach for Mental Health Professionals*. Washington, DC: American Psychological Press.
- Smith P, Yule W, Perrin S, Tranah T, Dalgleish T, & Clark DM (2007). Cognitive-behavioral therapy for PTSD in children and adolescents: A preliminary randomized controlled trial. *Journal of the American Academy of Child & Adolescent Psychiatry*, 46(8), 1051–1061. 10.1097/CHI.0b013e318067e288 [PubMed: 17667483]
- Sripada RK, Rauch SAM, & Liberzon I (2016). Psychological mechanisms of PTSD and its treatment. *Current Psychiatry Reports*, 18, 99. 10.1007/s11920-016-0735-9 [PubMed: 27671916]
- Steinberg AM, Brymer MJ, Decker KB, & Pynoos RS (2004). The University of California at Los Angeles post-traumatic stress disorder reaction index. *Current Psychiatry Reports*, 6, 96–100. 10.1007/s11920-004-0048-2 [PubMed: 15038911]
- Stiles WB (2009). Responsiveness as an obstacle for psychotherapy outcome research: It's worse than you think. *Clinical Psychology: Science and Practice*, 16(1), 86–91. 10.1111/j.1468-2850.2009.01148.x
- Tyler KA (2002). Social and emotional outcomes of childhood sexual abuse: A review of recent research. *Aggression and Violent Behavior*, 7(6), 567–589. 10.1016/S1359-1789(01)00047-7
- Webb C, Hayes AM, Grasso D, Laurenceau J, & Deblinger E (2014). Trauma-focused cognitive behavioral therapy for youth: Effectiveness in a community setting. *Psychological Trauma: Theory, Research, Practice, and Policy*, 6(5), 555–562. 10.1037/a0037364
- Weisz JR, & Kazdin AE (2010). The present and future of evidence-based psychotherapies for children and adolescents. In Weisz JR & Kazdin AE (Eds.), *Evidence-based psychotherapies for children and adolescents* (pp. 557–576). New York, NY: Guilford Press.
- Wolf EJ, Miller MW, Harrington KM, & Reardon A (2012). Personality-based latent classes of posttraumatic psychopathology: Personality disorders and the internalizing/externalizing model. *Journal of Abnormal Psychology*, 121(1), 256–262. 10.1037/a0023237 [PubMed: 21574669]
- Yasinski C, Hayes AM, Ready CB, Cummings JA, Berman IS, McCauley T, ... Deblinger, E. (2016). In-session caregiver behavior predicts symptom change in youth receiving trauma-focused cognitive behavioral therapy (TF-CBT). *Journal of Consulting and Clinical Psychology*, 84(12), 1066–1077. 10.1037/ccp0000147 [PubMed: 27618641]
- Zalta AK, Gillihan SJ, Fisher AJ, Mintz J, McLean CP, Yehuda R, & Foa EB (2014). Change in negative cognitions associated with PTSD predicts symptom reduction in prolonged exposure. *Journal of Consulting and Clinical Psychology*, 82(1), 171–175. 10.1037/a0034735 [PubMed: 24188512]
- Zbozinek TD, & Craske MG (2017). The role of positive affect in enhancing extinction learning and exposure therapy for anxiety disorders. *Journal of Experimental Psychopathology*, 8(1), 1–52. 10.5127/jep.052615

Table 1.

CHANGE coding categories with descriptions, examples of high levels of each variable, and intra-class correlations (ICCs) of inter-rater agreement. Content in each category had to be related to the trauma or its impact to be coded.

Coding Category	Description	Example	ICC
NEGATIVE RESPONSES			
Negative Self	Negative beliefs about the self, including expressions of negative self-worth, self-criticism, and feelings of failure.	"I feel like I'm broken. I don't think I'm strong enough to get past this."	.66
Negative View of Others	Perceived negative quality of relationships with others or interactions with others, including specific people and people in general.	"It makes me not want to trust people. I think most people probably want to hurt me like my dad hurt me."	.91
Negative Hope	Feelings of being stuck or having no way out, feeling tired of trying, or negative beliefs about the future.	"I feel so horrible, and it's always going to be like this. I can't see a way out."	.76
Negative Emotion	Rated based on the number and intensity of negative emotion words (e.g., anxious, sad, angry, ashamed, guilty) and quality of the emotional tone (e.g., crying).	"I hurt so bad. I feel guilty that my dad got arrested because of me, and I'm so embarrassed."	.82
Negative Behaviors	Maladaptive actions in which the person engages, rated based on number of behaviors and their intensity.	"Then I lost it and started yelling at him [brother]. I just kept yelling and telling him he didn't know what he was talking about."	.88
Negative Physiological	Negative physiological experiences resulting from a person's thinking or emotions.	"I feel really shaky, like I'm going to throw up."	.85
POSITIVE RESPONSES			
Positive Self	Positive beliefs about the self, including a sense of worth, competency, desirability, self-acceptance, and pride.	"I felt good that I was able to handle it when I got scared. I was proud of myself."	.78
Positive View of Others	Perceived positive quality of relationships with others or interactions with others, including specific people and people in general.	"I helped her [mother] with the cleaning, and we actually had a pretty good time. Then she took me out for dinner and it was really nice." (Child's relationship with their mother was previously strained for trauma-related reasons, so these interactions were considered therapeutic.)	.82
Positive Hope	Capacity to see possibility of change in the future, determination to making changes, or positive beliefs about the future.	"I have a scar on my heart, but it's healing. I really think I'll be okay."	.70
Positive Emotion	Rated based on the number and intensity of positive emotion words (e.g., happy, calm, excited) and quality of the emotional tone (e.g., genuine laughter).	"I feel better this week. I feel pretty calm. And I'm happy that I talked to my mom and told her my story."	.85
Positive Behaviors	Adaptive actions in which the person engages, or exhibiting control so as not to engage in a maladaptive behavior.	"Instead of punching him [boy in neighborhood], I took a deep breath and told him to stop talking to me like that."	.81
Positive Physiological	Positive physiological experiences resulting from a person's thinking or emotions.	"I tried doing those relaxation exercises you taught me, and I felt really calm and relaxed after."	.86

Note. ICC = intra-class correlation.

Table 2.

Descriptive statistics and intercorrelations among between-person outcome measures, including internalizing and externalizing scales of the Child Behavior Checklist (CBCL) and PTSD severity (UCLA PTSD Reaction Index for DSM-IV; UPID).

	Internalizing Baseline	Internalizing Posttreatment	Externalizing Baseline	Externalizing Posttreatment	PTSD Baseline	PTSD Posttreatment
Internalizing Baseline	–					
Internalizing Posttreatment	.57 ^{***}	–				
Externalizing Baseline	.64 ^{***}	.36 ^{**}	–			
Externalizing Posttreatment	.28 [*]	.61 ^{***}	.54 ^{***}	–		
PTSD Baseline	.25 [*]	–.02	.11	–.10	–	
PTSD Posttreatment	–.01	.04	.03	.14	.20	–
Mean	14.86	8.90	16.98	12.03	33.64	18.16
SD	9.00	7.92	13.17	11.12	13.20	11.93
Range	0–36	0–34	0–49	0–44	7–65	0–46

*
 $p < .05$

**
 $p < .01$

 $p < .001$

Regression results for within-person slopes of multimodal negative and positive responses to trauma-related content predicting posttreatment outcomes (internalizing, externalizing, and PTSD symptoms), controlling for baseline symptoms.

Table 3.

Model	Predictor (Within-Person Slope)	Posttreatment Outcome	b	β	SE	z	p	95% Confidence Interval	
								Lower	Upper
1	Multimodal Negative Responses: Linear	CBCL Internalizing Symptoms	61.84	.15	260.12	.24	.81	-448.00	571.68
	Multimodal Positive Responses: Linear		-138.32	-.52	177.89	-.78	.44	-486.98	210.35
2	Multimodal Negative Responses: Linear	CBCL Externalizing Symptoms	-33.69	-.05	1336.13	-.03	.98	-2652.51	2585.13
	Multimodal Positive Responses: Linear		-313.53	-.54	40.63	-7.72	<.01	-393.16	-233.89
3	Multimodal Negative Responses: Linear	UPID PTSD Symptoms	250.15	.38	506.22	.49	.62	-742.04	1242.33
	Multimodal Positive Responses: Linear		-121.33	-.26	721.22	-.17	.87	-1534.92	1292.25
4	Multimodal Negative Responses: Quadratic	CBCL Internalizing Symptoms	268.41	.21	50.83	5.28	<.01	168.78	368.03
5	Multimodal Negative Responses: Quadratic	CBCL Externalizing Symptoms	-714.59	-.72	535.32	-1.34	.18	-1763.81	334.65
6	Multimodal Negative Responses: Quadratic	UPID PTSD Symptoms	353.26	.18	67.63	5.22	<.01	220.70	485.81

Note. CBCL = Child Behavior Checklist. UPID = UCLA PTSD Reaction Index for DSM-IV. Multimodal Negative Responses is a composite variable representing the number of domains of pathological trauma responses (cognitive, emotional, behavioral, physiological) activated at a moderate to high level (2 on the 0–3 CHANGE scale) in response to trauma-related content during the trauma processing phase of trauma-focused cognitive behavioral therapy (TF-CBT). Multimodal Positive Responses is a composite variable representing the number of domains of adaptive trauma responses activated at the same threshold.

* $p < .05$

** $p < .01$