



EMORY
LIBRARIES &
INFORMATION
TECHNOLOGY

OpenEmory

Association of Tumor Necrosis Factor alpha Inhibitor Use with Diagnostic Features and Mortality of Tuberculosis in the United States, 2010-2017

[Kenneth Castro](#), *Emory University*
[Neela Goswami](#), *Emory University*
SS Katrak, *Calif Dept Publ Hlth*
R Li, *Ctr Dis Control & Prevent*
S Reynolds, *Ctr Dis Control & Prevent*
SM Marks, *Ctr Dis Control & Prevent*
JR Probst, *Emory University*
T Chorba, *Ctr Dis Control & Prevent*
K Winthrop, *Oregon Hlth & Sci Univ*

Journal Title: OPEN FORUM INFECTIOUS DISEASES

Volume: Volume 9, Number 2

Publisher: OXFORD UNIV PRESS INC | 2022-01-30, Pages ofab641-ofab641

Type of Work: Article

Publisher DOI: 10.1093/ofid/ofab641

Permanent URL: <https://pid.emory.edu/ark:/25593/vv2m8>

Final published version: <http://dx.doi.org/10.1093/ofid/ofab641>

Accessed December 9, 2022 12:47 AM EST

Association of Tumor Necrosis Factor α Inhibitor Use with Diagnostic Features and Mortality of Tuberculosis in the United States, 2010–2017

Shereen S. Katrak,^{1,2} Rongxia Li,³ Sue Reynolds,³ Suzanne M. Marks,³ Jessica R. Probst,⁴ Terence Chorbha,³ Kevin Winthrop,⁵ Kenneth G. Castro,⁴ and Neela D. Goswami^{3,4}

¹Tuberculosis Control Branch, California Department of Public Health, Richmond, California, USA, ²Division of Infectious Diseases, University of California, San Francisco, San Francisco, California, USA, ³Division of Tuberculosis Elimination, Centers for Disease Control and Prevention, Atlanta, Georgia, USA, ⁴Rollins School of Public Health, Emory University, Atlanta, Georgia, USA, and ⁵Division of Infectious Disease, Oregon Health and Science University, Portland, Oregon, USA

Background. An elevated risk of tuberculosis (TB) disease in persons who have received tumor necrosis factor alpha inhibitor medications (TNF- α inhibitors) has been reported for nearly two decades, but clinical diagnostic features and outcomes of TB in this population remain poorly described.

Methods. We analyzed national surveillance data for TB cases among persons aged 15 years and older reported in the United States during 2010–2017 and associated mortality data reported through 2019 to describe the clinical characteristics of those receiving TNF- α inhibitors.

Results. Of 70 129 TB cases analyzed, 504 (0.7%) of the patients had TNF- α inhibitor use reported at TB diagnosis. Patients with TNF- α inhibitor use at TB diagnosis were more likely than TB patients not receiving TNF- α inhibitors to have TB diagnosed in extrapulmonary sites in conjunction with pulmonary sites (28.8% vs 10.0%, $P < .001$). Patients receiving TNF- α inhibitors were less likely to have acid-fast bacilli noted on sputum smear microscopy (25.6% vs 39.1%, $P = .04$), and more likely to have drug-resistant disease (13.5% vs 10.0%, $P < .001$). TB-attributed deaths did not significantly differ between patients receiving and not receiving TNF- α inhibitors (adjusted odds ratio, 1.46 [95% confidence interval, .95–2.26]).

Conclusions. Clinicians evaluating TNF- α inhibitor–treated patients should have a high index of suspicion for TB and be aware that extrapulmonary or sputum smear–negative TB disease is more common in these patients. No significantly diminished survival of TB patients treated with TNF- α inhibitor therapy before TB diagnosis was noted.

Keywords. biologic; immunocompromised host; mycobacterial disease; tuberculosis; TNF- α inhibitor.

The risk of tuberculosis (TB) disease among persons with exposure to TB or with latent TB infection (LTBI) increases with use of tumor necrosis factor alpha (TNF- α) inhibitors, a class of medications that is widely used in treatment of autoimmune conditions such as rheumatoid arthritis, psoriasis, Crohn disease, and ulcerative colitis. TNF- α is an essential component of host response to mycobacterial diseases, including TB, and is required for regulation of cellular immunity and formation of granulomas [1–3]. Of the TNF- α inhibitors approved for use in the United States (US) during the period covered by this analysis (ie, infliximab, adalimumab, golimumab, etanercept, and certolizumab pegol),

infliximab and adalimumab are most associated with increased risk of LTBI progression to TB disease [4, 5], but all classes of TNF- α inhibitors can increase TB risk [1, 5, 6].

Elevated TB incidence in association with use of TNF- α inhibitors has been reported previously [7, 8]. However, the clinical presentation of TB in this context is not well-characterized, and most cohort data come from Asia [9–11], where baseline TB incidence is significantly higher than in the US. In a 2001 US report describing 70 incident cases of TB among patients receiving infliximab, >50% of cases were extrapulmonary, nearly 25% were disseminated, and >15% of patients died [12]. In contrast, a retrospective cohort study using Kaiser Health System data from 2000 to 2008 in the US found that among 16 TB cases in patients using TNF- α inhibitors, 11 (69%) had pulmonary disease only [13]. Although clinical outcomes of TB disease in patients receiving TNF- α inhibitors are not well-described, published data suggest that TB-associated death may be associated with use of immunosuppressive medication [14].

The purpose of this analysis is to describe clinical characteristics of US TB patients who were receiving TNF- α inhibitors at the time of their TB diagnosis, and to assess the relationship

Received 1 October 2021; editorial decision 14 December 2021; accepted 21 December 2021; published online 22 December 2021.

Correspondence: Shereen Katrak, MD, MPH, California Department of Public Health, 850 Marina Bay Parkway, Bldg P, Richmond, CA 94804, USA (shereen.katrak@cdph.ca.gov).

Open Forum Infectious Diseases®2022

© The Author(s) 2021. Published by Oxford University Press on behalf of Infectious Diseases Society of America. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work is properly cited. For commercial re-use, please contact journals.permissions@oup.com <https://doi.org/10.1093/ofid/ofab641>

between use of TNF- α inhibitors and TB-related death. Because of the body of published data, we hypothesized that use of TNF- α inhibitors would be associated with extrapulmonary disease in conjunction with pulmonary disease [15], as compared to pulmonary-only disease, and with increased mortality among TB patients [16].

METHODS

Design and Patient Population

TB is a nationally notifiable condition; verified cases of TB are reported to the Centers for Disease Control and Prevention's (CDC) National Tuberculosis Surveillance System (NTSS) [16]. We analyzed data for all patients with TB aged 15 years and older, reported to NTSS from the 50 states and District of Columbia during 2010–2017, and any associated mortality data on these patients reported through 2019. Data in the NTSS during this period were collected via the Report of Verified Case of Tuberculosis (RVCT) (office of management and budget number 0920-0026), which includes information on patient demographics, TB risk factors, diagnostics, initial drug regimen, and death before or during receipt of TB treatment. The RVCT form was updated in 2009 to include additional risk factors, new drug treatments, and additional diagnostic tests; thus, only reported cases after 2009 were included in this analysis. NTSS data are protected under an Assurance of Confidentiality provided under the authority granted to CDC by Sections 306 and 308(d) of the Public Health Service Act (42 US Code [U.S.C.] 242k and 242m[d]), which prevents disclosure of any information that could be used to identify patients directly or indirectly. This project was determined not to be human subjects research by CDC and did not require approval by an institutional review board because data were collected and analyzed as part of routine public health surveillance.

Exposure and Outcomes

The primary exposure studied was use of TNF- α inhibitor therapy at the time of TB diagnosis. This was documented as a checkbox completed by TB program staff on the RVCT for each TB case under "additional TB risk factors". Covariates of interest assessed at the time of TB diagnosis included age at diagnosis, sex, race/ethnicity, birth outside the US, diabetes, end-stage renal disease (ESRD), human immunodeficiency virus (HIV) infection, excess alcohol use in the past year, drug use (injecting or noninjecting) in the past year, homelessness in the past year, private provider outpatient care, residence in a long-term care institution at the time of diagnosis, residence in a correctional facility at the time of diagnosis, employment status, presence of smear positivity (for both pulmonary disease and extrapulmonary disease), site of TB disease, and presence of first-line drug resistance. Since cavitary pulmonary disease and miliary TB were highly correlated with site of disease,

these variables were excluded from logistic regression models. Covariates were chosen based on clinical relevance or previous association with TB-related morbidity or mortality [14].

The primary outcome of this analysis was TB-associated death, before starting TB treatment or during TB treatment. TB-associated death was defined by an RVCT response of (1) "status at TB diagnosis" reported as "dead" with "was TB a cause of death" reported as "yes"; or (2) "reason therapy stopped or never started" reported as "died" and "cause of death" reported as "related to TB disease" or "related to TB therapy." The outcome of death was considered complete for patients reported no later than 2010–2017 since jurisdictions have up to 2 years to submit the follow-up RVCT form (<https://www.cdc.gov/tb/statistics/reports/2019/outcomes.htm>).

Statistical Analysis

We described baseline characteristics of all patients and of patients with TB-associated deaths, stratified by receipt of TNF- α inhibitors. Both χ^2 tests or Fisher exact tests (when the expected cell size was <5) were used to examine whether unadjusted associations existed between TNF- α inhibitor use and all demographic and clinical risk factors, after exclusion of missing data. We used multiple imputation to impute missing values for the following variables: age at diagnosis, sex, race/ethnicity, birth outside the US, HIV infection, excess alcohol use in the past year, drug use in the past year, homelessness in the past year, employment status, and residence in a correctional facility at the time of diagnosis. The above variables were assumed to be missing at random [17] and the fully conditional specification technique [3] was employed to impute missing values. Five imputed datasets were generated, and point estimates were averaged and standard errors were derived following Rubin's rules [17]. Multiple imputation was only used to input values for missing variables considered to be missing at random. For variables not suspected to be missing at random, including outpatient provider type or smear results, the values for these variables were retained as "missing" in multivariable analyses. Odds ratios and associated 95% confidence intervals (CIs) were calculated to assess unadjusted associations between TNF- α inhibitor use and death. Statistically significant variables at the α level of .05, or those considered clinically relevant, along with TNF- α inhibitor status, were entered into a multivariable model. Multivariable logistic regression was used to estimate adjusted odds ratios (aORs) and associated 95% CIs of factors associated with TB-related death. All statistical analyses were performed using R statistical software [18].

RESULTS

In total, 70 129 TB cases were included in these analyses. Unadjusted results for TB patients by TNF- α inhibitor therapy status are presented in Table 1; missing data are shown in

Table 1. Characteristics of Patients With Tuberculosis by Tumor Necrosis Factor Alpha Inhibitor Therapy Status, United States, 2010–2017 (N = 70 129)^a

Characteristic	TNF- α Inhibitor (n = 504)		No TNF- α Inhibitor (n = 69 625)		P Value
	No.	(%)	No.	(%)	
Age at diagnosis, y					<.001
15–24	21	(4.2)	7 786	(11.2)	
25–44	130	(25.8)	23 832	(34.2)	
45–64	207	(41.1)	22 866	(32.8)	
\geq 65	146	(29.0)	15 134	(21.7)	
Sex					<.001
Female	269	(53.4)	26 981	(38.8)	
Male	235	(46.6)	42 637	(61.2)	
Race					<.001
Asian	206	(41.0)	23 164	(33.3)	
Black	50	(9.9)	15 212	(21.9)	
Hispanic	106	(21.1)	19 649	(28.3)	
Other ^b	9	(1.8)	1 784	(2.6)	
White	132	(26.2)	9 655	(13.9)	
Place of birth					.30
Non-US born	335	(66.5)	47 807	(68.7)	
US born	169	(33.5)	21 778	(31.3)	
Diabetes					.04
Not reported	434	(86.1)	57 496	(82.6)	
Reported	70	(13.9)	12 129	(17.4)	
End-stage renal disease					.97
Not reported	494	(98)	68 157	(97.9)	
Reported	10	(2)	1 468	(2.1)	
HIV status					<.001
HIV negative	426	(99.3)	56 951	(93.4)	
HIV positive	3	(0.7)	4 010	(6.6)	
Excess alcohol use past year					<.001
No	488	(97.6)	60 962	(88.7)	
Yes	12	(2.4)	7 826	(11.3)	
Drug use past year ^c					<.001
No	493	(98.0)	63 690	(92.6)	
Yes	10	(2.0)	5 104	(7.4)	
Homelessness past year ^d					<.001
No	503	(99.8)	65 317	(94.4)	
Yes	1	(0.2)	3 851	(5.6)	
Private outpatient provider					<.001
No	311	(61.7)	53 857	(77.5)	
Yes	193	(38.3)	15 661	(22.5)	
Resident long-term care facility ^e					.84
No	496	(98.6)	68 367	(98.4)	
Yes	7	(1.4)	1 117	(1.6)	
Resident of corrections facility ^f					<.001
No	501	(99.6)	66 517	(95.8)	
Yes	2	(0.4)	2 908	(4.2)	
Employment status					.002
Employed	250	(49.9)	29 100	(42.9)	
Not employed	251	(50.1)	38 676	(57.1)	
Initial sputum smear result					.04
Negative	291	(57.7)	33 970	(48.8)	
Positive	129	(25.6)	27 215	(39.1)	
Not done	84	(16.7)	8 403	(12.1)	
Initial smear result, any ^g					.34
Negative	227	(45.0)	31 708	(45.5)	
Positive	271	(53.8)	36 451	(52.4)	
Not done	6	(1.2)	1 454	(2.1)	

Table 1. Continued

Characteristic	TNF- α Inhibitor (n = 504)		No TNF- α Inhibitor (n = 69 625)		P Value
	No.	(%)	No.	(%)	
Site of TB disease					<.001
Both	145	(28.8)	6949	(10.0)	
Extrapulmonary	140	(27.8)	14 387	(20.7)	
Pulmonary	219	(43.5)	48 261	(69.3)	
Drug resistance ^h					<.001
No	333	(66.1)	47 199	(67.8)	
Yes	68	(13.5)	6941	(10.0)	
Not done	89	(17.7)	13 018	(18.7)	

Abbreviations: HIV, human immunodeficiency virus; TB, tuberculosis; TNF- α , tumor necrosis factor alpha; US, United States.

^aIncludes all TB cases age 15 years or older diagnosed in the US between 2010 and 2017. Percentages may not sum to 100% due to rounding. Missing data for each variable are presented in [Supplementary Table 1](#); column totals for each variable may not sum to total due to missing data. Percentages and P values were calculated excluding missing data.

^bThis category includes persons of American Indian/Alaska Native and Native Hawaiian/Other Pacific Islander races, as well as persons of multiple races.

^cInjection or noninjection drug use within 12 months prior to TB diagnosis.

^dHomeless within 12 months prior to TB diagnosis.

^eResident of long-term institution at time of diagnosis.

^fResident of correctional facility at time of diagnosis.

^gInitial smear result for any available specimen with reported acid-fast bacilli studies, including sputum, body fluid, or tissue.

^hPresence of resistance to isoniazid, rifampin, pyrazinamide, and/or ethambutol among culture-positive cases with drug susceptibility results.

Supplementary Table 1. Of the TB cases, 504 (0.7%) had TNF- α inhibitor use reported at time of TB diagnosis. Compared to patients not receiving TNF- α inhibitors, those receiving TNF- α inhibitors differed significantly in terms of race ($P < .001$) and age distribution ($P < .001$), and included higher proportions of Asian (41.0% vs 33.3%) and non-Hispanic White patients (26.2% vs 13.9%), patients aged 45–64 years (41.1% vs 32.8%) or ≥ 65 years (29.0% vs 21.7%), and women (53.4% vs 38.8%, $P < .001$). The proportion of patients born outside the US was similar in the 2 groups (66.5% vs 68.7%, $P = .3$). Compared to those not receiving TNF- α inhibitors, a higher percentage of TB patients receiving TNF- α inhibitors received outpatient TB care from a private healthcare provider (38.3% vs 22.5%, $P < .001$). Other reported sources of outpatient TB care included the public health department, Indian Health Service or tribal health department, hospital only, and correctional or institutional providers. A lower percentage of patients receiving TNF- α inhibitors were non-Hispanic Black (9.9% vs 21.9%) or Hispanic (21.1% vs 28.3%), reported HIV infection (0.7% vs 6.6%, $P < .001$), reported homelessness in the year preceding diagnosis (0.2% vs 5.6%, $P < .001$), or were unemployed (50.1% vs 57.1%, $P = .002$).

In unadjusted analyses ([Table 1](#)), site of TB disease varied significantly based on TNF- α inhibitor receipt ($P < .001$), with a greater percentage of patients who received TNF- α inhibitors manifesting disease in both pulmonary and extrapulmonary sites (28.8% vs 10.0%) or extrapulmonary only sites (27.8% vs 20.7%) compared with patients with only pulmonary disease. Patients receiving TNF- α inhibitors were less likely to have acid-fast bacilli (AFB) noted on sputum smear microscopy (25.6% vs 39.1%, $P = .04$). Among TNF- α inhibitor users with sputum specimens collected and examined, more than half were AFB

sputum smear negative (291/420 [69.3%]). TB patients with TNF- α inhibitor use were more likely to be diagnosed with drug-resistant disease than patients without reported TNF- α inhibitor use (13.5% vs 10.0%, $P < .001$).

Characteristics of TB patients who had TB-associated deaths are presented in [Table 2](#). After adjustment for confounders, TNF- α inhibitor use was not statistically associated with TB-related death (aOR, 1.46 [95% CI, .95–2.26]). To explore the potential moderating effect of site of TB disease in the relationship between TNF- α inhibitor use and death, an interaction term was estimated, but was not statistically significant and was excluded from the final multivariable model. A small sample size limited our ability to compare characteristics of patients with TB-associated death, based on TNF- α inhibitor use ([Table 3](#)).

DISCUSSION

This analysis indicates that over an 8-year period in the US, few TB cases were reported as recipients of TNF- α inhibitor medications, but patients receiving TNF- α inhibitors at diagnosis and had different clinical characteristics from those of the general population of TB patients, with no significantly increased odds of TB-related death among TB patients using TNF- α inhibitors.

We found that more patients receiving TNF- α inhibitors were diagnosed with disseminated TB disease involving both pulmonary and extrapulmonary sites of disease than patients not receiving TNF- α inhibitors. This finding is consistent with most published data [[12](#), [15](#), [19–23](#)], with the exception of a small ($N = 16$) US-based cohort study that found most were diagnosed with only pulmonary disease [[13](#)]. Our analysis also found that patients with pulmonary TB disease receiving TNF- α inhibitors were more often diagnosed with AFB sputum smear

Table 2. Univariate and Multivariate Analyses of Characteristics Associated With Tuberculosis-Associated Death Among Patients With Tuberculosis, United States, 2010–2017 (n = 69 730)^a

Characteristic	OR	(95% CI)	PValue	aOR	(95% CI)	PValue
TNF-α inhibitor therapy						
No	Ref	
Yes	1.54	(1.02–2.32)	.04	1.46	(.95–2.26)	.09
Age at diagnosis, y						
15–24	Ref	
25–44	2.56	(1.79–3.78)	<.01	2.26	(1.78–3.80)	<.01
45–64	7.83	(5.46–11.24)	<.01	6.23	(4.32–8.99)	<.01
≥ 65	22.18	(15.51–31.71)	<.01	16.14	(11.23–23.22)	<.01
Sex						
Female	Ref	
Male	1.28	(1.17–1.40)	<.01	1.11	(1.00–1.22)	.04
Race/ethnicity						
White	Ref	
Black	0.70	(.61–.80)	<.01	0.95	(.82–1.10)	.48
Asian	0.65	(.57–.74)	<.01	1.13	(.96–1.34)	.15
Hispanic	0.65	(.57–.74)	<.01	1.20	(1.03–1.41)	.02
Other ^b	1.27	(1.01–1.59)	.04	1.56	(1.23–1.99)	<.01
Place of birth						
US-born	Ref	
Non-US born	0.58	(.53–.63)	<.01	0.71	(.61–.82)	<.01
Diabetes						
Not reported	Ref	
Reported	1.89	(1.72–2.08)	<.01	1.10	(.99–1.22)	.08
End-stage renal disease						
Not reported	Ref	
Reported	5.81	(4.99–6.77)	<.01	3.55	(3.00–4.20)	<.01
HIV status						
Negative	Ref	
Positive	1.64	(1.41–1.92)	<.01	2.37	(1.97–2.85)	<.01
Excess alcohol use past year						
No	Ref	
Yes	1.54	(1.37–1.73)	<.01	1.47	(1.27–1.69)	<.01
Drug use past year^c						
No	Ref	
Yes	0.87	(.73–1.03)	.11	
Homelessness past year^d						
No	Ref	
Yes	1.25	(1.06–1.48)	<.01	0.92	(.76–1.11)	.38
Private outpatient provider						
No	Ref	
Yes	0.57	(.50–.64)	<.01	0.48	(.42–.54)	<.01
Resident of long-term care facility^e						
No	Ref	
Yes	6.73	(5.70–7.94)	<.01	3.35	(2.80–4.02)	<.01
Resident of corrections facility^f						
No	Ref	
Yes	0.25	(.16–.37)	<.01	0.42	(.27–.63)	<.01
Employment status						
Employed	Ref	
Not employed	3.90	(3.48–4.38)	<.01	1.99	(1.74–2.27)	<.01
Initial smear result, any^g						
Negative	Ref	
Positive	2.17	(1.97–2.39)	<.01	1.74	(1.57–2.93)	<.01
Not done	4.85	(3.61–6.50)	<.01	4.34	(3.14–5.99)	<.01
Site of TB disease						
Pulmonary only	Ref	

Table 2. Continued

Characteristic	OR	(95% CI)	PValue	aOR	(95% CI)	PValue
Extrapulmonary only	0.53	(.46–.61)	<.01	0.80	(.69–.93)	<.01
Both	1.91	(1.70–2.13)	<.01	1.98	(1.76–2.24)	<.01
Drug resistance^h						
No	Ref	
Yes	0.94	(.82–1.07)	.36	1.15	(1.00–1.32)	.05
Not done	0.17	(.14–.21)	<.01	0.25	(.20–.31)	<.01

Abbreviations: aOR, adjusted odds ratio; CI, confidence interval; HIV, human immunodeficiency virus; OR, odds ratio; TB, tuberculosis; TNF- α , tumor necrosis factor alpha; US, United States.

^aIncludes all TB cases age 15 years or older diagnosed in the US between 2010 and 2017 with confirmed death outcomes, excluding patients with missing information on outpatient provider type or smear result.

^bThis category includes persons of American Indian/Alaska Native and Native Hawaiian/Other Pacific Islander races, as well as persons of multiple races.

^cInjection or noninjection drug use within 12 months prior to TB diagnosis, not included in multivariate model due to insignificance in the univariate model.

^dHomeless within 12 months prior to TB diagnosis.

^eResident of long-term institution at time of diagnosis.

^fResident of correctional facility at time of diagnosis.

^gInitial smear result for any available specimen with reported acid-fast bacilli studies, including sputum, body fluid, or tissue.

^hPresence of resistance to isoniazid, rifampin, pyrazinamide, and/or ethambutol among culture-positive cases with drug susceptibility results; included in the multivariate model despite the fact it shows insignificance in the univariate model.

microscopy–negative disease. This could be the result of a pathophysiologic link to treatment with biologics, similar to the presentation of paucibacillary TB disease in immunosuppressed patients with HIV infection [24], who more often have extrapulmonary, smear-negative, and noncavitary disease. This finding could also be related to earlier diagnosis in patients with increased access to specialty private doctors, with TNF- α inhibitor treatment serving as a proxy for the ability to have continuity of medical care with a rheumatologist. It should be noted, however, that initial symptoms of TB disease can mimic an inflammatory disease that a treating clinician may empirically treat with local or systemic immunosuppression, leading to delayed TB diagnosis, and potentially a higher burden of disease outside the lung [12, 25, 26].

The finding of higher resistance to first-line drugs among TB patients treated with TNF- α inhibitors is interesting for clinical management considerations, but the cause of the drug resistance was uncertain, as this database for analysis did not include extensive patient history prior to TB diagnosis, such as prior LTBI treatment and adherence, or risk for contact with other patients with drug-resistant disease.

In our analysis, we did not find a significantly increased risk of TB-related mortality among TNF- α inhibitor users. Published studies comparing TB mortality in TNF- α inhibitor users vs nonusers are limited, and prospective studies are needed. In a single retrospective study of TB mortality by Beavers et al during 2005–2006, multivariable analysis demonstrated that receiving immunosuppressive medications was significantly associated with TB-related deaths; this analysis included patients receiving both prednisone and TNF- α inhibitors [14]. Similar to TB-related mortality associations in our analyses, the Beavers et al study found a TB-related mortality association with smear positivity, drug resistance, HIV, long-term care residence, and age. In a study of TB patient mortality from 2009 to 2013 by Hannah et al [27], TNF- α inhibitor use was combined with post-organ transplantation to

define “any immunosuppression,” which was significantly associated with an increased adjusted odds of TB-related mortality (aOR, 2.20 [95% CI, 1.71–2.83]), controlling for HIV, end-stage renal disease, multidrug-resistant TB, and increasing age. Finding similar variables significantly related to TB mortality as these other 2 studies lends confidence to our model estimates. It is worth noting that patients with reported TNF- α inhibitor use in our study varied significantly from nonusers in terms of key structural determinants of health such as race and employment, and that other unmeasured structural determinants may be acting as confounders.

The primary limitation of this analysis is its reliance on surveillance data and its possible underascertainment of TNF- α inhibitor use and TB-related mortality. Because TNF- α inhibitor therapy reported on the RVCT may underestimate or fail to capture the true number of patients who have recently received TNF- α inhibitors at the time of TB diagnosis, our analysis may not capture differences between groups and our results may be biased toward the null hypothesis. Our analyses did not capture the outcomes of death after therapy, slow response to therapy, or disability, nor are they able to describe whether patients had TNF- α inhibitors held during TB treatment, which could affect mortality. Although prior work suggests that different types of TNF- α inhibitors are associated with differential risk of TB disease [4, 28], our database did not include a population denominator of all TNF- α inhibitors in use, and also did not include data on type TNF- α inhibitor therapy or concurrent prednisone use. Prior studies have suggested a median duration of 12 weeks between start of infliximab treatment to onset of TB [12] with longer durations outside the US [29, 30]; however, our work was not able to capture duration of TNF- α inhibitor therapy prior to TB diagnosis. Future work should assess both infectious and noninfectious risks associated with individual drugs, as well as the contribution of specific inflammatory disorders, which may independently impact TB risk [6].

Table 3. Characteristics of Persons With Tuberculosis Who Were Dead at Diagnosis or Died During Treatment by Tumor Necrosis Factor Alpha Inhibitor Status, United States, 2010–2017

Characteristic	TNF- α Inhibitor (n = 24)		No TNF- α Inhibitor (n = 2310)		P Value ^a
	No.	(%)	No.	(%)	
Age at diagnosis, y					.41 ^b
15–24	1	(4.2)	33	(1.4)	
25–44	2	(8.3)	261	(11.3)	
45–64	6	(25.0)	739	(32.0)	
≥ 65	15	(62.5)	1277	(55.3)	
Sex					.02
Female	14	(58.3)	762	(33)	
Male	10	(41.7)	1548	(67)	
Race/ethnicity					.01 ^b
Asian	8	(33.3)	683	(29.6)	
Black	0	(0)	493	(21.4)	
Hispanic	6	(25.0)	584	(25.3)	
Other ^c	0	(0)	99	(4.3)	
White	10	(41.7)	441	(19.1)	
Place of birth					.63
Non-US born	15	(62.5)	1278	(55.3)	
US born	9	(37.5)	1024	(44.3)	
Diabetes					.33
No	20	(83.3)	1672	(72.4)	
Yes	4	(16.7)	638	(27.6)	
ESRD					.72 ^b
No	23	(95.8)	2089	(90.54)	
Yes	1	(4.2)	221	(9.6)	
HIV status					.40 ^b
Negative	14	(58.3)	1270	(55.0)	
Positive	0	(0)	177	(7.7)	
Homeless past year ^d					.41 ^b
No	24	(100)	2115	(91.6)	
Yes	0	(0)	161	(7.0)	
Private outpatient provider					<.01
No	15	(62.5)	1990	(86.1)	
Yes	9	(37.5)	316	(13.7)	
Employment status					1.00 ^b
Employed	4	(16.7)	371	(16.1)	
Not employed	19	(79.2)	1839	(79.6)	
Initial smear result, any ^e					1.00 ^b
Negative	7	(29.2)	642	(27.8)	
Not done	0	(0)	84	(3.6)	
Positive	17	(70.8)	1583	(68.5)	
Site of TB disease					<.01 ^b
Both	10	(41.7)	427	(18.5)	
Extrapulmonary only	4	(16.7)	247	(10.7)	
Pulmonary only	10	(41.7)	1636	(70.8)	
Drug resistance ^f					.41 ^b
No	20	(83.3)	1826	(79.0)	
Yes	2	(8.3)	255	(11.0)	
Not done	2	(8.3)	89	(3.9)	

Column totals for each variable may not sum to total due to missing data.

Abbreviations: ESRD, end-stage renal disease; HIV, human immunodeficiency virus; TB, tuberculosis; TNF- α , tumor necrosis factor alpha; US, United States.

^aP values calculated without missing values.

^bFisher exact test.

^cThis category includes persons of American Indian/Alaska Native and Native Hawaiian/Other Pacific Islander races, as well as persons of multiple races.

^dHomeless within 12 months prior to TB diagnosis.

^eInitial smear result for any available specimen with reported acid-fast bacilli studies, including sputum, body fluid, or tissue.

^fPresence of resistance to isoniazid, rifampin, pyrazinamide, and/or ethambutol among culture-positive cases with drug susceptibility results.

TNF- α inhibitors were initially approved for use in treatment of rheumatoid arthritis, but in the decades since their initial approval, their use has expanded dramatically and continues to grow. Inevitably, clinicians communicating with patients on these therapies will be faced with questions about TB risk, diagnosis, and clinical outcomes. TB prevention through testing for and treatment of latent TB infection is recommended for persons with immunocompromising diseases and before administration of immunosuppressive medications [31].

While this study did not demonstrate a statistically significant association between TNF- α inhibitor use and mortality, clinicians evaluating TNF- α inhibitor-treated patients with extrapulmonary manifestations should have a high index of suspicion for TB and consider additional tests before ruling out TB. The lack of a definitive diagnostic test for LTBI represents an urgent research gap that impacts many TB patients [32]; an improved diagnostic test for LTBI with predictive value for progression to TB disease would be particularly helpful for patients embarking on TNF- α inhibitor therapy. Additionally, as use of both TNF- α inhibitor and other biologic therapy increases, improved TB surveillance efforts that capture type and duration of immune-modulating therapy are urgently needed.

Supplementary Data

Supplementary materials are available at *Open Forum Infectious Diseases* online. Consisting of data provided by the authors to benefit the reader, the posted materials are not copyedited and are the sole responsibility of the authors, so questions or comments should be addressed to the corresponding author.

Notes

Acknowledgments. The authors gratefully acknowledge the patients who contributed data to this analysis, and the local tuberculosis (TB) programs that provided care and case management to the patients.

Disclaimer. The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention.

Financial support. There were no sources of funding for this analysis and manuscript.

Potential conflicts of interest. All authors: No reported conflicts of interest. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

References

- Harris J, Keane J. How tumour necrosis factor blockers interfere with tuberculosis immunity. *Clin Exp Immunol* **2010**; 161:1–9.
- Gardam MA, Keystone EC, Menzies R, et al. Anti-tumour necrosis factor agents and tuberculosis risk: mechanisms of action and clinical management. *Lancet Infect Dis* **2003**; 3:148–55.
- Algood HM, Lin PL, Flynn JL. Tumor necrosis factor and chemokine interactions in the formation and maintenance of granulomas in tuberculosis. *Clin Infect Dis* **2005**; 41(Suppl 3):S189–93.
- Dixon WG, Hyrich KL, Watson KD, et al. Drug-specific risk of tuberculosis in patients with rheumatoid arthritis treated with anti-TNF therapy: results from the British Society for Rheumatology Biologics Register (BSRBR). *Ann Rheum Dis* **2010**; 69:522–8.
- Wallis RS, Broder MS, Wong JY, Hanson ME, Beenhouwer DO. Granulomatous infectious diseases associated with tumor necrosis factor antagonists. *Clin Infect Dis* **2004**; 38:1261–5.

- Zhang Z, Fan W, Yang G, et al. Risk of tuberculosis in patients treated with TNF- α antagonists: a systematic review and meta-analysis of randomised controlled trials. *BMJ Open* **2017**; 7:e012567.
- Hanauer SB. Review article: safety of infliximab in clinical trials. *Aliment Pharmacol Ther* **1999**; 13(Suppl 4):16–22.
- Núñez Martínez O, Ripoll Noisieux C, Carneros Martín JA, González Lara V, Gregorio Marañoñ HG. Reactivation tuberculosis in a patient with anti-TNF- α treatment. *Am J Gastroenterol* **2001**; 96:1665–6.
- Cantini F, Niccoli L, Goletti D. Tuberculosis risk in patients treated with non-anti-tumor necrosis factor- α (TNF- α) targeted biologics and recently licensed TNF- α inhibitors: data from clinical trials and national registries. *J Rheumatol Suppl* **2014**; 91:56–64.
- Jo KW, Hong Y, Jung YJ, et al. Incidence of tuberculosis among anti-tumor necrosis factor users in patients with a previous history of tuberculosis. *Respir Med* **2013**; 107:1797–802.
- Muñoz L, Casas S, Juanola X, Bordas X, Martínez C, Santin M. Prevention of anti-tumor necrosis factor-associated tuberculosis: a 10-year longitudinal cohort study. *Clin Infect Dis* **2015**; 60:349–56.
- Keane J, Gershon S, Wise RP, et al. Tuberculosis associated with infliximab, a tumor necrosis factor alpha-neutralizing agent. *N Engl J Med* **2001**; 345:1098–104.
- Winthrop KL, Baxter R, Liu L, et al. Mycobacterial diseases and antitumor necrosis factor therapy in USA. *Ann Rheum Dis* **2013**; 72:37–42.
- Beavers SF, Pascopella L, Davidow AL, et al. Tuberculosis mortality in the United States: epidemiology and prevention opportunities. *Ann Am Thorac Soc* **2018**; 15:683–92.
- Mohan AK, Coté TR, Block JA, Manadan AM, Siegel JN, Braun MM. Tuberculosis following the use of etanercept, a tumor necrosis factor inhibitor. *Clin Infect Dis* **2004**; 39:295–9.
- Centers for Disease Control and Prevention. Reported tuberculosis in the United States, 2015. Atlanta, GA: CDC; **2016**.
- Rubin DB. Multiple imputation for nonresponse in surveys. New York: John Wiley and Sons; **1987**.
- R Core Team. R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; **2020**. <https://www.R-project.org/>.
- Centers for Disease Control and Prevention. Tuberculosis associated with blocking agents against tumor necrosis factor- α —California, 2002–2003. *MMWR Morb Mortal Wkly Rep* **2004**; 53:683–6.
- Saliu OY, Sofer C, Stein DS, Schwander SK, Wallis RS. Tumor-necrosis-factor blockers: differential effects on mycobacterial immunity. *J Infect Dis* **2006**; 194:486–92.
- Lee EH, Kang YA, Leem AY, et al. Active tuberculosis incidence and characteristics in patients treated with tumor necrosis factor antagonists according to latent tuberculosis infection. *Sci Rep* **2017**; 7:6473.
- Dimakou K, Papaioannides D, Latsi P, Katsimboula S, Korantzopoulos P, Orphanidou D. Disseminated tuberculosis complicating anti-TNF- α treatment. *Int J Clin Pract* **2004**; 58:1052–5.
- Meira L, Chaves C, Araújo D, et al. Predictors and outcomes of disseminated tuberculosis in an intermediate burden setting. *Pulmonology* **2019**; 25:320–7.
- Sterling TR, Pham PA, Chaisson RE. HIV infection-related tuberculosis: clinical manifestations and treatment. *Clin Infect Dis* **2010**; 50(Suppl 3):S223–30.
- Abadir AP, Han JY, Youssef FA. Intestinal tuberculosis masquerading as Crohn's disease? A case of disseminated tuberculosis after anti-TNF therapy for suspected Crohn's disease. *Case Rep Gastrointest Med* **2019**; 2019:6053503.
- Koschny R, Junghans T, Mischnik A, et al. Development of military tuberculosis under infliximab in a patient with spondyloarthritis and suspected Crohn's disease. *Z Gastroenterol* **2013**; 51:1177–83.
- Hannah HA, Miramontes R, Gandhi NR. Sociodemographic and clinical risk factors associated with tuberculosis mortality in the United States, 2009–2013. *Public Health Rep* **2017**; 132:366–75.
- Wang X, Wong SH, Wang XS, et al. Risk of tuberculosis in patients with immune-mediated diseases on biological therapies: a population-based study in a tuberculosis endemic region. *Rheumatology* **2019**; 58:803–10.
- Abitbol Y, Laharie D, Cosnes J, et al. Negative screening does not rule out the risk of tuberculosis in patients with inflammatory bowel disease undergoing anti-TNF treatment: a descriptive study on the GETAID cohort. *J Crohns Colitis* **2016**; 10:1179–85.
- Agarwal A, Kedia S, Jain S, et al. High risk of tuberculosis during infliximab therapy despite tuberculosis screening in inflammatory bowel disease patients in India. *Intest Res* **2018**; 16:588–98.
- Singh JA, Saag KG, Bridges SL Jr, et al. 2015 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. *Arthritis Rheumatol* **2016**; 68:1–26.
- World Health Organization. Evidence gaps and research needs identified during tuberculosis policy guideline development. WHO/CDS/TB/2018.27. Geneva, Switzerland: WHO; **2018**.