



EMORY
LIBRARIES &
INFORMATION
TECHNOLOGY

OpenEmory

Preventing Infection in Implant-based Breast Reconstruction: Evaluating the Evidence for Common Practices and Standardized Protocols

Nusaiba F. Baker, *Emory University School of Medicine*

Owen Brown, *Emory University School of Medicine*

[Alexandra Hart](#), *Emory University*

Dora Danko, *Emory University School of Medicine*

Christopher M. Stewart, *Emory University School of Medicine*

[Peter Thompson](#), *Emory University*

Journal Title: Plastic and Reconstructive Surgery - Global Open

Volume: Volume 10, Number 3

Publisher: Wolters Kluwer Health, Inc. | 2022-03-22, Pages E4208-E4208

Type of Work: Article | Final Publisher PDF

Publisher DOI: 10.1097/GOX.0000000000004208

Permanent URL: <https://pid.emory.edu/ark:/25593/vvnkw>

Final published version: <http://dx.doi.org/10.1097/GOX.0000000000004208>

Copyright information:

© 2022 The Authors.

This is an Open Access work distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (<https://creativecommons.org/licenses/by-nc-nd/4.0/rdf>).

Accessed December 3, 2023 2:36 AM EST

Preventing Infection in Implant-based Breast Reconstruction: Evaluating the Evidence for Common Practices and Standardized Protocols

Nusaiba F. Baker, PhD*
Owen Brown, MD†
Alexandra M. Hart, MD†
Dora Danko, BS*
Christopher M. Stewart, MD†
Peter W. Thompson, MD†

Background: Infection following implant-based breast reconstruction (IBBR) results in increased rates of hospital readmission, reoperation, patient and hospital expenses, and reconstructive failure. IBBR is a complex, multistep procedure, and there is a relative lack of high-quality plastic surgery evidence regarding “best practices” in the prevention of implant infections. In the absence of strong data, standardizing procedures based on available evidence can reduce error and improve efficacy and outcomes.

Methods: We performed a focused literature review of the available evidence supporting specific interventions for infection prevention in the preoperative, intraoperative, and postoperative phases of care that are applicable to IBBR. In addition, we examined previously published standardized perioperative protocols for implant reconstruction.

Results: Preoperative, intraoperative, and postoperative planning and organization is crucial in IBBR. Preoperative planning involves skin decolonization in advance of surgery with either chlorhexidine gluconate or mupirocin. Intraoperative methods that have shown potential benefit include double-gloving, breast pocket irrigation, separate closing instruments, and the utilization of “no-touch” techniques. In the postoperative period, the duration of drain removal and postoperative antibiotic administration play an important role in the prevention of surgical site infection.

Conclusions: There is a crucial need to establish an evidence-based set of “best practices” for IBBR, and there exists a paucity of evidence in the breast literature. These data can be utilized to develop a standardized protocol as part of a rigorous quality improvement methodology. (*Plast Reconstr Surg Glob Open* 2022;10:e4208; doi: 10.1097/GOX.0000000000004208; Published online 22 March 2022.)

INTRODUCTION

Implant-based breast reconstruction (IBBR) is the most common breast reconstruction technique.¹ A devastating complication of this technique is periprosthetic implant infection, with a reported incidence of 1%–30%.^{2–6} Implant infections are associated with wound healing, capsular contracture, suboptimal aesthetic outcome, and reconstructive failure, all potentially resulting in poor patient satisfaction.^{7,8} In addition, it has been

demonstrated that implant infections increase rates of hospital readmission, reoperation, and total cost of care.^{9,10}

Certain perioperative infection prevention measures, such as preoperative antibiotic prophylaxis and patient warming, are supported by robust evidence and are widely accepted as standard of care. Outside of these accepted best practices, surgeons who perform IBBR employ a variety of infection reduction measures that are often based largely on individual experience or anecdote. An American Society of Plastic Surgeons (ASPS) survey of 253 members found “considerable heterogeneity” in how surgeons apply infection reduction measures such as implant placement technique, pocket irrigation, and perioperative antibiotics.¹¹ Even the recent ASPS-published “Evidence-Based Clinical Practice Guideline: Breast Reconstruction with Expanders and Implants” discussed patient risk factor modification and the impact of adjuvant therapy on outcomes, but it did not provide guidance regarding specific perioperative practices that may help to reduce infection.¹² This is in large part due to the relative lack

From the *Emory University School of Medicine, Atlanta, Ga.; and †Emory Department of Surgery, Division of Plastic Surgery, Atlanta, Ga.

Received for publication October 15, 2021; accepted January 25, 2022.

Copyright © 2022 The Authors. Published by Wolters Kluwer Health, Inc. on behalf of The American Society of Plastic Surgeons. This is an open-access article distributed under the terms of the [Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 \(CCBY-NC-ND\)](#), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

DOI: 10.1097/GOX.0000000000004208

Disclosures: The authors have no financial interest to declare in relation to the content of this article.

of high-quality, plastic surgery-specific evidence regarding best practices in infection prevention specific to IBBR.

The purpose of this report is to critically review the evidence for the most common implant infection prevention measures used by plastic surgeons. We do not wish advocate that any specific technique is the optimal approach; rather, we seek to provide a synthesized and summarized reference for surgeons to help facilitate objective and evidence-based decisions regarding infection prevention practices.

METHODS

First, a literature review was performed to identify previously published protocols targeting implant infection, and the composition of these protocols was used to guide and narrow down the specific interventions discussed in this manuscript.^{5,13,14} The frequency with which specific interventions appeared in the three protocols is listed in Table 1.

Next, a comprehensive literature review using PubMed, Embase, Cochrane Library, and Web of Science databases in accordance with the PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-analyses) guidelines was performed for each specific intervention using the search terms highlighted in Table 2. Applicable literature from other specialties was included given the relative lack of plastic surgery-specific literature. Exclusion criteria included duplicate studies, non-English language studies, and studies that did not utilize patient data.

Once eligible results were identified, all publications were reviewed, and data were compiled for inclusion in the manuscript. These data were then reviewed by two independent reviewers within our Division of Plastic Surgery as well as the senior author. The final list of infection prevention measures discussed in this review were those used most often, with the highest amount of evidence basis, and the highest potential for efficacy and ease of implementation.

RESULTS

Preoperative Interventions

Skin Decolonization

Cutaneous bacteria are normal human microflora, with *Staphylococcal* and *Streptococcal* species being the most prevalent. Carriage rates for *S. aureus* are approximately 37.2%, and positive carrier status has been shown to be associated with a 7.1-fold increased relative risk of developing an infection following any type of surgery.²⁵ Likewise, *Staphylococcal* organisms and other Gram-positive skin flora are found to be the causative pathogen in most breast implant infections.²⁰

Preoperative skin decolonization protocols could potentially decrease the incidence of postoperative implant infection by reducing or eliminating the asymptomatic carriage. Twelve studies evaluated the efficacy of a chlorhexidine gluconate (CHG) or mupirocin regimen. Each of the prospective, nonrandomized studies demonstrated a significant improvement in infection

Takeaways

Question: What are the data for current practices employed by plastic surgeons regarding preoperative, intraoperative, and postoperative implant-based breast reconstruction?

Findings: We have compiled the data supporting several infection prevention interventions that are commonly employed in IBBR. Furthermore, we perform a comprehensive review of the available evidence supporting several specific interventions that have been included in these protocols.

Meaning: While data supporting specific interventions may not be particularly robust in study design or specificity to IBBR, it is possible that when bundled as components of a standard protocol, the benefit on infection reduction may be additive.

rates following implementation of the skin decolonization protocol.^{15–17,21–23,27} Of the randomized controlled trials (RCTs), two demonstrated a significant improvement in surgical site infections (SSIs) with use of the skin decolonization protocol,^{24,26} two demonstrated a trend towards fewer infections which was not statistically significant,^{18,19} and one demonstrated no difference.²⁸

A multicenter prospective study published in 2015 included over 14,000 screened *S. aureus* carriers undergoing either orthopedic joint arthroplasty or cardiac surgery.¹⁷ The rate of SSI following implementation of the protocol was significantly lower in both groups (relative risk [RR] 0.48 and 0.86, respectively). In an RCT published in the *New England Journal of Medicine*, 917 *S. aureus* carriers were randomized to receive 5 days of showers with CHG and intranasal mupirocin twice daily, or placebo.²⁹ The rate of *S. aureus* infection in the treatment group was significantly lower (RR 0.42), and the effect was most profound to deep SSIs (RR 0.21). Smith et al²⁶ evaluated preoperative skin decolonization versus placebo in 1350 *S. aureus* noncarriers undergoing Mohs surgery and demonstrated a 50% reduction in the rate of infection (2% versus 4%, $P = 0.03$). The effect of duration of decolonization duration was studied by Kline et al³⁰ in an RCT, demonstrating that a 5-day regimen of showers with CHG and intranasal mupirocin was more effective than 2 days of showers alone. In most published studies, a 5-day CHG/mupirocin regimen is considered the standard for skin decolonization.

The only plastic surgery-specific study was an RCT published by Veiga et al³¹ One hundred and fifty patients undergoing plastic surgery (including 16 breast reconstructions and 9 breast augmentations) were randomized to receive either no treatment, a shower with CHG preoperatively, or shower with placebo. There was no significant difference between any group in this study; however, the overall infection rate was very low (two infections total).

Summary: Though evidence specific to IBBR is lacking, there is relevant high-level evidence from other specialties that a 5-day skin decolonization protocol with CHG and mupirocin can decrease rate of SSI.

Table 1. Search Criteria for Literature Review

Search Terms	Identified Studies*	Included Studies†	Surgical Fields
Skin Decolonization			
1. "Preoperative skin decolonization" OR "mupirocin"	312	14	Orthopedics Cardiac Dermatologic General ENT Neurologic Plastic
2. "Chlorhexidine"			
Double gloving			
1. "Double gloving"	40	4	Obstetric General Plastic Vascular/trauma
2. "Glove perforation"			
3. "Surgical site infection" OR "Breast implant infection"			
Pocket irrigation			
1. "Breast"	239	13	Plastic
2. "Irrigation"			
3. "Infection"			
Minimal touch techniques			
1. "No touch" OR "Keller" OR "Minimal touch"	17	7	Cardiothoracic Hepatobiliary
2. "Breast reconstruction"			Orthopedics Plastic
Closing instruments			
1. "Closing instruments"	5	3	Colorectal
2. "Surgical site infections"			General
3. "Sterile"			
ADM			
1. "Acellular dermal matrix"	4	3	Plastic
2. "Surgical site infection"			
3. "Sterile"			
Postoperative antibiotics			
1. "Postoperative"	76	3	Plastic
2. "Antibiotics"			
3. "Surgical site infection"			
Drains			
1. "Drain care"	3	3	Plastic
2. "Breast reconstruction"			
Implant infection prevention protocols			
1. "Implant-based breast reconstruction"	18	3	Plastic
2. "Evidence based"			
3. "Protocol"			

ADM, acellular dermal matrix. "OR" refers to the Boolean operator used to focus search results.

*Studies were identified in accordance with the PRISMA guidelines using PubMed, Embase, Cochrane Library, and Web of Science databases. Eligible studies included RCTs, retrospective and prospective cohort studies, case-control, and cross-sectional studies.

†Exclusion criteria included duplicate studies, non-English language studies, and those not utilizing patient data.

Intraoperative Interventions

Double Gloving

Gloves in the operating room serves two purposes: as personal protective equipment for surgeons and as a barrier to contamination of the surgical field. Four studies compared the rate of innermost glove perforation in double versus single glove; each demonstrated a significantly higher rate of skin barrier failure with single gloving vs. double gloving.^{32-34,67} The largest of these was an RCT published by Laine and Aarnio⁶⁷ in which all surgeons across

all specialties at a single institution were randomized to wear either single or double gloves and found that the innermost glove was found to be perforated in only 6.8% of cases when the surgeons double gloved compared to 36.8% of cases when only a single glove was worn. In 2009, Misteli et al.³⁴ prospectively evaluated the rate of SSI following 4147 vascular and trauma procedures in cases with or without surgeon glove perforation. The authors found that there was a significantly higher rate of SSI in the setting of inner- or single glove perforation; however, multivariate logistic regression demonstrated that this effect was only statistically significant when preoperative surgical microbial prophylaxis was not administered. There were no studies specific to IBBR or plastic surgery which evaluated the practice of double gloving as it relates to infection.

Summary: Glove perforation with skin exposure, particularly when single gloving, may represent an occult source of surgical field contamination. There is evidence that double gloving reduces the rate of innermost glove perforation; however, there is no evidence definitively linking inner glove perforation to SSI.

Table 2. Incidence of Specific Measures Used in Previously Published Protocols

Preoperative	
Skin decolonization	3/3 (100%) ^{5,13,14}
Intraoperative	
Double gloving	1/3 (33%) ¹⁴
Pocket irrigation	3/3 (100%) ^{5,13,14}
"No-touch" techniques	3/3 (100%) ^{5,13,14}
Closing instruments	1/3 (33%) ¹⁴
Postoperative	
Postoperative antibiotics	2/3 (67%) ^{5,13}
Drains and drain care	3/3 (100%) ^{5,13,14}

Pocket Irrigation

In a 2019 ASPS survey, 63% of reconstructive surgeons reported using some form of pocket irrigant. Triple-antibiotic solution (TAS) (50,000 IU Bacitracin, 1g Ancef, and 80mg Gentamicin³⁶) was most common and was used by 41% of surgeon responders³⁷; 12.7% utilized betadine at varying concentrations, whereas others used Hibiclens (CHG; 0.9% of respondents), Irrisept (0.05% CHG in sterile water, 0.6% of respondents), Clorpectin wcs-90 (oxychlorosene sodium, 0.6% of respondents), and PhaseOne wound irrigation (hypochlorous acid; 0.3% of respondents).³⁷

The optimal irrigant has been evaluated in several studies. Merceron et al⁴³ performed a retrospective study that evaluated the effectiveness of CHG irrigation against TAS in IBBR. In this study, CHG proved to be superior to TAS in terms of infection reduction and rate of overall complications; however, there was no significant difference in rate of capsular contracture. Haynes et al compared CHG alone, TAS alone, and a combination of the two in IBBR.⁴⁴ The authors found that TAS combined with CHG resulted in significant protection against surgical complications over either irrigant alone. These data suggest a role for the inclusion of CHG irrigation compared to TAS alone. To our knowledge, there have not been any prospective or retrospective studies performed to validate the in vitro efficacy of PhaseOne or other hypochlorous acid derived irrigation techniques.

Summary: Pocket irrigation with some form of antibiotic solution has been shown to reduce the rate of infection and capsular contracture in some studies involving breast augmentation. Although this data can be extrapolated to apply to breast reconstruction, the optimal type and method of irrigation has yet to be determined.

“No-touch” Techniques

Another important source of surgical site contamination is that from native skin flora that may persist despite standard preoperative skin preparation. A variety of “no-touch” techniques have been developed, and range from self-retaining sterile retractor systems to the Keller Funnel (Allergan, Madison, N.J.) that has traditionally been used in augmentation mammoplasty.^{49,50} There is data to support the underlying theory that these techniques minimize possible cutaneous contamination of the sterile breast implant. Moyer et al⁵¹ compared the Keller Funnel versus the typical digital insertion method. In this study, the authors demonstrated that using the Keller Funnel resulted in a 27-fold decrease in implant-to-skin contact. Separately, the authors also swabbed the cadaver breasts with methicillin-sensitive *S. aureus* before implant placement and found that implant contamination as measured by cultures from the implant surface was twice as common using the digital insertion technique.

We identified one study specific to IBBR with tissue expanders (TEs) that not only demonstrated the benefit of a “no-touch” technique, but also directly evaluated its effect on reconstructive failure due to infection. Wilson described a technique using a transparent drape and a series of hooks to create a self-retaining retractor

system. This study found a statistically significant six-fold reduction in infection-related reconstruction failure from 11.5% to 1.9%.^{49,52}

Summary: There is evidence that the use of “no-touch” implant placement techniques can reduce implant-to-skin contamination during implant placement. There is also evidence that these techniques may translate to a reduction in rates of infection, including infection-related reconstructive failure in IBBR, though this data is relatively limited.

Closing Instruments

It is common during certain procedures to exchange surgical instruments for a new, sterile set of instruments before the wound is closed. This practice is particularly common in colorectal surgery, which has the highest rate of SSI in the literature (3%–30%). We identified two studies that investigated closing instruments in colorectal surgery before wound closure.^{53,54} However, neither study identified any benefit with regards to SSI. In contrast, a retrospective study in pancreatic surgery literature found a significant reduction in incisional SSI when using closing instruments and new drapes when compared to standard methods. The authors demonstrated a significant reduction in SSI from 12.4% in the control group versus to 2.2% in the clean group.⁵⁵ There is a lack of evidence regarding similar practice in cosmetic breast surgery or breast reconstruction.

Summary: The available literature is equivocal regarding the clinical benefit of using new, sterile instruments during skin closure. However, this has not yet been studied with regards to IBBR.

Postoperative Interventions

Postoperative Antibiotics

Preoperative antibiotics are standard of care in breast reconstruction. However, the decision to prescribe postoperative antibiotics is highly variable.^{56,57,68,69} A 2011 ASPS survey of providers performing IBBR demonstrated a lack of uniformity, with postoperative antibiotics prescribed by 72% of respondents.⁷⁰ Of those that prescribed antibiotics, 46% preferred to discontinue at time of drain removal, whereas 52% tended toward a specific postoperative day.

Five studies may help to inform decisions about antibiotic duration in IBBR.^{58–60,71,72} The largest retrospective report, published by McCullough et al⁵⁸ found that there was a slight trend towards more infections in the less than 24 hours group (RR 1.12), though this was not statistically significant. A noninferiority RCT by Phillips et al⁷³ involving patients undergoing IBBR with TEs compared oral antibiotic discontinuation at 24 hours postoperatively versus at the time of drain removal. A similar rate of infections was found between the two groups; however, patients in the extended group were more likely to require IV antibiotics and had a higher rate of TE loss (14.0% versus 4.8%).⁷³ The authors concluded that 24 hours of antibiotics was equivalent to extended oral antibiotic therapy in regards to reducing SSI.

Wang et al⁷⁴ corroborated the findings of Phillips et al via a systematic review of a total of 953 patients. The overall

risk of infection trended upward with less than 24 hour duration of antibiotic therapy compared to greater than 24 hours (19% and 14%, respectively, RR 1.3), though this result did not achieve statistical significance. Therefore, the authors concluded that prolonged antibiotic therapy was not effective in reducing SSI or implant loss.

Summary: There is significant heterogeneity regarding the decision to prescribe postoperative antibiotics. However, there is level II evidence suggesting there is no added benefit to continuing postoperative antibiotics longer than 24 hours postoperatively.

Drains and Drain Care

Closed suction drains play an important role in reducing dead space, thereby reducing the incidence of seroma in breast surgery.⁸ The practice of keeping drains in place for a prolonged period has been theorized to contribute to implant infections. We identified three studies relevant to drain duration and care.^{75–77} Hanna et al⁷⁵ investigated the association between time to drain removal and the subsequent incidence of infection in 323 patients undergoing IBBR with TEs. In multivariate analysis, drain use for longer than 21 days was independently associated with a 3.3-fold increased risk of infection. However, the authors recognize that this does not necessarily indicate causation: though patients with drains left in place longer appeared to have higher infection rates, prolonged high drain output may simply represent an early manifestation of infection rather than infection resulting from the drain itself.

Some surgeons believe that drains represent a potential communication between the sterile implant cavity and the outside world, which may allow for bacterial translocation into the implant cavity. In a retrospective, consecutive cohort study of 200 IBBRs using TEs, Murray et al investigated a protocol in which all reconstructions received mupirocin 2% cream to the drain sites compared to control. The authors found that this protocol led to a significant reduction in infections.⁷⁶ It should be noted that there were significant limitations of this study: not only was there significant sample size discrepancy between the two groups, but drains were removed nearly two weeks earlier in the mupirocin group (7.5 versus 20 days postoperatively).

Another proposed method to reduce risk of infection is the use of a Biopatch (Ethicon, Somerville, N.J.), a CHG-impregnated disk that is placed at drain exit sites. The idea to use Biopatch in IBBR stems from data demonstrating its efficacy in decreasing central line-associated bloodstream infections.⁶¹ However, a retrospective review performed by Weichman et al⁷⁷ failed to demonstrate a statistical reduction in overall infections when using Biopatch.

Summary: There is evidence that early drain removal may decrease risk of implant infections. While common practice to use local antimicrobials such as mupirocin or Biopatch to dress the drain insertion site, there is no strong evidence supporting their routine use.

DISCUSSION

In this review, we have systematically compiled the best available evidence regarding infection prevention

interventions that are commonly employed in IBBR. We provide clear and concise summaries of the evidence for each intervention, which can serve as quick references for providers when considering intervention implementation or when seeking to augment an already established set of practices. Though some studies were limited by sample size, design, or were published in other specialty literature, we feel that the data presented represents the “best available evidence” for each individual measure. As with any proposed change, the risks of implementing these interventions must be weighed against the potential benefit toward reducing the risk of infection. Some interventions discussed in this review could have unintended negative consequences; for example, the use of topical antibiotics could result in an allergic hypersensitivity reaction, or the use and management of closing instruments operative room staff may prolong operative time.

Some of the evidence regarding any one intervention in isolation may be equivocal, yet there remains a potential for standardized protocols to have a significant positive impact on patient outcomes in IBBR. The concept of “care bundles,” as advocated by the Institute of Healthcare Innovation (IHI), may demonstrate the effectiveness of standardization of processes based on the “best available evidence.” IHI defines a care bundle as a “structured way of improving the processes of care and patient outcomes: a small, straightforward set of evidence-based practices that, when performed collectively and reliably, have been proven to improve patient outcomes.⁶² Bundles have been proven to be effective in terms of reducing rates of central line-associated blood stream infection⁶³ and ventilator-associated pneumonia.^{63,64} Within a particular bundle, each individual component may have a small and potentially even controversial benefit, and evidence for this benefit may come from a variety of sources other than specialty-specific level-one data. However, the effect of the whole bundle may be greater than the sum of its individual parts, resulting in a synergistic impact on patient outcomes.

As a complex, multistep process with high potential for variability amongst individual providers, IBBR is an appealing target for standardization. There are several published “proof-of-concept” protocols which suggests that implementation of standard protocols in IBBR can significantly improve postoperative infection rates (Table 3). In the largest of these studies to our knowledge, Khansa et al¹³ compared 198 patients undergoing IBBR with tissue expanders using a new infection prevention protocol with a historical cohort of 305 patients. The authors found that patients exposed to the protocol were 55% less likely to develop a SSI after controlling for potential confounders (OR 0.45, $P = 0.022$).¹³ Dassoulas et al¹⁵ then formulated a protocol again for IBBR with subpectoral tissue expanders which was also independently associated with a decrease in infection risk (OR 0.244, $P = 0.021$). Most recently, Knight et al¹⁴ published a more inclusive protocol used in patients undergoing IBBR using both implant types placed in either the subpectoral or prepectoral plane. Despite this promising data in support of protocols in IBBR, more study is needed. There remains a need to demonstrate that such protocols can be broadly effective

Table 3. Description of Previously Published Protocols, Including Patient Population, Protocol Components, and Outcomes

Publication	Protocol Specifics			Outcomes
	Preoperative	Intraoperative	Postoperative	
Khansa et al ¹⁵	<ul style="list-style-type: none"> -Submuscular TE reconstruction only: before protocol implementation (2005–2010) versus after (2010–2012) 	<ul style="list-style-type: none"> -Preoperative IV antibiotics: cefazolin, or clindamycin if PCN allergy -Chlorhexidine skin prep, draping with plastic surgery team present -Soak TE in triple-antibiotic solution after opening (50,000 units of bacitracin, 1 g of cefazolin, and 80 mg of gentamicin in 500 mL of normal saline) -Pocket irrigation with antibiotic solution prior to implant placement -Reprepare around pocket with povidone-iodine and redrape with sterile towels -TE handled only by single surgeon after glove change -TE inserted with as little skin contact as possible - use of ADM (used serratus for inferior coverage) -Surgical drains always inserted -Layered skin closure -Preoperative IV antibiotics: cefazolin, or clindamycin if PCN allergy -Chlorhexidine skin prep -ADM soaked in Triple-antibiotic solution (cefazolin, gentamicin, bacitracin) -Pocket irrigation with antibiotic solution AND povidone-iodine prior to implant placement -ADM placed with as little skin contact as possible -Re-prep around pocket with chlorhexidine and Redrape with sterile towels -All surgeons required to change outer gloves -Irrigate pocket again with antibiotic solution AND povidone-iodine -Soak TE/implant in triple-antibiotic solution after opening -TE/implant inserted with as little skin contact as possible -Surgical drains always inserted -Layered skin closure with Dermabond -Preoperative IV antibiotics: teicoplanin and gentamicin -Intraoperative personnel reduction and avoid door opening (use of locks and signs) -Reduce operative time: two surgeons for bilateral procedures -Chlorhexidine skin prep -Nipple shields for unilateral cases -Surgeons require to double glove; must change outer glove prior to implant handling -Implant handled only by single surgeon after glove change -Pocket irrigated and implant soaked in Vancomycin 1 g and Gentamicin 160 mg solution with sterile water -Implants <500 cc only -Unused sterile instruments only after implant opened -Surgical drains always inserted -Bacteriostatic suture with skin glue 	<ul style="list-style-type: none"> -Perioperative antibiotics (cefazolin, or clindamycin if PCN allergy) for 24 hours -Discharge on PO antibiotics until final drain removal -Drain removal when output ≤ 30cc/day -Expansion started 2-4 weeks postoperatively (or rapid expansion prior to radiation therapy) 	<ul style="list-style-type: none"> -305 patients (456 total reconstructions) before protocol; 198 patients (313 total reconstructions) after protocol -Fewer patients experienced infections in protocol group (11.6% vs 18.4%, $P = 0.042$); fewer total infected TE in protocol patients (9.3% vs 13.2%, $P = 0.097$) -Protocol significantly reduced odds of infection on multivariate analysis (OR 0.45, $P = 0.022$) -235 patients (358 total reconstructions) before protocol; 85 patients (135 total reconstructions) after protocol -Reduced incidence of infection after protocol implementation (2.9% versus 9.5%, $p=0.013$) -Protocol independently associated with decrease infection risk (OR 0.244, $p=0.021$)
Dassoulas et al ⁶	<ul style="list-style-type: none"> -All immediate implant-based: before protocol implementation vs (2010–2014) after (implemented in 2015) -Excluded autologous reconstruction -Two breast surgeons; three plastic surgeons 	<ul style="list-style-type: none"> -Chlorhexidine scrub three days prior to surgery (specific instruction to pay attention to axilla, chest wall, IMF) -Intra-nasal mupirocin BID for 5 days before surgery 	<ul style="list-style-type: none"> -Discharge on PO antibiotics until final drain removal (if previous XRT, prescribed TMP-SMX double-strength BID; if not, Keflex) -Drain removal when output ≤ 30cc/day or by POD21 	<ul style="list-style-type: none"> -54 patients (77 total implant-based reconstructions) before protocol; 106 patients (129 total reconstructions) after protocol -Reduced rate of implant loss at three months after protocol implementation (14% vs 0%, $P < 0.00001$)
Knight et al ¹⁴	<ul style="list-style-type: none"> -All immediate implant-based before protocol (2012–2014) versus after (2015–2017) -Excluded LDF and two-stage reconstructions -Excluded patients with >1 risk factor (BMI > 30, smoker; DM, radiotherapy, neoadjuvant chemotherapy) -Four breast surgeons; seven plastic surgeons 	<ul style="list-style-type: none"> -Patient selection: only included those with ≤1 risk factor (BMI > 30, smoker, DM, radiotherapy, neoadjuvant chemotherapy) -MSSA/MRSA screening 48 h before surgery -No shaving or waxing for operation -Chlorhexidine shower the night before and morning of surgery 	<ul style="list-style-type: none"> -Oral doxycycline 100mg BD until final drain removal -Drain removal when output <30 cc/day on 2 consecutive days or by day 10 	<ul style="list-style-type: none"> -54 patients (77 total implant-based reconstructions) before protocol; 106 patients (129 total reconstructions) after protocol -Reduced rate of implant loss at three months after protocol implementation (14% vs 0%, $P < 0.00001$)

ADM, acellular dermal matrix; BID, bis in die; BMI, body mass index; DM, diabetes mellitus; IMF, inframammary fold; MRSA, Methicillin-resistant *Staphylococcus aureus*; MSSA, Methicillin-sensitive *Staphylococcus aureus*; PCN, penicillin; PO, per oral administration; TE, Tissue expander; TMP-SMX, trimethoprim/sulfamethoxazole; XRT, radiation therapy.

in IBBR regardless of prosthetic type, plane of placement, and patient comorbidities.

This review has several important limitations. One must remember that outcomes are closely related to the “quality” of the mastectomy skin flaps: when thin and relatively de-vascularized, tissue healing can be compromised which plays a critical role in infection development.^{65,66} However, not only is this factor largely out of the control of the plastic surgeon but it is also out of the intended scope of this review. Second, the data analyzed are likely subject to publication bias, whereby results which demonstrate a positive treatment effect are more likely to be published, resulting in an over-estimation of the potential benefit of a given intervention. In addition, the design and objectives of the available studies supporting implant infection prevention protocols may be subject to the Hawthorne effect. In this form of observation research bias, investigators and study participants may alter their behavior due to awareness of participation in an experiment, potentially affecting outcomes of interest. Longer-term follow-up after implementation of an infection prevention bundle can help determine if outcome improvement is due to the direct effect of the interventions, or simply a self-limited observation bias.

CONCLUSIONS

Implant-based breast reconstruction is a ubiquitous practice in plastic surgery, and implant infections can have significant adverse effects on patient outcome and costs of care. In this review, we scrutinized the “best available evidence” in support of several infection prevention interventions that may aid in provider decision-making. Although data supporting specific interventions may not be particularly robust in study design or specificity to IBBR, it is possible that when bundled as components of a standard protocol, the benefit on infection reduction may be additive. Further investigation of this approach within the framework of a rigorous quality improvement methodology is necessary.

Peter W. Thompson, MD, FACS

Emory Division of Plastic Surgery
1365 Clifton Rd NE
Clinic B, Rm 4016
Atlanta, GA 30322
E-mail: pwthomp@emory.edu

REFERENCES

- American Society of Plastic Surgeons. 2020 Plastic Surgery Statistics Report. American Society of Plastic Surgeons. Available at <https://www.plasticsurgery.org/documents/News/Statistics/2020/plastic-surgery-statistics-full-report-2020.pdf>. Accessed February 23, 2022.
- Gabriel SE, Woods JE, O'Fallon WM, et al. Complications leading to surgery after breast implantation. *N Engl J Med*. 1997;336:677–682.
- Brand KG. Infection of mammary prostheses: a survey and the question of prevention. *Ann Plast Surg*. 1993;30:289–295.
- Olsen MA, Nickel KB, Fox IK, et al. Incidence of surgical site infection following mastectomy with and without immediate reconstruction using private insurer claims data. *Infect Control Hosp Epidemiol*. 2015;36:907–914.
- Dassoulas KR, Wang J, Thuman J, et al. Reducing infection rates in implant-based breast reconstruction: impact of an evidence-based protocol. *Ann Plast Surg*. 2018;80:493–499.
- Wilkins EG, Hamill JB, Kim HM, et al. Complications in post-mastectomy breast reconstruction: one-year outcomes of the Mastectomy Reconstruction Outcomes Consortium (MROC) Study. *Ann Surg*. 2018;267:164–170.
- Fanouf N, Salem I, Tawilé C, et al. Absence of capsular contracture in 319 consecutive augmentation mammoplasties: dependent drains as a possible factor. *Can J Plast Surg*. 2004;12:193–197.
- Ooi ASH, Song DH. Reducing infection risk in implant-based breast-reconstruction surgery: challenges and solutions. *Breast Cancer (Dove Med Press)*. 2016;8:161–172.
- Badia JM, Casey AL, Petrosillo N, et al. Impact of surgical site infection on healthcare costs and patient outcomes: a systematic review in six European countries. *J Hosp Infect*. 2017;96:1–15.
- Yan C, Fischer JP, Wes AM, et al. The cost of major complications associated with immediate two-stage expander/implant-based breast reconstruction. *J Plast Surg Hand Surg*. 2015;49:166–171.
- Gowda AU, Chopra K, Brown EN, et al. Preventing breast implant contamination in breast reconstruction: a national survey of current practice. *Ann Plast Surg*. 2017;78:153–156.
- Evidence-Based Clinical Practice Guideline. Breast reconstruction with expanders and implants. American Society of Plastic Surgeons. Available at <https://www.plasticsurgery.org/for-medical-professionals/quality/evidence-based-clinical-practice-guidelines>. Accessed January 15, 2021.
- Khansa I, Hendrick RG Jr, Shore A, et al. Breast reconstruction with tissue expanders: implementation of a standardized best-practices protocol to reduce infection rates. *Plast Reconstr Surg*. 2014;134:11–18.
- Knight HJ, Musgrove JJ, Youssef MMG, et al. Significantly reducing implant loss rates in immediate implant-based breast reconstruction: a protocol and completed audit of quality assurance. *J Plast Reconstr Aesthet Surg*. 2020;73:1043–1049.
- Lefebvre J, Buffet-Bataillon S, Henaux PL, et al. *Staphylococcus aureus* screening and decolonization reduces the risk of surgical site infections in patients undergoing deep brain stimulation surgery. *J Hosp Infect*. 2017;95:144–147.
- Saraswat MK, Magruder JT, Crawford TC, et al. Preoperative *Staphylococcus aureus* screening and targeted decolonization in cardiac surgery. *Ann Thorac Surg*. 2017;104:1349–1356.
- Schweizer ML, Chiang HY, Septimus E, et al. Association of a bundled intervention with surgical site infections among patients undergoing cardiac, hip, or knee surgery. *JAMA*. 2015;313:2162–2171.
- Shuman AG, Shuman EK, Hauff SJ, et al. Preoperative topical antimicrobial decolonization in head and neck surgery. *Laryngoscope*. 2012;122:2454–2460.
- Sousa RJ, Barreira PM, Leite PT, et al. Preoperative *Staphylococcus aureus* screening/decolonization protocol before total joint arthroplasty—results of a small prospective randomized trial. *J Arthroplasty*. 2016;31:234–239.
- Cohen JB, Carroll C, Tenenbaum MM, et al. Breast implant-associated infections: the role of the National Surgical Quality Improvement Program and the local microbiome. *Plast Reconstr Surg*. 2015;136:921–929.
- George S, Leasure AR, McGovern A, et al. Reduction of postoperative infections through routine preoperative universal decolonization of advanced heart failure patients with chlorhexidine and mupirocin before left ventricular assist device implantation: a single-center observational study. *Dimens Crit Care Nurs*. 2020;39:312–320.
- Rao N, Cannella B, Crossett LS, et al. A preoperative decolonization protocol for *Staphylococcus aureus* prevents orthopaedic infections. *Clin Orthop Relat Res*. 2008;466:1343–1348.
- Sporer SM, Rogers T, Abella L. Methicillin-resistant and methicillin-sensitive *Staphylococcus aureus* screening and decolonization to

- reduce surgical site infection in elective total joint arthroplasty. *J Arthroplasty*. 2016;31(Suppl 9):144–147.
24. Cherian P, Gunson T, Borchard K, et al. Oral antibiotics versus topical decolonization to prevent surgical site infection after Mohs micrographic surgery—a randomized, controlled trial. *Dermatol Surg*. 2013;39:1486–1493.
 25. Kluytmans J, van Belkum A, Verbrugh H. Nasal carriage of *Staphylococcus aureus*: epidemiology, underlying mechanisms, and associated risks. *Clin Microbiol Rev*. 1997;10:505–520.
 26. Smith H, Borchard K, Cherian P, et al. Randomized controlled trial of preoperative topical decolonization to reduce surgical site infection for *Staphylococcus aureus* nasal swab-negative Mohs micrographic surgery patients. *Dermatol Surg*. 2019;45:229–233.
 27. Kohler P, Sommerstein R, Schönrrath F, et al. Effect of perioperative mupirocin and antiseptic body wash on infection rate and causative pathogens in patients undergoing cardiac surgery. *Am J Infect Control*. 2015;43:e33–e38.
 28. Rohrer F, Nötzli H, Risch L, et al. Does preoperative decolonization reduce surgical site infections in elective orthopaedic surgery? A prospective randomized controlled trial. *Clin Orthop Relat Res*. 2020;478:1790–1800.
 29. Bode LG, Kluytmans JA, Wertheim HF, et al. Preventing surgical-site infections in nasal carriers of *Staphylococcus aureus*. *N Engl J Med*. 2010;362:9–17.
 30. Kline SE, Neaton JD, Lynfield R, et al. Randomized controlled trial of a self-administered five-day antiseptic bundle versus usual disinfectant soap showers for preoperative eradication of *Staphylococcus aureus* colonization. *Infect Control Hosp Epidemiol*. 2018;39:1049–1057.
 31. Veiga DF, Damasceno CA, Veiga-Filho J, et al. Randomized controlled trial of the effectiveness of chlorhexidine showers before elective plastic surgical procedures. *Infect Control Hosp Epidemiol*. 2009;30:77–79.
 32. Kovavisarath E, Jaravechson S. Comparison of perforation between single and double-gloving in perineorrhaphy after vaginal delivery: a randomized controlled trial. *Aust N Z J Obstet Gynaecol*. 1998;38:58–60.
 33. Marín-Bertolín S, González-Martínez R, Giménez CN, et al. Does double gloving protect surgical staff from skin contamination during plastic surgery? *Plast Reconstr Surg*. 1997;99:956–960.
 34. Misteli H, Weber WP, Reck S, et al. Surgical glove perforation and the risk of surgical site infection. *Arch Surg*. 2009;144:553–558; discussion 558.
 35. Baker NF, Hart AM, Carlson GW, et al. A systematic review of breast irrigation in implant-based breast surgery. *Ann Plast Surg*. 2021;86:359–364.
 36. Adams WP Jr, Conner WC, Barton FE Jr, et al. Optimizing breast-pocket irrigation: the post-betadine era. *Plast Reconstr Surg*. 2001;107:1596–1601.
 37. Epps MT, Langsdon S, Pels TK, et al. Pocket irrigation and technique during reconstructive surgery: an American Society of Plastic Surgery Survey of Current Practice. *Ann Plast Surg*. 2019;82(6S Suppl 5):S427–S432.
 38. Araco A, Gravante G, Araco F, et al. Infections of breast implants in aesthetic breast augmentations: a single-center review of 3,002 patients. *Aesthetic Plast Surg*. 2007;31:325–329.
 39. Pfeiffer P, Jørgensen S, Kristiansen TB, et al. Protective effect of topical antibiotics in breast augmentation. *Plast Reconstr Surg*. 2009;124:629–634.
 40. Blount AL, Martin MD, Lineberry KD, et al. Capsular contracture rate in a low-risk population after primary augmentation mammoplasty. *Aesthet Surg J*. 2013;33:516–521.
 41. Lynch JM, Sebai ME, Rodriguez-Unda NA, et al. Breast pocket irrigation with antibiotic solution at implant insertion: a systematic review and meta-analysis. *Aesthetic Plast Surg*. 2018;42:1179–1186.
 42. Boustany AN, Elmaraghi S, Agochukwu N, et al. A breast prosthesis infection update: two-year incidence, risk factors and management at single institution. *Indian J Plast Surg*. 2018;51:7–14.
 43. Merceron TK, Betarbet U, Hart A, Baker N, Carlson GW, Losken A. Comparison of complications following implant-based breast reconstruction using triple antibiotic solution versus low concentration chlorhexidine gluconate solution. *Modern Plastic Surgery*. 2019;9:74–85.
 44. Haynes D. A Clinical Comparison of Irrisept®, Triple Antibiotic, and Combination Breast Pocket Irrigation. *Proceedings of the 61st Annual Scientific Meeting, Southeastern Society of Plastic and Reconstructive Surgeons*. West Palm Beach, FL. 2018 June 17-20.
 45. Haley CE, Marling-Cason M, Smith JW, et al. Bactericidal activity of antiseptics against methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol*. 1985;21:991–992.
 46. Septimus EJ, Schweizer ML. Decolonization in prevention of health care-associated infections. *Clin Microbiol Rev*. 2016;29:201–222.
 47. Wiener TC. Betadine and breast implants: an update. *Aesthet Surg J*. 2013;33:615–617.
 48. Jewell ML, Adams WP Jr. Betadine and breast implants. *Aesthet Surg J*. 2018;38:623–626.
 49. Wilson HB. Early results show reduced infection rate using no-touch technique for expander/ADM breast reconstruction. *Plast Reconstr Surg Glob Open*. 2015;3:e317.
 50. Newman AN, Davison SP. Effect of Keller funnel on the rate of capsular contracture in periareolar breast augmentation. *Plast Reconstr Surg Glob Open*. 2018;6:e1834.
 51. Moyer HR, Ghazi B, Saunders N, et al. Contamination in smooth gel breast implant placement: testing a funnel versus digital insertion technique in a cadaver model. *Aesthet Surg J*. 2012;32:194–199.
 52. Wilson HB. “No-touch” enhancement significantly reduces the risk of infection-related failure in immediate breast reconstruction. *Ann Plast Surg*. 2019;82(6S Suppl 5):S410–S416.
 53. Ortiz H, Armendariz P, Kreisler E, et al. Influence of rescrubbing before laparotomy closure on abdominal wound infection after colorectal cancer surgery: results of a multicenter randomized clinical trial. *Arch Surg*. 2012;147:614–620.
 54. Ghuman A, Chan T, Karimuddin AA, et al. Surgical site infection rates following implementation of a colorectal closure bundle in elective colorectal surgeries. *Dis Colon Rectum*. 2015;58:1078–1082.
 55. Hashimoto D, Chikamoto A, Arima K, et al. Unused sterile instruments for closure prevent wound surgical site infection after pancreatic surgery. *J Surg Res*. 2016;205:38–42.
 56. Phillips BT, Bishawi M, Dagum AB, Bui DT, Khan SU. A systematic review of infection rates and associated antibiotic duration in acellular dermal matrix breast reconstruction. *Eplasty*. 2014;14:e42.
 57. Sánchez JC. The impact of an entrepreneurship education program on entrepreneurial competencies and intention. *J Small Bus Manage*. 2013;51:447–465.
 58. McCullough MC, Chu CK, Duggal CS, et al. Antibiotic prophylaxis and resistance in surgical site infection after immediate tissue expander reconstruction of the breast. *Ann Plast Surg*. 2016;77:501–505.
 59. Phillips BT, Halvorson EG. Antibiotic prophylaxis following implant-based breast reconstruction: what is the evidence? *Plast Reconstr Surg*. 2016;138:751–757.
 60. Townley WA, Baluch N, Bagher S, et al. A single pre-operative antibiotic dose is as effective as continued antibiotic prophylaxis in implant-based breast reconstruction: a matched cohort study. *J Plast Reconstr Aesthet Surg*. 2015;68:673–678.

61. Levy I, Katz J, Solter E, et al. Chlorhexidine-impregnated dressing for prevention of colonization of central venous catheters in infants and children: a randomized controlled study. *Pediatr Infect Dis J*. 2005;24:676–679.
62. Institute for Healthcare Improvement: What is a bundle? Institute for Healthcare Improvement. Available at <http://www.ihl.org/resources/Pages/ImprovementStories/WhatsaBundle.aspx>. Accessed April 25, 2021.
63. O'Neil C, Ball K, Wood H, et al. A central line care maintenance bundle for the prevention of central line-associated bloodstream infection in non-intensive care unit settings. *Infect Control Hosp Epidemiol*. 2016;37:692–698.
64. Wip C, Napolitano L. Bundles to prevent ventilator-associated pneumonia: how valuable are they? *Curr Opin Infect Dis*. 2009;22:159–166.
65. Long C, Sue GR, Chattopadhyay A, et al. Critical evaluation of risk factors of infection following 2-stage implant-based breast reconstruction. *Plast Reconstr Surg Glob Open*. 2017;5:e1386.
66. Reish RG, Damjanovic B, Austen WG Jr, et al. Infection following implant-based reconstruction in 1952 consecutive breast reconstructions: salvage rates and predictors of success. *Plast Reconstr Surg*. 2013;131:1223–1230.
67. Laine T, Aarnio P. How often does glove perforation occur in surgery? Comparison between single gloves and a double-gloving system. *Am J Surg*. 2001;181:564–566.
68. Phillips BT, Bishawi M, Dagum AB, et al. A systematic review of antibiotic use and infection in breast reconstruction: what is the evidence? *Plast Reconstr Surg*. 2013;131:1–13.
69. Pannucci CJ, Antony AK, Wilkins EG. The impact of acellular dermal matrix on tissue expander/implant loss in breast reconstruction: an analysis of the tracking outcomes and operations in plastic surgery database. *Plast Reconstr Surg*. 2013;132:1–10.
70. Phillips BT, Wang ED, Mirrer J, et al. Current practice among plastic surgeons of antibiotic prophylaxis and closed-suction drains in breast reconstruction: experience, evidence, and implications for postoperative care. *Ann Plast Surg*. 2011;66:460–465.
71. Avashia YJ, Mohan R, Berhane C, et al. Postoperative antibiotic prophylaxis for implant-based breast reconstruction with acellular dermal matrix. *Plast Reconstr Surg*. 2013;131:453–461.
72. Clayton JL, Bazakas A, Lee CN, et al. Once is not enough: withholding postoperative prophylactic antibiotics in prosthetic breast reconstruction is associated with an increased risk of infection. *Plast Reconstr Surg*. 2012;130:495–502.
73. Phillips BT, Fourman MS, Bishawi M, et al. Are prophylactic postoperative antibiotics necessary for immediate breast reconstruction? results of a prospective randomized clinical trial. *J Am Coll Surg*. 2016;222:1116–1124.
74. Wang F, Chin R, Piper M, et al. Do prolonged prophylactic antibiotics reduce the incidence of surgical-site infections in immediate prosthetic breast reconstruction? *Plast Reconstr Surg*. 2016;138:1141–1149.
75. Hanna KR, Tilt A, Holland M, et al. Reducing infectious complications in implant based breast reconstruction: impact of early expansion and prolonged drain use. *Ann Plast Surg*. 2016;76(Suppl 4):S312–S315.
76. Murray JD, Elwood ET, Jones GE, et al. Decreasing expander breast infection: a new drain care protocol. *Can J Plast Surg*. 2009;17:17–21.
77. Weichman KE, Clavin NW, Miller HC, et al. Does the use of bio-patch devices at drain sites reduce perioperative infectious complications in patients undergoing immediate tissue expander breast reconstruction? *Plast Reconstr Surg*. 2015;135:9e–17e.