



## **Alarming downtrend in mechanical thrombectomy rates in African American patients during the COVID-19 pandemic-Insights from STAR**

[Jonathan Grossberg](#), *Emory University*

[Brian Howard](#), *Emory University*

S Al Kasab, *Medical University of South Carolina*

E Almallouhi, *Medical University of South Carolina*

A Alawieh, *Emory University*

P Jabbour, *Thomas Jefferson Univ Hosp*

A Sweid, *Thomas Jefferson Univ Hosp*

RM Starke, *University of Miami*

V Saini, *University of Miami*

SQ Wolfe, *Wake Forest Sch Med*

*Only first 10 authors above; see publication for full author list.*

---

**Journal Title:** JOURNAL OF NEUROINTERVENTIONAL SURGERY

**Volume:** Volume 13, Number 4

**Publisher:** BMJ PUBLISHING GROUP | 2021-04-01, Pages 304-307

**Type of Work:** Article

**Publisher DOI:** 10.1136/neurintsurg-2020-016946

**Permanent URL:** <https://pid.emory.edu/ark:/25593/vw1zm>

---

Final published version: <http://dx.doi.org/10.1136/neurintsurg-2020-016946>

*Accessed November 27, 2022 11:35 AM EST*

Original research

# Alarming downtrend in mechanical thrombectomy rates in African American patients during the COVID-19 pandemic-Insights from STAR

Sami Al Kasab <sup>1,2</sup>, Eyad Almallouhi,<sup>1,2</sup> Ali Alawieh <sup>3,4</sup>, Pascal Jabbour <sup>5</sup>, Ahmad Sweid,<sup>5</sup> Robert M Starke,<sup>6</sup> Vasu Saini <sup>6</sup>, Stacey Q Wolfe <sup>7</sup>, Kyle M Fargen,<sup>8</sup> Adam S Arthur <sup>9,10</sup>, Nitin Goyal,<sup>10</sup> Abhi Pandhi,<sup>11</sup> Ilko Maier <sup>12</sup>, Jonathan A Grossberg <sup>13</sup>, Brian M Howard <sup>14,15</sup>, Stavropoula I Tjoumakaris <sup>5</sup>, Ansaar Rai,<sup>16</sup> Min S Park,<sup>17</sup> Justin R Mascitelli,<sup>18</sup> Marios N Psychogios,<sup>19</sup> Alejandro M Spiotta<sup>1</sup>

► Additional material is published online only. To view, please visit the journal online (<http://dx.doi.org/10.1136/neurintsurg-2020-016946>).

For numbered affiliations see end of article.

## Correspondence to

Dr Sami Al Kasab, Neurosurgery, Medical University of South Carolina, Charleston, SC 29425, USA; [alkasab@musc.edu](mailto:alkasab@musc.edu)

SAK and EA are joint first authors.

Received 6 October 2020  
Revised 11 December 2020  
Accepted 12 December 2020

## ABSTRACT

**Background** The coronavirus disease (COVID-19) pandemic has affected stroke care globally. In this study, we aim to evaluate the impact of the current pandemic on racial disparities among stroke patients receiving mechanical thrombectomy (MT).

**Methods** We used the prospectively collected data in the Stroke Thrombectomy and Aneurysm Registry from 12 thrombectomy-capable stroke centers in the US and Europe. We included acute stroke patients who underwent MT between January 2017 and May 2020. We compared baseline features, vascular risk factors, location of occlusion, procedural metrics, complications, and discharge outcomes between patients presenting before (before February 2020) and those who presented during the pandemic (February to May 2020).

**Results** We identified 2083 stroke patients: of those 235 (11.3%) underwent MT during the COVID-19 pandemic. Compared with pre-pandemic, stroke patients who received MT during the pandemic had longer procedure duration (44 vs 38 min,  $P=0.006$ ), longer length of hospitalization (6 vs 4 days,  $P<0.001$ ), and higher in-hospital mortality (18.7% vs 11%,  $P<0.001$ ). Importantly, there was a lower number of African American patients undergoing MT during the COVID-19 pandemic (609 (32.9%) vs 56 (23.8%);  $P=0.004$ ).

**Conclusion** The COVID-19 pandemic has affected the care process for stroke patients receiving MT globally. There is a significant decline in the number of African American patients receiving MT, which mandates further investigation.

Evidence from recently published studies suggested that this decline disproportionately affects African American (AA) patients, widening the already existed racial disparity in stroke care.<sup>6</sup> In this study, we aimed to investigate the impact of the COVID-19 pandemic on the racial disparities related to MT.

## METHODS

### Patient population and collected variables

Prospectively collected data from 12 thrombectomy-capable stroke centers included in the Stroke Thrombectomy and Aneurysm Registry (STAR) collaboration was interrogated to identify stroke patients who received mechanical thrombectomy (MT) between January 2017 and May 2020. Eleven of the included centers are located in the United States and one in Germany. Patients who presented before February 2020 were considered pre-COVID-19 pandemic and patients who presented after February 2020 were considered during the COVID-19 pandemic. We compared baseline demographics, procedural metrics, and outcomes between patients undergoing MT before and during the pandemic. Approval from the Institutional Review Board at the Medical University of South Carolina was obtained, and no consent was needed per the institutional policy.

### Statistical analysis

We used univariate analysis to report patient demographic and clinical characteristics using median and IQR for continuous variables and percentages for categorical variables. Characteristics of groups were compared using the Mann–Whitney  $U$  test, and chi-square as appropriate. To assess for the relationship between the rate of AA (or Black Europeans of African ancestry) patients receiving MT and COVID-19 pandemic controlling for the center of presentation, we used the Cochran Mantel Haenszel test. An alpha level of 0.05 was used as the level of statistical significance. The analysis was conducted using SPSS v25 (IBM Corporation, New York, NY).

## INTRODUCTION

The Coronavirus disease of 2019 (COVID-19) pandemic has affected all aspects of medical care for various conditions.<sup>1</sup> Several studies reported a potential relationship between COVID-19 infection and ischemic stroke secondary to large vessel occlusion (LVO).<sup>2,3</sup> Importantly, recent studies reported a significant decline in the number of patients presenting with acute stroke, raising concerns that stroke patients are avoiding hospitals for various reasons including fear of COVID-19 infection.<sup>4,5</sup>



© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Al Kasab S, Almallouhi E, Alawieh A, et al. *J NeuroIntervent Surg* Epub ahead of print: [please include Day Month Year]. doi:10.1136/neurintsurg-2020-016946

**Table 1** Characteristics, procedural metrics, and outcomes of stroke patients receiving mechanical thrombectomy before and during the COVID-19 pandemic

	Thrombectomy patients during COVID-19 pandemic (n=235)	Thrombectomy patients before COVID-19 pandemic (n=1848)	P- value*
Age, median (IQR)	71 (59–79)	69 (58–79)	0.058
Females, n (%)	113 (48.1%)	901 (48.8%)	0.846
Racial/ethnic distribution:			<b>0.01</b>
White, n (%)	146 (62.1%)	1047 (56.7%)	
African American, n (%)	56 (23.8%)	609 (32.9%)	
Hispanic, n (%)	31 (13.2%)	188 (10.2%)	
Others, n (%)	2 (0.9%)	4 (0.2%)	
Admission NIHSS, median (IQR)	16 (10–21)	16 (10–21)	0.898
IV-tPA, n (%)	106 (45.1%)	831 (45%)	0.968
ASPECTS, median (IQR)†	9 (7–10)	9 (7–10)	0.225
Posterior circulation occlusion, n (%)	21 (8.9%)	195 (10.6%)	0.444
Symptom onset to groin in minutes, median (IQR)	343 (202–576)	288 (178–575)	0.065
Number of passes, median (IQR)	2 (1–3)	2 (1–3)	0.921
mTICI ≥2 b, n (%)	196 (83.4%)	1588 (85.9%)	0.228
Procedure duration in minutes, median (IQR)	44 (25–73)	38 (21–63)	<b>0.006</b>
Procedure complications, n (%)	20 (8.5%)	163 (8.8%)	0.874
siCH, n (%)	15 (6.4%)	108 (5.8%)	0.741
Length of hospitalization in days, median (IQR)	6 (3–10)	4 (2–5)	<b>&lt;0.001</b>
DC mRS, median (IQR)	4 (2–5)	3 (2–4)	<b>0.015</b>
mRS 0–2 on discharge, n (%)	77 (32.8%)	637 (34.5%)	0.604
In-hospital mortality, n (%)	44 (18.7%)	203 (11%)	<b>&lt;0.001</b>

\*Calculated using chi-square test for categorical variables and Mann–Whitney U test for the continuous variables.  
 †Only for patients with anterior circulation occlusion.  
 ASPECTS, Alberta Stroke Program Early CT score; COVID-19, coronavirus disease of 2019; IQR, interquartile range; IV-tPA, intravenous tissue plasminogen activator; LKN, last known normal; mRS, modified Rankin Scale; mTICI, modified Thrombolysis in Cerebral Infarction; NIHSS, National Institute of Health Stroke Scale; siCH, symptomatic intracerebral hematoma.


**RESULTS**

**Patients' characteristics**

A total of 2083 stroke patients were included in this study, 1848 (88.7%) underwent MT before the COVID-19 pandemic and 235 (11.3%) during the COVID-19 pandemic. The median age was 69 (IQR 58–79) years, admission National Institute of Health Stroke Scale (NIHSS) 16 (10-21), and Alberta Stroke Program Early CT Score (ASPECTS) 9 (IQR 7–10). Online supplemental table I summarizes baseline features and clinical outcomes. The racial/ethnic distribution of the included patients was as follows: 1197 (57.8%) white, 665 (31.9%) AA, 215 (10.3%) Hispanic, and six (0.3%) others.

**Before vs. during the COVID-19 pandemic**

Table 1 summarizes the comparison between stroke patients who underwent MT before vs during the COVID-19 pandemic. Five (2.1%) of the MT patients during the pandemic were positive for COVID-19 polymerase chain reaction (PCR) in nasopharyngeal swab. Significant differences in patients undergoing MT during the COVID-19 pandemic include: longer procedure duration (44 vs 38 min, P=0.006), longer length of hospitalization (6 vs 4, P<0.001), lower discharge mRS (4 vs 3, P=0.015), and higher in-hospital mortality (18.7% vs 11%, P<0.001), compared with patients who underwent MT before the pandemic. While the difference between the symptom onset to groin time was not significant in the overall population (P=0.065) and in non-AA patients (P=0.278), AA patients had longer onset to groin time during the COVID-19 pandemic (362 vs 275 min, P=0.047). In addition, the percentage of AA patients who underwent MT in the included centers decreased from 609/1848 (32.9%) before the COVID-19 pandemic to 56/235 (23.8%) during the COVID-19 pandemic (P=0.004). No interaction was noted between the percentage of AA patients receiving thrombectomy and the centers that performed the thrombectomy before and after the COVID-19 pandemic (P=0.192). No other statistically significant differences were noted in other races/ethnicities. Figure 1 shows the trend in the rate of AA patients receiving mechanical thrombectomy during the study period. Online supplemental tables II and III show the comparison between AA and non-AA thrombectomy patients before and during the COVID-19 pandemic.



**Figure 1** The trend in the rate of African American patients receiving mechanical thrombectomy during the study period.

## DISCUSSION

In this multicenter, international study, we evaluated the procedural metrics and outcomes of MT patients presenting during the COVID-19 pandemic compared with patients presenting during the 3 prior years. We have found that patients who presented during the pandemic required a longer procedure time and had a higher rate of mortality. While only 2.1% of these patients were positive for COVID-19 PCR, these findings likely reflect the effect of COVID-19 precautions and the changes in health-care workflow protocols.

Additional significant findings from our analyses included a reduction in the number of AA patients undergoing MT, which is highly alarming because racial disparities in cerebrovascular diseases care already exist. AA patients have higher vascular risk factors, and experience higher stroke-related in-hospital mortality.<sup>7</sup> Also, studies have shown that AA/Hispanic patients are more likely to have received aneurysm treatment following a subarachnoid hemorrhage rather than getting treatment in the early stages for unruptured aneurysm.<sup>8</sup>

The reduction in the number of AA patients receiving MT during the COVID-19 pandemic threatens to widen the gap in racial disparities given that most patients with emergent LVO would suffer from significant long-term disability if they do not receive treatment.<sup>9–11</sup> A recent study evaluating racial disparities among patients evaluated over a large telestroke network reported a significant drop among AA patients with acute ischemic stroke.<sup>6</sup> The overall drop in the number of AA patients presenting with acute stroke is a likely reason that there is a drop in the number of AA patients receiving MT. It remains unclear the reasons behind this drop. However, studies have shown that AA patients have a higher incidence of COVID-19 infection and increased mortality<sup>12,13</sup> which could be the reason why AA patients are reluctant to seek medical attention over concerns for safety due to fear of acquiring COVID-19.

Interestingly, there was an increase in the symptom onset to groin time in AA, but not non-AA patients which may also reflect the abundance of caution from presenting to hospitals during the early COVID-19 pandemic. A similar delay in care in all stroke patients during the COVID-19 pandemic was reported recently by Schirmer et al.<sup>14</sup> One of the proposed causes for this delay is the changes in stroke triage protocols during the pandemic.<sup>15</sup>

Our study has a few limitations. First, our study is limited by its observational nature. Also, our study does not provide information related to the relationship between COVID-19 infection and emergent LVO. However, our study provides multicentric, early observations stroke centers in different regions.

## CONCLUSION

Stroke patients receiving MT during the COVID-19 pandemic had a longer procedure duration and a higher mortality rate compared with patients presenting before the pandemic. Also, there is a decline in the number of AA patients receiving MT during the pandemic, which demonstrates the need for more work in the public health domain to educate patients with the importance of prompt medical evaluation for all those suffering from stroke symptoms.

### Author affiliations

<sup>1</sup>Neurosurgery, Medical University of South Carolina, Charleston, SC, USA

<sup>2</sup>Neurology, Medical University of South Carolina, Charleston, SC, USA

<sup>3</sup>Neurosurgery, Emory University, Atlanta, GA, USA

<sup>4</sup>Microbiology and Immunology, Medical University of South Carolina, Charleston, SC, USA

<sup>5</sup>Neurological Surgery, Thomas Jefferson University Hospital, Philadelphia, PA, USA

<sup>6</sup>Neurological Surgery, University of Miami Miller School of Medicine, Miami, FL, USA

<sup>7</sup>Neurosurgery, Wake Forest School of Medicine, Winston-Salem, NC, USA

<sup>8</sup>Neurosurgery, Wake Forest University, Winston-Salem, NC, USA

<sup>9</sup>Neurosurgery, Semmes-Murphey Neurologic and Spine Institute, Memphis, Tennessee, USA

<sup>10</sup>Neurosurgery, University of Tennessee Health Science Center, Memphis, Tennessee, USA

<sup>11</sup>Neurology, University of Tennessee Health Science Center College of Medicine, Memphis, Tennessee, USA

<sup>12</sup>Neurology, University Medicine Goettingen, Goettingen, NS, Germany

<sup>13</sup>Neurosurgery and Radiology, Emory University School of Medicine, Atlanta, Georgia, USA

<sup>14</sup>Neurosurgery, Emory University School of Medicine, Atlanta, Georgia, USA

<sup>15</sup>Radiology and Imaging Sciences, Emory University School of Medicine, Atlanta, Georgia, USA

<sup>16</sup>Radiology, West Virginia University Hospitals, Morgantown, West Virginia, USA

<sup>17</sup>Neurosurgery, Barrow Neurological Institute, Phoenix, Arizona, USA

<sup>18</sup>Neurosurgery, University of Texas Health Science Center at San Antonio, San Antonio, Texas, USA

<sup>19</sup>Department of Neuroradiology, Clinic of Radiology and Nuclear Medicine, University Hospital Basel, Basel, Switzerland

**Twitter** Sami Al Kasab @samialkasab, Pascal Jabbour @PascalJabbourMD, Ahmad Sweid @AhmadSweidMD, Robert M Starke @Starke\_neurosurgery, Adam S Arthur @AdamArthurMD, Abhi Pandhi @abhipandhi, Brian M Howard @BrianHoward\_MD and Alejandro M Spiotta @alex\_spiotta

**Collaborators** STAR collaborators: Dileep R Yavagal, MD; Eric C Peterson, MD; Alex Brehm, MD; Patrick A Brown, MD; M. Reid Gooch, MD; Nabeel Herial, MD; Dr. med. Jan Liman; Daniel Alan Hoit MD MPH; Violiza Inoa-Acosta MD; Christopher Nickele MD; Lucas Elijovich MD; Michael Cawley, MD; Gustavo Pradilla, MD

**Contributors** All authors have: provided a substantial contribution to the conception and design of the studies and/or the acquisition and/or the analysis of the data and/or the interpretation of the data. They have drafted the work or revised it for significant intellectual content and approved the final version of the manuscript. They agree to be accountable for all aspects of the work, including its accuracy and integrity.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial, or not-for-profit sectors.

**Competing interests** PJ: consultant-Medtronic, Microvention. RMS: consulting and teaching agreements with Penumbra, Abbott, Medtronic, InNeuroCo, and Cerenovus. ASA: consultant—Balt, Johnson and Johnson, Leica, Medtronic, Microvention, Penumbra, Scientia, Siemens, and Stryker; research support—Cerenovus, Microvention, Penumbra, and Siemens; and shareholder—Bendit, Cerebrotech, Endostream, Magneto, Marblehead, Neurogami, Serenity, Synchron, Triad Medical, Vascular Simulations. AR: consulting agreement with Stryker, Cerenovus, and Microvention. MNP: travel grants/honoraria—Phenox, Stryker, Siemens. AMS: consultant—Penumbra, Microvention, and Pulsar Vascular; travel grants/honoraria—Penumbra, Pulsar Vascular, Microvention, Stryker.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request. Data are available upon reasonable request.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

This article is made freely available for use in accordance with BMJ's website terms and conditions for the duration of the covid-19 pandemic or until otherwise determined by BMJ. You may use, download and print the article for any lawful, non-commercial purpose (including text and data mining) provided that all copyright notices and trade marks are retained.

### ORCID iDs

Sami Al Kasab <http://orcid.org/0000-0001-8909-9761>

Ali Alawieh <http://orcid.org/0000-0003-2601-8850>

Pascal Jabbour <http://orcid.org/0000-0002-1544-4910>

Vasu Saini <http://orcid.org/0000-0002-6796-5881>

Stacey Q Wolfe <http://orcid.org/0000-0001-7603-2728>

Adam S Arthur <http://orcid.org/0000-0002-1536-1613>

Ilko Maier <http://orcid.org/0000-0001-6988-8878>

Jonathan A Grossberg <http://orcid.org/0000-0002-1152-8826>

Brian M Howard <http://orcid.org/0000-0001-9134-0817>

Stavropoula I Tjoumakaris <http://orcid.org/0000-0002-1054-9414>

## REFERENCES

- 1 Blumenthal D, Fowler EJ, Abrams M, *et al.* Covid-19 – implications for the health care system. *N Engl J Med* 2020;383:1483–8.
- 2 Majidi S, Fifi JT, Ladner TR, *et al.* Emergent large vessel occlusion stroke during New York City's COVID-19 outbreak: clinical characteristics and paraclinical findings. *Stroke* 2020;51:2656–63.
- 3 Oxley TJ, Mocco J, Majidi S, *et al.* Large-vessel stroke as a presenting feature of Covid-19 in the young. *N Engl J Med* 2020;382:e60.
- 4 Jasne AS, Chojecka P, Maran I, *et al.* Stroke code presentations, interventions, and outcomes before and during the COVID-19 pandemic. *Stroke* 2020;51:2664–73.
- 5 Nguyen-Huynh MN, Tang XN, Vinson DR, *et al.* Acute stroke presentation, care, and outcomes in community hospitals in northern California during the COVID-19 pandemic. *Stroke* 2020;51:2918–24.
- 6 Cummings C, Almallouhi E, Al Kasab S, *et al.* Blacks are less likely to present with strokes during the COVID-19 pandemic: observations from the buckle of the stroke belt. *Stroke* 2020;51:3107–11.
- 7 Cruz-Flores S, Rabinstein A, Biller J, *et al.* Racial-ethnic disparities in stroke care: the American experience: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke* 2011;42:2091–116.
- 8 Rinaldo L, Rabinstein AA, Cloft HJ, *et al.* Racial and economic disparities in the access to treatment of unruptured intracranial aneurysms are persistent problems. *J Neurointerv Surg* 2019;11:833–6.
- 9 Tabb LP, Ortiz A, Judd S, *et al.* Exploring the spatial patterning in racial differences in cardiovascular health between blacks and whites across the United States: the REGARDS study. *J Am Heart Assoc* 2020;9:e016556.
- 10 Ajinkya S, Almallouhi E, Turner N, *et al.* Racial/ethnic disparities in acute ischemic stroke treatment within a telestroke network. *Telemed J E Health* 2020;26:1221–5.
- 11 Goyal M, Menon BK, van Zwam WH, *et al.* Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *Lancet* 2016;387:1723–31.
- 12 Yancy CW. COVID-19 and African Americans. *JAMA* 2020;323:1891–2.
- 13 Reyes CHN, Gutowski C. *Chicago's coronavirus disparity: black Chicagoans are dying at nearly six times the rate of white residents, data show.* Chicago Tribune, 2020.
- 14 Schirmer CM, Ringer AJ, Arthur AS, *et al.* Delayed presentation of acute ischemic strokes during the COVID-19 crisis. *J Neurointerv Surg* 2020;12:639–42.
- 15 Al Kasab S, Almallouhi E, Alawieh A, *et al.* International experience of mechanical thrombectomy during the COVID-19 pandemic: insights from StAR and ENRG. *J Neurointerv Surg* 2020;12:1039–44.