



Consumption of Fortified Wheat Flour and Associations with Anemia and Low Serum Ferritin in Colombia.

Amy Fothergill, *Emory University.*

Zulma YF Centeno, *Universidad Nacional de Colombia, Observatorio de Soberanía y Seguridad Alimentaria y Nutricional-OBSSAN, Bogotá, Colombia.*

Paul René Ocampo Téllez, *Universidad Nacional de Colombia. Bogotá, Colombia.*

Helena Pachón, *Emory University*

Journal Title: Perspect Nutr Hum

Volume: Volume 21, Number 2

Publisher: PERSPECTIVAS EN NUTRICIÓN HUMANA | 2019, Pages 159-171

Type of Work: Article | Post-print: After Peer Review

Publisher DOI: 10.17533/udea.penh.v21n2a03

Permanent URL: <https://pid.emory.edu/ark:/25593/w0wd7>

Final published version: <http://dx.doi.org/10.17533/udea.penh.v21n2a03>

Copyright information:

This is an Open Access work distributed under the terms of the Creative Commons Attribution 4.0 International License (<https://creativecommons.org/licenses/by/4.0/rdf>).

Accessed November 10, 2024 12:43 PM EST



HHS Public Access

Author manuscript

Perspect Nutr Hum. Author manuscript; available in PMC 2021 September 15.

Published in final edited form as:

Perspect Nutr Hum. 2019 ; 21(2): 159–171. doi:10.17533/udea.penh.v21n2a03.

Consumption of Fortified Wheat Flour and Associations with Anemia and Low Serum Ferritin in Colombia

Amy Fothergill^{1,*}, Zulma Y. Fonseca Centeno², Paul René Ocampo Téllez³, Helena Pachón⁴

¹Emory University, Atlanta USA. Cornell University, Ithaca NY 14853, USA.

²Universidad Nacional de Colombia, Observatorio de Soberanía y Seguridad Alimentaria y Nutricional-OBSSAN, Bogotá, Colombia.

³Universidad Nacional de Colombia. Bogotá, Colombia.

⁴Emory University, Atlanta USA. Food Fortification Initiative, Atlanta, USA

Abstract

Background—Colombia’s mandatory wheat flour fortification program has yet to be evaluated.

Objective—Examine associations between consumption of fortified wheat flour and low serum ferritin (LSF) and anemia prevalence.

Materials and Methods—A secondary analysis of the 2005 national nutrition survey (ENSIN) was completed for 3988 children 2–4 y, 5669 children 5–12 y and 2053 non-pregnant women 13–49 y. The relationship between consumption (quartiles) of wheat flour containing food (WFCF) and LSF and anemia was examined using chi-square analyses and logistic regression models.

Results—In unadjusted analyses, the prevalence of LSF was similar across all quartiles of WFCF consumption in all age groups. The highest prevalence of anemia was observed in the lowest WFCF consumption quartiles in all age groups, but was not significantly different in non-pregnant women 13–49 y. In adjusted models this relationship between WFCF and anemia remained for children 2–4 y when comparing the highest WFCF intake quartile with the lowest quartile (OR: 0.7, 95 % CI: 0.6–0.9). No association between WFCF and LSF was observed in adjusted (or unadjusted) models.

Conclusions—In Colombia, consumption of wheat flour containing foods is associated with lower levels of anemia in pre-school children.

*Corresponding author. Phone: +1 603-630-0948. Fax: +1 607-254-6225. af544@cornell.edu.

AUTHORS CONTRIBUTIONS

H.P and A.F designed research; Z.Y.F.C. and P.R.O.T. helped secure the dataset; Z.Y.F.C. classified foods as containing or not containing wheat flour; A.F. analyzed data; and A.F and H.P wrote the paper. A.F had primary responsibility for final content. All authors read and approved the final manuscript.

CONFLICT OF INTEREST

None.

DISCLAIMER

The findings and conclusions of this report are solely the responsibility of the authors and do not necessarily represent the official position of the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD), the National Institutes of Health, or the US Centers for Disease Control and Prevention.

Resumen

la fortificación obligatoria de harina de trigo en Colombia no se ha evaluado.

examinar las asociaciones entre el consumo de harina y ferritina sérica baja (BSF) y la prevalencia de anemia.

se completó un análisis secundario de la Encuesta Nacional de la Situación Nutricional de 2005 (ENSIN) para 3988 niños de 2–4 años, 5669 escolares de 5–12 años y 2053 mujeres no embarazadas de 13–49 años. Se examinó la relación entre el consumo (cuantiles) de alimentos que contenían harina (ACHT) y la BSF y la anemia mediante análisis de ji al cuadrado y modelos de regresión logística.

en modelos ajustados, controlando por el estado socioeconómico, alimentos de origen animal y la ingesta de suplementos en las 24 horas anteriores, existía relación entre ACHT y la anemia en los preescolares. al comparar el cuartil más alto del consumo de ACHT con el cuartil más bajo (OR: 0.7; IC 95 %: 0,6–0,9). No se observó asociación entre ACHT y BFS en modelos ajustados.

en Colombia, el consumo de alimentos que contienen harina de trigo se asocia con niveles más bajos de anemia en preescolares.

Keywords

Ferritin; hemoglobin; fortification; wheat flour; enrichment; effectiveness; evaluation

Keywords

ferritina; hemoglobina; fortificación; harina de trigo; enriquecimiento; efectividad; evaluación

INTRODUCTION

Globally 85 countries have legislation mandating fortification of wheat flour, including the majority of countries in Latin America and the Caribbean. Countries with wheat flour fortification mandates typically fortify with at least iron and folic acid (1). A recent literature review (2) examining the effectiveness of flour fortification found that anemia was most likely to be reduced if a World Health Organization (WHO) recommended iron compound was used in fortification (3). Overall, reductions in anemia were only observed in one third of the groups studied, and reductions in the prevalence of low ferritin were only consistently observed in women, not children (2). Compared to the number of countries that currently fortify, few effectiveness evaluations were available. The Colombian standard for wheat flour fortification allows mills to choose among three iron fortificants (ferrous fumarate, ferrous sulfate, and reduced iron) (4); the first two are WHO recommended while the third is not (3). If a non-recommended form of iron is used to fortify, positive nutritional effects may not be seen at the national level (2,5,6).

In 1996 Colombia mandated fortification of wheat flour with vitamins B1, B2, B3, B9 (folic acid), and iron, while calcium is optional (4). A clarification released in 2013 explicitly states that all domestically produced and imported foods with wheat flour as the main

ingredient must use fortified wheat flour (7). To date, no impact evaluations of the country's wheat flour fortification program have been conducted. This study examines the relationship between consumption of wheat flour during the wheat flour fortification program in Colombia and low serum ferritin and anemia status, using nationally representative data from 2005.

MATERIALS AND METHODS

The 2005 National Nutrition Situation Survey (ENSIN), was conducted in a subset of the households targeted by the 2005 Demographic and Health Survey (DHS, n=37,211) and included 17,740 households (8,9). The DHS survey used probabilistic sampling to ensure it represented 98 % of the national population (8), and the ENSIN survey was representative at the national, urban/rural, regional, and sub-regional levels, and included people aged 0–64 y. Individuals aged 2–64 y participated in a 24 h dietary recall survey and a subset of those completing the survey had blood samples collected. The household selection process and survey administration methods are described else-where (8). Briefly, within an identified household, every person 2–20 y and every 3rd person 21–64 y participated in the 24 h dietary recall; every 4th woman 13–49 y, every other child 5–12 y, and every child 1–4 y gave a blood sample (9).

Study population

Children 2–12 y and non-pregnant women 13–49 y who completed the 24 h dietary recall and for whom hemoglobin, serum ferritin and C-reactive protein (CRP) levels were measured were included (Figure 1). Descriptive statistics were calculated for the entire study population with dietary and biological data to examine the general characteristics of individuals present in the dataset (Table 1).

Does consumption of wheat flour containing foods (WFCF) influence iron status and/or anemia status?—This analysis aimed to examine low serum ferritin and anemia status and their relation to consumption of fortified wheat flour. However, the composition of WFCF, specifically the percent composed of wheat flour, was unavailable. The consumption of WFCF was therefore used as a proxy for wheat flour consumption. All unique foods reported in the dietary recall (n=1027) were grouped into broad food types (n=310) and also classified as either containing (n=32) or not containing wheat flour (n=281). Bienestarina is a food that contains fortified wheat flour (9). It is produced and distributed to recipients of the Family Welfare program in Colombia. Bienestarina was commonly reported in the 24 h recall and was categorized as a WFCF. Some of the broad food groups classified as not containing wheat flour had foods within them that did contain wheat flour; these foods were classified individually as containing wheat flour or not.

Additionally, some foods were classified as being an animal source food, and micronutrient supplement intake was documented, although the micronutrient formulation was unknown.

Biomarker outcomes were limited to iron status and hemoglobin. Iron is the only nutrient included in the fortification program that was measured in biological samples (serum ferritin). Hemoglobin levels can be influenced by multiple nutrients added to fortified

wheat flour: iron, and B-vitamins such as folate, B1, B2, and B3 (10). The main outcomes of interest were iron status (as determined through serum ferritin) and anemia status (as determined through altitude-adjusted hemoglobin levels (11)).

Laboratory Analyses

Hemoglobin concentrations were determined in real-time via HemoCue, which was calibrated daily and after every 20 measurements (9). All venous samples were processed in the field and frozen in liquid nitrogen until they were sent to the National Health Institute's Nutritional Laboratory for analysis (9). Plasma ferritin concentrations were measured using the ADVIA Centaur chemiluminescence assay (9). Plasma CRP was measured with a turbidometric method (9).

Statistical Analyses

Linear regression models were used to estimate the effect of WFCF consumption on biomarker levels as continuous variables (serum ferritin, hemoglobin), and logistic regression models to estimate the effect of WFCF consumption on biomarker levels as categorical variables (low serum ferritin (LSF), anemia). For categorical analyses, biomarker levels were examined as binary outcomes (LSF vs. not LSF). Consistent with how the ENSIN analyzed the biomarker data in 2005 (8), individuals were included in low serum ferritin analyses if their CRP levels were <1.2 mg/dL; individuals were considered to have low serum ferritin if they had eligible CRP levels and their serum ferritin was <12 µg/L. Also consistent with the 2005 ENSIN analyses, hemoglobin levels were adjusted for altitude using the Dirren et al. method (11), and adjusted hemoglobin levels were used in conjunction with age to determine anemia status. Smoking status was unavailable, therefore hemoglobin levels were unable to be adjusted for smoking. Children 2–4 y were considered anemic if their adjusted hemoglobin was <11 g/dL, children 5–12 y and non-pregnant women 13–49 y were considered anemic if their hemoglobin was <12 g/dL. When looking at the effects of WFCF consumption on iron and anemia status, all analyses were conducted within age groups: 2–4 y, 5–12 y and non-pregnant women 13–49 y.

All analyses were conducted in SAS (version 9.3; SAS Institute) using procedures for complex survey design utilizing clusters, strata, and statistical weights to account for differing response rates. All analyses were conducted with and without the weights to see if the outcomes were affected. No differences were found in the outcomes, and therefore weighted counts and models are reported. Methods proposed by Rothman and Greenland (12) were used to adjust all models for confounding. El Sistema de Selección de Beneficiarios para Programas Sociales (SISBEN) level (a measure of socioeconomic status (13)), having consumed any animal source food in the previous 24 h, and having consumed any micronutrient supplements in the previous 24 h were included in the final adjusted models.

SISBEN levels were available for every household participating in the ENSIN survey (9). SISBEN scores are assigned to households based on a variety of factors and used to provide the Colombian government with a ranking system to help identify households in need of

social welfare programs. Households with scores of 1–2 are in most need, while households with a score of 6 are considered to have the least need (13).

Ethics Considerations

The Instituto Colombiano de Bienestar Familiar (Colombian Institute of Family Wellbeing) and the Instituto Nacional de Salud (INS; National Institute of Health) coordinated the collection and analysis of blood samples; ethical approval was granted by the INS. Written consent was obtained from the adult interviewee or parents of children for the blood draws. The analysis presented here is a secondary analysis of a large, nationally representative dataset. For this analysis, data were provided in a de-identified format; thus, no additional ethical review was sought.

RESULTS

The main characteristics of the women and children from the 2005 ENSIN included in this analysis are summarized in Table 1. The majority of the study population had consumed both WFGF and animal source foods in the previous 24 h. The most commonly consumed WFCF were bread, pasta, cookies, and Bienestarina while the most commonly consumed animal source foods were milk, eggs, and beef. Approximately 13 % of children 2–4 y had taken a supplement in the previous 24 h, while less than 8 % of children 5–12 y and non-pregnant women 13–49 y had taken a supplement (Table 1). Despite being highly prevalent in the population, WFCF consumption varied significantly between socioeconomic strata in all age groups (data not shown, see discussion). The median intake of WFCF ranged from 58.9 g/d in children 2–4 y to 77.3 g/d in children 5–12 y.

Low serum ferritin was prevalent in 26.4 %, 11.9 % and 20.3 % of children 2–4 y, children 5–12 y and non-pregnant women 13–49 y, respectively (Table 1). Anemia was 26.3 %, 37.3 %, and 33.7 % for children 2–4 y, children 5–12 y and non-pregnant women 13–49 y, respectively. The prevalence of low serum ferritin anemia (LSF-A) was <12 % in all age groups. The prevalence of LSF was similar across all levels of WFCF consumption quartiles for all age groups (Figure 2). The highest prevalence of anemia was observed in the lowest WFCF consumption quartiles for all age groups, and the prevalence of anemia was found to be statistically different within age group by quartile of consumption in all age groups except non-pregnant women 13–49 y (Figure 3).

In logistic regression models controlling for SISBEN, having consumed animal source foods and supplements in the previous 24 h, the observed relationship between WFCF and low serum ferritin was statistically significant when comparing quartile 3 vs. 1 in non-pregnant women 13–49 y, but not for any other age group (Table 2). The relationship between WFCF consumption and anemia was statistically significant for children 2–4 y when comparing quartile 2 vs. 1 and quartile 4 vs. 1, but not for any other age group (Table 3). When serum ferritin and hemoglobin were examined as continuous variables, relationships between hemoglobin and continuous WFCF consumption were statistically significant for all age groups in adjusted models, with increased WFCF consumption associated with increased hemoglobin concentrations. Relationships between serum ferritin and continuous WFCF consumption were not statistically significant for any age group.

DISCUSSION

In Colombia, consuming WFCF was associated with a lower prevalence of anemia in children 2–4 y, with the highest prevalence of anemia being observed in the lowest quartiles of WFCF consumption for this age group.

In this evaluation, we were able to estimate individual intake of WFCF rather than household consumption, but our estimates were limited by our knowledge of the flour content in WFCFs consumed.

For example, foods such as cookies and sandwiches were coded as containing wheat flour, because they were likely made with fortified wheat flour. However, we did not know the proportion of wheat flour in each food, so the entire weight of the cookie/sandwich contributed to the grams of WFCF consumed. We completed analyses for all WFCF and those WFCF where more than 50 % of the weight was estimated by us to contain flour; the results did not differ. Additionally, we were unable to calculate iron intake from WFCF. To partially address this, we controlled for consumption of foods known to contain high amounts of iron in all analyses. Knowing the wheat flour composition of all foods consumed and the contributions of each food to total iron intake would allow for a more accurate assessment of the fortified flour exposure.

Despite these limitations, this study is different from other impact evaluation studies because rather than comparing pre/post fortification levels (or changes therein) of low serum ferritin and anemia, (14–16) we looked at individual consumption levels and relationships with health outcomes. Grimm et al. also looked at consumption levels and health outcomes in Oman, but they examined per capita household flour consumption. (17) They found a lower prevalence of iron deficiency in non pregnant women from households with 1 kg of monthly wheat flour consumption compared to women from households with <1 kg per capita consumption per month.

Other evaluation studies included information on whether the flour was adequately fortified based on government monitoring reports (18) or from sampling and testing flour samples (19–21). In this study, such information was not available for 2005. Regulatory monitoring data from 2011, 2012–13, and 2015 show that 17.5 %, 7.4 % and 7.0 % of flour samples, respectively, did not meet standards (Claudia Patricia Moreno Barrera, Ministers de Salud y Protección Social, personal communication, 12 September, 2016), suggesting that during the 2011–15 period, most samples met the country's flour fortification standards. The type and amount of fortificant being used to fortify wheat flour can differ from the country standard; knowing what was used in Colombia would strengthen the analyses reported here. Imported wheat flour products are supposed to be produced with fortified wheat flour, according to the 2013 government clarification (7). In some small stores it was recently noted (anecdotally) that imported WFCF such as pasta were not all labeled as using fortified wheat flour. A better understanding of the compliance of import regulations in Colombia would positively contribute to this evaluation.

In 2009, WHO released recommendations for wheat flour fortification with iron and four other nutrients (3). Colombia was fortifying its flour prior to the release of these

recommendations (4). Using non-recommended forms of iron, or less than the recommended amount, may reduce the effectiveness of wheat flour fortification (2,5,6). After Costa Rica changed and intensified the iron component of its food fortification program, researchers found reductions in iron deficiency, anemia and iron-deficiency anemia in children 1–7 y and reductions in anemia in women 14–45 y (18). In 1958 Costa Rica mandated fortification of wheat flour with 45 mg/kg of iron as reduced iron (not a WHO recommended fortificant), but switched in 2002 to 55 mg/kg of iron as ferrous fumarate (a WHO recommended, more bioavailable fortificant).

WHO fortification recommendations are based on wheat flour consumption or availability (3). The median intake for WFCF in Colombia observed in this analysis (58.9–77.3 g/d) puts it in the lowest tier of WHO recommendations (<75 g/d). For this tier, the recommended iron levels for wheat flour are 40 mg/kg as ferric sodium ethylenediaminetetraacetate (NaFeEDTA), 60 mg/kg as ferrous sulfate, or 60 mg/kg as ferrous fumarate. At consumption levels this low, electrolytic iron is not recommended. Currently the Colombia requirement of 44 mg/kg of iron (4) is below the WHO recommendations for ferrous sulfate and ferrous fumarate (3). One of the allowable fortificants in Colombia, reduced iron, is an ineffective fortificant (5). Reduced iron is the least expensive among the three compounds allowed (22), but it is unclear how often it is utilized by flour mills. Increasing the requirement to 60 mg/kg and eliminating the option of using reduced iron, might improve the iron status across the population.

In this study population, low serum ferritin contributed to approximately 1/3 of the anemia in children 2–4 y and non-pregnant women 13–49 y, while contributing to a smaller portion in children 5–12 y (~13 %). The anemia in Colombia may be caused by other nutritional deficiencies as well as non-nutritional causes. Given the limitations with the iron compounds and levels used to fortify wheat flour, it is plausible that the other nutrients in fortified wheat flour (vitamins B1, B2, B3, and B9) are contributing to improvements in hemoglobin synthesis. This could be contributing to the associations between WFCF consumption and hemoglobin in all age groups and between WFCF consumption and anemia in children 2–4 y. In a recent analysis among children in Colombia, micronutrient powders (MNP) were not found to improve hemoglobin status. The authors hypothesize that this could be due to the lower prevalence of iron deficiency anemia (23) which is in line with our observations here.

WFCF consumption differed significantly between certain groups, with urban dwellers and people in higher SISBEN levels more likely to report WFCF consumption. In models looking at WFCF and anemia, WFCF was associated with anemia in 5–12 y children until SISBEN was added (and SISBEN was significantly related to anemia on its own). If accessibility and cost of WFCFs are preventing people in more rural areas and lower SISBEN levels from consuming WFCF, anemia interventions (including the fortification of alternative food products) that target rural populations and lower SISBEN levels may be beneficial.

WHO recommends adjusting women's hemoglobin data for smoking and lack of these data in this analysis is a limitation (24). Additionally, the current analyses utilized cutoffs for low serum ferritin and anemia that were originally used in the ENSIN report, rather than those

recommended by WHO (24,25). This decision was made to remain consistent with reports previously published by the Colombian government. This limits the ability to compare these results to those from other countries that use the WHO cutoffs.

Despite not having information on percent composition of foods, food-specific iron contributions, and fortificants currently in use in Colombia and compliance with standards, this study has several strengths. The sample size is very large and nationally representative of the Colombian population and we were able to control for CRP in the low serum ferritin analyses.

In conclusion, the consumption of WFCF was associated with lower levels of anemia but not low serum ferritin in Colombia, in specific age groups. In adjusted analyses, among children between the ages of 2–4 y, lower anemia prevalence was observed among groups with a higher consumption of wheat flour containing foods. This analysis utilized a large, nationally representative sample with biomarker and food recall data collected in close time proximity. Information on iron fortificants used in Colombia, fortification compliance in the country, and percent composition of wheat flour containing food could strengthen further evaluations of Colombia's wheat flour fortification program.

ACKNOWLEDGMENTS

The authors would like to thank Dr. Yaw Addo, Dr. Penelope Howards, Dr. Katarzyna Kordas, Ms. Sayda Pico, Ms. Sarah Zimmerman, Dr. Melanie Jacobson and Dr. Jennifer Richards for their help with this manuscript.

Funding for this research was provided solely by way of salary support. Research reported in this publication was supported by the National Institutes of Health under award 5 T32 HD087137 (Amy Fothergill). Helena Pachón's time was supported by an agreement between the United States Centers for Disease Control and Prevention (CDC), McKing Consulting Corporation, and Emory University.

References

1. Food Fortification Initiative. Global Progress of Industrially Milled Cereal Grains. [Internet], [Cited October 2018], Available from: http://ffinetwork.org/global_progress/index.php
2. Pachón H, Spohrer R, Mei Z, & Serdula MK. Evidence of the effectiveness of flour fortification programs on iron status and anemia: a systematic review. *Nutr Rev.* 2015;73(11):780–95. DOI: 10.1093/nutrit/nuv037 [PubMed: 26433017]
3. WHO, FAO, UNICEF, GAIN, MI, & FFI. Recommendations on wheat and maize flour fortification. Meeting Report: Interim Consensus Statement, Geneva, World Health Organization, 2009. [Internet], [Cited October 2018], Available from: www.who.int/nutrition/publications/micronutrients/wheat_maize_fort.pdf
4. Forero de Saade TPE. (1996). Decreto Número 1994 de 1996. [Internet], [Cited October 2018], Available from: https://www.invima.gov.co/images/stories/aliementos/decreto_1944_1996.pdf
5. Hurrell R, Ranum P, De Pee S, Biebinger R, Hulthen L, Johnson Q, et al. Revised recommendations for iron fortification of wheat flour and an evaluation of the expected impact of Current national wheat flour fortification programs. *Food and Nutr Bull.* 2010;31 (suppl1):S7–S21. DOI: 10.1177/15648265100311S102 [PubMed: 20629349]
6. Hurrell RF. Flour fortification as a strategy to prevent anaemia. *Brit J Nutr.* 2015; 114(04),501–2. DOI: 10.1017/S0007114515002147 [PubMed: 26198077]
7. República de Colombia, Instituto Nacional de Vigilancia de Medicamentos y Alimentos INVIMA. Procedimiento Para Importacion de Productos Cuyo Ingrediente Principal es la Harina de Trigo 2013. [Internet], [Cited October 2018], Disponible en: <https://paginaweb.invima.gov.co/informacion-de>

intres/alimentos-y-bebidas-info/4378-procedimiento-para-importaci%C3%B3n-de-productos-cuyo-
ingrediente-principal-es-la-harina-de-trigo.html

8. Ojeda G, Ordóñez M, Ochoa LH. Salud sexual y reproductiva. Resultados Encuesta Nacional de Demografía y Salud. Profamilia, ICBF, USAID, el Ministerio de la Protección Social MPS, UNFPA. 2005. Bogotá: Disponible en: <https://www.minsalud.gov.co/sites/rid/Lists/BibliotecaDigital/RIDE/VS/ED/GCFI/Base%20de%20datos%20ENDS%202005%20informe.pdf>
9. Profamilia, Ministerio de la Protección Social, ICBF. Encuesta nacional de la situación nutricional en Colombia ENSIN 2005. Bogotá: Instituto Colombiano de Bienestar Familiar; 2006. 440 pp. Disponible en: https://www.icbf.gov.co/sites/default/files/libro_2005.pdf
10. Kraemer K, Zimmermann MB. Nutritional Anemia. Base: Sight and Life Press Basel; 2007. 400 pp. Available from: http://ernaehrungsdenkwerkstatt.de/fileadmin/user_upload/EDWText/TextElemente/Ernaehrungswissenschaft/Naehrstoffe/nutritional_anemia_book.pdf
11. Dirren H, Logman MH, Barclay DV, Freire WB. Altitude correction for hemoglobin. *Eur J Clin Nutr.* 1994;48(9):625–32. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/8001519> [PubMed: 8001519]
12. Rothman KJ, Greenland S, Lash T. *Modern Epidemiology*. 3rd ed. Philadelphia (PA): Lippincott Williams & Williams. 2005. 56–60 pp.
13. Velez CE, Castaño E, Deutsch R. An Economic Interpretation of Colombia's SISBEN: A Composite Welfare Index Derived from the Optimal Scaling Algorithm An Economic Interpretation of Colombia's SISBEN: A Composite Household Welfare Index Derived from the Optimal Scaling Algorithm. 1998. Available from: <http://idbdocs.iadb.org/wsdocs/getdocument.aspx?docnum=364066>
14. Tazhibayev S, Dolmatova O, Ganiyeva G, Khairov K, Ospanova F, Oyunchimeg D, et al. Evaluation of the potential effectiveness of wheat flour and salt fortification programs in five Central Asian countries and Mongolia. *Food and Nutr Bull.* 2002;29(4):255–65. DOI: 10.1177/156482650802900402
15. Assunção MC, Santos IS, Barros AJ, Gigante DP, Victora CG. Flour fortification with iron has no impact on anaemia in urban Brazilian children. *Public Health Nutr.* 2012;15(10):1796–801. DOI: 10.1017/S1368980012003047 [PubMed: 22704130]
16. Sadighi J, Mohammad K, Sheikholeslam R, Amirkhani MA, Torabi P, Salehi F, et al. Anaemia control: Lessons from the flour fortification programme. *Public Health.* 2009;123(12):794–9. DOI:10.1016/J.PUHE.2009.09.024 [PubMed: 19914671]
17. Grimm KA, Sullivan KM, Alasfoor D, Parvanta I, Mohammed-Suleiman AJ, Kaur M, et al. Iron-fortified wheat flour and iron deficiency among women. *Food and Nutr Bull.* 2012;33(3): 180–5. DOI: 10.1177/156482651203300302 [PubMed: 23156120]
18. Martorell R, Ascencio M, Tacsan L, Alfaro T, Young M F, Addo OY, et al. Effectiveness evaluation of the food fortification program of Costa Rica: impact on anemia prevalence and hemoglobin concentrations in women and children. *Am J Clin Nutr.* 2015;101(1):210–7. DOI: 10.3945/ajcn.114.097709 [PubMed: 25527765]
19. Layrisse M, Garcia-Casal MN, Mendez-Castellano H, Jimenez M, Olavarria C H, Chavez J, et al. Impact of fortification of flours with iron to reduce the prevalence of anemia and iron deficiency among schoolchildren in Caracas, Venezuela: A followup. *Food and Nutr Bull.* 2002;23(4) 384–9 DOI: 10.1177/156482650202300412 [PubMed: 16619746]
20. Hund L, Northrop-Clewes C, Nazario R, et al. A novel approach to evaluating the iron and folate status of women of reproductive age in Uzbekistan after 3 years of flour fortification with micronutrients. *PLOS ONE.* 2013; 8 (11):1–11. DOI: 10.1371/journal.pone.0079726
21. Schultz J, Vatucawaqa P. Impact of Iron Fortified Flour in Child Bearing Age (CBA) Women in Fiji. 2010 Report. Fiji: National Food and Nutrition Centre, UNICEF, WHO, Ministry of health Shaping Fiji's Health; 2012, 61 pp. Available from: <http://www.ffinetwork.org/monitor/Documents/Fiji.pdf>
22. Lofti M, Venkatesh MG, Merx RJHM, Naber-van den Heuvel P. Micronutrient Fortification of Foods Current practices, research, and opportunities. Ontario, Canada: The Micronutrient Initiative, International Development Research Centre, International Agricultural Centre. 1996. 109 pp. Available from: <https://idl-bnc-idrc.dspacedirect.org/bitstream/handle/10625/14002/104386.pdf?sequence=1>

23. Andrew A, Attanasio O, Fitzsimons E, Rubio-Codina M. Why is multiple micronutrient powder ineffective at reducing anaemia among 12–24 month olds in Colombia? Evidence from a randomised controlled trial. *SSM - Popul Health*. 2016;2:95–104. DOI: 10.1016/J.SSMPH.2016.02.004 [PubMed: 29349132]
24. WHO. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. Vitamin and Mineral Nutrition Information System. Geneva: WHO; 2011. 5 pp. Available from: https://apps.who.int/iris/bitstream/handle/10665/85839/WHO_NMH_NHD_MNM_11.1_eng.pdf?ua=1
25. WHO. Serum ferritin concentrations for the assessment of iron status and iron deficiency in populations. Vitamin and Mineral Nutrition Information System. Geneva: WHO; 2011.6 pp. Available from: http://www.who.int/vmnis/indicators/serum_ferritin.pdf




Figure 1. Population flow from Colombia Demographic Health Survey (DHS), through Encuesta Nacional de Situación Nutricional (ENSIN) and included here.




Figure 2. Quartiles of wheat flour containing food consumption, and the unadjusted relationship with low serum ferritin.

Low serum ferritin was defined as having serum ferritin levels $<12 \mu\text{g/L}$ and CRP levels $<1.2 \text{ mg/dL}$. All participants with CRP levels $>1.2 \text{ mg/dL}$ were removed from analyses per ENSIN 2005 analytical procedures (8). The sample size per group by quartile: Children 2–4 y (975,997,1029, and 1029), Children 5–12 y (1382,1409,1399,1485), and Non-pregnant women, 13–49 y (620,426,482,531). Estimated prevalence based on weighted counts for low serum ferritin within each quartile of consumption.




Figure 3. Quartiles of wheat flour containing food consumption, and the unadjusted relationship with anemia.

Anemia was defined as having low hemoglobin levels (<11 g/dL for children 2–4 y and <12 g/dL for children 5–12 y and non-pregnant women 13–49 y), after adjusting for altitude using the Dirren et al. method (11). The sample size per group by quartile: Children 2–4 y (975,997, 1029, and 1029), Children 5–12 y (1382,1409,1399,1485), and Non-pregnant women, 13–49 y (620,426,482,531). Estimated prevalence based on weighted counts for anemia within each quartile of consumption. * Statistically significant; Chi-square $p < 0.01$.

Table 1. Overall population characteristics of women and children in the 2005 ENSIN survey weighted for survey

	Children 2–4 y	Children 5–12 y	Non-pregnant women 13–49 y
Total N [*] (both dietary and biomarker data)	3988	5669	2053
Weighted population N [†] (both dietary and biomarker data)	4030	5676	2059
Consumption in previous 24 h [‡]			
Wheat flour containing food, n (%)	3254 (80.7)	4489 (79.0)	1441 (69.9)
Animal source food, n (%)	3874 (96.1)	5488 (96.7)	1991 (96.7)
Micronutrient supplement, n (%)	533 (13.2)	423 (7.5)	137 (6.7)
Grams WFCF consumed, median (95 % CI)	58.9 (56.4, 61.5)	77.3 (72.8, 81.8)	70.9 (65.0, 76.9)
Serum ferritin in µg/L, median (95 % CI)	26.7 (25.9, 27.5)	32.8 (31.9, 33.7)	27.7 (26.4, 28.9)
Hemoglobin in g/dL, median (95 % CI)	11.6 (11.5, 11.6)	12.3 (12.2, 12.4)	12.4 (12.3, 12.5)
C-reactive protein in mg/dL, median (95 % CI)	0.1 (0.09, 0.1)	0.09 (0.09, 0.1)	0.1 (0.09, 0.1)
Prevalence of low serum ferritin [§] , n (%)	1010 (26.4)	658 (11.9)	401 (20.3)
Prevalence of anemia , n (%)	1059 (26.3)	2119 (37.3)	696 (33.7)
Prevalence of low serum ferritin anemia [§] , n (%)	338 (8.4)	274 (4.8)	232 (11.3)
Sex, n (%)			
Female	1944 (48.2)	2823 (49.7)	2059 (100)
Male	2086 (51.8)	2853 (50.3)	0 (0)
Geographic location, n (%)			
Urban	2849 (70.7)	4014 (70.7)	1616 (78.5)
Rural	1181 (29.3)	1662 (29.3)	443 (21.5)
SISBEN [¶] , n (%)			
1	1577 (39.1)	2246 (39.6)	690 (33.5)
2	1505 (37.3)	2144 (37.8)	817 (39.7)
3	862 (21.4)	1178 (20.8)	495 (24)
4–6	86 (2.1)	108 (1.9)	57 (2.8)
Altitude ^{**} in MASL, n (%)			
<200	1638 (40.7)	2444 (43.1)	903 (43.9)

	Children 2–4 y	Children 5–12 y	Non-pregnant women 13–49 y
>200	2392 (59.3)	3232 (56.9)	1156 (56.1)

WFCE- Wheat flour containing food

* The total study population had dietary data and both serum ferritin and hemoglobin values. All individuals who had serum ferritin values also had values for C-reactive protein (CRP).

[†]Weighted analyses were conducted to account for the survey design. All reported counts and percentages in the table are based on the weighted populations.

[‡]All consumption was self-reported. Items were categorized as a wheat flour containing food, an animal source food, or micronutrient supplements.

[§]Low serum ferritin was defined as having serum ferritin levels <12 µg/L and CRP levels <1.2 mg/dL. All participants with CRP levels 1.2 mg/dL were removed from analyses per ENSIN 2005 analytical procedures (8).

// Anemia was defined as having low hemoglobin levels (<11 g/dL for children 2–4 y and <12 g/dL for children 5–12 y and non-pregnant women 13–49 y), after adjusting for altitude using the Dirren et al. method (11).

[¶]An individual's SISBEN (El Sistema de Selección de Beneficiarios para Programas Sociales) score is a measure of their socioeconomic status, with higher numbers corresponding to higher status (12).

^{**}The Dirren et al. method adjusts hemoglobin for all altitudes > 200 meters above sea level (MASL) (10).

Table 2. Association between wheat flour containing food (WFCF) consumption and being classified as low serum ferritin

WFCF consumption	Association with low serum ferritin*					
	n	Unadjusted			Adjusted [†]	
		OR	95% CI	OR	95% CI	OR
Children 2–4 y						
Quartile 1 (13.3 g)	975	1.00	Reference	1.00	Reference	
Quartile 2 (13.4–43 g)	997	0.77	0.6–1.0	0.76	0.6–0.9	
Quartile 3 (43.1–89 g)	1029	0.86	0.7–1.1	0.84	0.7–1.1	
Quartile 4 (90 g)	1029	0.88	0.7–1.1	0.85	0.7–1.1	
Children 5–12 y						
Quartile 1 (14.25 g)	1382	1.00	Reference	1.00	Reference	
Quartile 2 (14.26–56 g)	1409	0.83	0.7–1.0	0.81	0.6–1.0	
Quartile 3 (56.1–112 g)	1399	1.0	0.8–1.3	0.99	0.8–1.3	
Quartile 4 (112.1 g)	1485	0.96	0.7–1.2	0.91	0.7–1.2	
Non-pregnant women 13–49 y						
Quartile 1 (0 g)	619	1.00	Reference	1.00	Reference	
Quartile 2 (0.1–40 g)	426	1.13	0.8–1.6	1.12	0.8–1.6	
Quartile 3 (40.1–97 g)	482	1.38	1.1–1.8	1.39	1.1–1.8	
Quartile 4 (97.1 g)	532	1.18	0.9–1.5	1.16	0.9–1.5	

WFCF- Wheat flour containing food

* Low serum ferritin was defined as having serum ferritin levels <12 µg/L and CRP levels <1.2 mg/dL. All participants with CRP levels 1.2 mg/dL were removed from analyses per ENSIN 2005 analytical procedures (8).

[†] Adjusted models controlled for SISBEN level (12), having taken a micronutrient supplement and having consumed an animal source food in previous 24 h.

Table 3. Association between wheat flour containing food (WFCF) consumption and being classified as anemic*

WFCF consumption	Unadjusted			Adjusted [†]		
	n	OR	95% CI	OR	95% CI	Reference
Children 2–4 y						
Quartile 1 (13.3 g)	975	1.00	Reference	1.00	Reference	
Quartile 2 (13.4–43 g)	997	0.66	0.5, 0.8	0.67	0.5, 0.8	
Quartile 3 (43.1–89 g)	1029	0.85	0.7, 1.0	0.89	0.7, 1.1	
Quartile 4 (90 g)	1029	0.66	0.5, 0.8	0.7	0.6, 0.9	
Children 5–12 y						
Quartile 1 (14.25 g)	1382	1.00	Reference	1.00	Reference	
Quartile 2 (14.26–56 g)	1409	0.89	0.8, 1.0	0.97	0.8, 1.1	
Quartile 3 (56.1–112 g)	1399	0.97	0.8, 1.2	1.1	0.9, 1.3	
Quartile 4 (112.1 g)	1485	0.75	0.6, 0.9	0.9	0.7, 1.1	
Non-pregnant women 13–49 y						
Quartile 1 (0 g)	619	1.00	Reference	1.00	Reference	
Quartile 2 (0.1–40 g)	426	0.99	0.8, 1.2	1.05	0.8, 1.3	
Quartile 3 (40.1–97 g)	482	0.95	0.7, 1.2	0.99	0.8, 1.3	
Quartile 4 (97.1 g)	532	0.88	0.7, 1.2	0.91	0.7, 1.2	

WFCF- Wheat flour containing food

* Anemia was defined as having low hemoglobin levels (<11 g/dL for children 2–4 y and <12 g/dL for children 5–12 y and non-pregnant women 13–49 y), after adjusting for altitude using the Dirren et al. method (11).

[†] Adjusted models controlled for SISBEN level (12), and having taken a micronutrient supplement and having consumed an animal source food in previous 24 h.